



FCT Fundação para a Ciência e a Tecnologia
MINISTÉRIO DA EDUCAÇÃO E CIÊNCIA



Joana Vanessa Cordeiro Melro Mourão

**CONTRIBUTION OF BIOCIDES TO THE SELECTION OF
SALMONELLA NON-TYPHOID RESISTANT TO
ANTIBIOTICS: A MULTILAYERED APPROACH**

*Tese do 3.º Ciclo de Estudos Conducente ao Grau de Doutoramento em Ciências
Farmacêuticas na Especialidade de Microbiologia*

Trabalho realizado sob a orientação da Professora Doutora Patrícia Antunes (Faculdade de Ciências da Nutrição e Alimentação, Universidade do Porto, Portugal) e coorientação da Professora Doutora Luísa Peixe e Professora Doutora Carla Novais (Faculdade de Farmácia, Universidade do Porto, Portugal).

February 2016

É AUTORIZADA A REPRODUÇÃO INTEGRAL DESTA TESE APENAS PARA EFEITOS DE INVESTIGAÇÃO, MEDIANTE DECLARAÇÃO ESCRITA DO INTERESSADO, QUE A TAL SE COMPROMETE;

Joana Vanessa Cordeiro Melro Mourão

Joana Vanessa Cordeiro Melro Mourão

To my parents for teaching me to never give up

AGRADECIMENTOS (ACKNOWLEDGMENTS)

Ao concluir esta Tese de Doutoramento quero deixar expresso o meu reconhecimento a instituições, e acima de tudo, a todas as pessoas que me acompanharam pelo mundo da Ciência por todo o auxílio que me prestaram e que, de uma forma decisiva, contribuíram para que a realização desta tese fosse possível. **MUITO OBRIGADA!**

Em primeiro lugar à minha orientadora Prof. Doutora Patrícia Antunes, e coorientadoras Prof. Doutora Luísa Peixe e Prof. Doutora Carla Novais quero deixar expresso o meu profundo agradecimento por terem aceite esta tarefa tão árdua que é a orientação de um aluno de Doutoramento. Agradeço todo o apoio durante estes quatro anos e a partilha de conhecimentos constante.

À Prof. Doutora Patrícia Antunes e à Prof. Doutora Carla Novais, não quero deixar passar esta oportunidade para expressar um agradecimento muito especial. Sem dúvida que a conclusão desta tese não teria sido possível sem o vosso esforço e contributos inigualáveis. Muito obrigada do fundo do coração pela paciência, pelo estímulo diário, pelos conhecimentos transmitidos, pelas opiniões e críticas, pelas longas discussões, pela constante disponibilidade e por fomentarem o meu espírito crítico e independência todos os dias. Sou sem dúvida uma pessoa muito diferente graças ao vosso contributo valioso.

Em segundo lugar quero agradecer todo o apoio financeiro concedido ao longo destes quatro anos: Fundação para a Ciência e a Tecnologia (FCT), Ministério da Ciência, Tecnologia e Ensino Superior de Portugal (Bolsa SFRH/BD/77518/2011, QREN – POPH – Tipologia 4.1 – Formação Avançada, participado pelo Fundo Social Europeu e por fundos sociais do MEC). Adicionalmente agradeço também à Unidade de Investigação/FCT – UCIBIO/REQUIMTE (através do projeto PEst-C/EQB/LA0006/2011 e UID/MULTI/04378/2013).

Ao Dr. Jorge Machado do Instituto Nacional de Saúde Dr. Ricardo Jorge de Lisboa, muito obrigada pela disponibilidade constante.

À Prof. Doutora São José Nascimento agradeço ter-me dado a oportunidade de realizar este trabalho no Laboratório de Microbiologia da Faculdade de Farmácia da Universidade do Porto. A todos os Prof. Doutores do Laboratório de Microbiologia quero agradecer todas as palavras de carinho, disponibilidade e a simpatia com que me receberam.

À Cristina Pinto da Costa um agradecimento muito especial, por toda a ajudar que me deste, pelos desabafos, pela companhia e acima de tudo pela amizade que vou guardar para sempre no meu coração. Muito obrigada por tudo. Ao Nuno Oliveira agradeço todo o apoio, risadas e incentivo ao longo destes anos.

Aos companheiros assíduos e confidentes de longa data João Pires, Joana Campos, Eduarda Silveira e Carla Rodrigues quero agradecer todos os momentos de confidências, companheirismo, amizade, risadas, partilha e ajuda nos bons e nos maus momentos. Sem dúvida que tornaram este caminho muito mais fácil de ultrapassar. Muito obrigada por estarem sempre presentes.

Aos membros e companheiros mais recentes, mas igualmente importantes João Botelho, Liliana Silva e Teresa Ribeiro, o meu muito obrigado pelo companheirismo, carinho, simpatia e amizade.

À Ângela Novais, Ana Freitas e Filipa Grosso quero agradecer todo a disponibilidade e partilha de conhecimentos.

A todos os alunos de projeto e de iniciação à investigação da Faculdade de Farmácia e da Faculdade de Ciências da Nutrição e Alimentação da Universidade do Porto (Laura, Liliana, Paula, Sara) muito obrigado pelo carinho, simpatia e por toda a ajuda.

A todos os alunos de ERASMUS (António, Leonardo, Jennifer, Houyem e Aymen) muito obrigada por todos os momentos únicos que passei com vocês. Foi sem dúvida um prazer conhecer-vos.

A todos os meus amigos, mas em especial à Márcia e à Marta por compreenderem mais do que tudo que uma verdadeira amizade pode durar anos mesmo estando longe da vista, mas perto do coração.

Finalmente aos meus pais, ao meu irmão, à minha Tia Branca e à minha prima Mónica agradeço pelo apoio incondicional em todos os sentidos. Sei que terei uma dívida de gratidão eterna para convosco.

ABSTRACT

Non-typhoidal *Salmonella* is a worldwide major foodborne zoonotic pathogen with a main reservoir in the intestinal tract of food-producing animals, from which it is readily transmitted to farm surrounding environments and the food chain. Expansion of successful multidrug-resistant (MDR) *Salmonella* serotypes and clones have been globally reported, but the factors imperative for their success, besides antibiotic resistance-ABR, remains scarcely explored. Tolerance to diverse antimicrobial non-antibiotic compounds (biocides) widely used in food-producing animals might constitute an adaptive feature to diverse environments and hosts. Among them are metals, often used in biocidal products (e.g. copper; silver-disinfectants, antiseptics/medicines or preservatives), in animal feed additives (e.g. copper, zinc) or in soil fertilisers/pesticides (e.g. arsenic, tellurium). Consequently, they can remain and accumulate in the animal-production setting and other environments, representing a long-term selective pressure potentially driving co-selection of metal-tolerant and/or ABR bacteria. In this context, for the elaboration of this thesis it was **hypothesized** that tolerance to metals/biocides contributes for the survival and emergence/persistence of particular MDR *Salmonella* serotypes and clones in diverse environments and hosts. The **main goal** was to assess the occurrence of acquired metal tolerance-Me^T genes as well as to characterize their associated genetic platforms and tolerance phenotype in different *Salmonella* serotypes and emergent clones. This assessment included 406 *Salmonella* isolates (2000-2014), belonging to 50 serotypes and diverse clones with different rates of frequency and emergence in the last 15 years. They were selected by source, antibiotic susceptibility phenotypes/genotypes, PFGE profiles and/or Sequence Types (ST) from a collection of ~2500 Portuguese isolates, and they were recovered in several Portuguese regions and different sources (humans, food products, food-animal production and the environment).

The search of acquired Me^T genes (PCR/sequencing) showed that those coding for copper (*pcoA-pcoD±silA-silE*), silver (*silA-silE*) and mercury (*merA*) tolerance were more frequent than those coding for arsenic (*arsB*) and tellurite (*terF*) tolerance. The occurrence of these diverse genes in isolates from variable sources, clones and serotypes suggests a genetic flow in different ecological communities and hosts. Of note are *Salmonella* belonging to recently emerging pig-associated serotypes/clones (MDR *S.* 4,[5],12:i:- and *S.* Typhimurium European clones/ST34 and *S.* Rissen/ST469) often isolated in the food chain and implicated in human infections in Europe/Portugal, which were more enriched in *sil±pco* genes than those associated with other food-animal settings (e.g. *S.* Enteritidis-poultry/eggs). The use of high copper concentrations allowed in feed for pigs comparing with other animals might have contributed for the selection and persistence of such emerging *sil±pco Salmonella*. Also, these isolates were more resistant to antibiotics than others without *sil±pco*, highlighting the importance of carrying diverse adaptive features to survival in environments contaminated with variable stressors as metals or antibiotics.

The characterisation of the genetic platforms carrying Me^T genes [genomic location (I-CeuI/S1-PFGE/hybridization), plasmid analysis (plasmid-based replicon typing/sequencing) and conjugation/transformation assays] identified variable elements contributing for the co-dispersion

of Me^T and ABR genes in diverse *Salmonella* serotypes. Large plasmids of different major families (e.g. Inc A/C, R, FIIA, HI1, HI2, N, P, I1), some transferable and described to be widespread in *Enterobacteriaceae*, were found to carry biocide (*oqxAB*-quaternary ammonium compounds), Me^T (*sil±pco*, *merA*, *arsB*, *terF*) and ABR genes [e.g. *tet(A)*, *sul1* and/or *sul3*-class 1 integrons] including clinically relevant ones (*bla*_{CTX-M-9} and *aac(6')-Ib-cr/oqxAB*). Although some of these plasmids have already been described as carrying Me^T genes, we unveiled the relevance of IncFIIA and IncN types in the dispersion of such genes. The co-location of both Me^T and ABR in the same plasmids also alert for the importance of single horizontal transfer events for their co-dispersion among bacteria sharing the same ecosystems and subjected to diverse selective pressures. Variable Me^T genes (*sil±pco*, *merA*, *terF*) were also found in non-transferable chromosomal regions with other ABR genes [e.g. *bla*_{TEM}, *tet(B)*, *sul2*] among several *Salmonella* serotypes, but mostly in the recently emergent MDR *S. 4,[5],12:i:-* and *S. Typhimurium* European clones and *S. Rissen*. These data highlight the importance of clones environmental adaptation not only for their spread but also for maintenance of Me^T genes. Concerning *sil±pco* genetic environments, our results and an *in silico* genomic analysis showed that those clusters are often adjacent and mostly inserted in a Tn7-like element (Tn6230) with different levels of recombination potentially implicated in its genome fixation, particularly in recently emergent *Salmonella* serotypes/clones.

Phenotypic susceptibility assays to copper and silver (MIC_{CuSO₄} and MIC_{AgNO₃} in aerobic and anaerobic atmospheres by standard agar dilution method) showed that *Salmonella* isolates (different serotypes/clones; wild type and transconjugants) carrying the *sil* efflux system (chromosomal or plasmids) were associated with higher tolerance to CuSO₄ in anaerobiosis and to AgNO₃ after prior exposure to silver in aerobiosis. These data demonstrate the role of *sil* gene cluster for *Salmonella* adaptation to different metal stressors present in food-animal settings, particularly to overcome the more toxic Cu⁺ occurring in reducing environments (e.g. animal gut, manure). Moreover, those modified methodological approaches in Cu/Ag tolerance assays were critical to identify isolates carrying *sil±pco* gene clusters and for the proposal of tolerance cut-offs to CuSO₄/AgNO₃, that can be used in further studies and allow comparison of data.

The **general conclusion** of this thesis is that the acquisition of metal tolerance genes, particularly to copper and silver, confers an advantage to clinically relevant multidrug-resistant *Salmonella* serotypes/clones by facilitating their better survival and persistence in environments contaminated with those metals across the food chain. The new information provided by this study might help to support the implementation of more efficient intervening measures associated with the use and/or accumulation of metals in diverse environments to prevent a wider expansion of foodborne zoonotic pathogens such as MDR *Salmonella* serotypes/clones or the emergence of new ones.

Keywords: *Salmonella*, Clones, Multidrug-resistance, Metals, Biocides

RESUMO

Salmonella não tifóide é um patógeno zoonótico de origem alimentar relevante a nível mundial. O seu principal reservatório é o trato intestinal dos animais, a partir do qual é facilmente transmitido para o ambiente de produção animal e cadeia alimentar. A expansão de serótipos e clones de *Salmonella* bem-sucedidos e que apresentam resistência a múltiplos antibióticos-MDR tem sido globalmente reportada. Contudo, outros fatores além da resistência a antibióticos-ABR que podem contribuir para esse sucesso permanecem pouco explorados. A tolerância a diversos compostos antimicrobianos não incluídos no grupo dos antibióticos (ex. biocidas), amplamente utilizados em produção animal, poderá constituir um fator adaptativo de *Salmonella* a diversos ambientes e hospedeiros. Entre estes compostos incluem-se os metais frequentemente utilizados em produtos biocidas (ex. cobre, prata-desinfetantes, antissépticos/medicamentos ou conservantes), aditivos na alimentação animal (ex. cobre, zinco) ou fertilizantes/pesticidas (ex. arsénio, telúrio). Na sequência do seu uso, os metais podem permanecer e acumular-se em diferentes ambientes, nomeadamente nos de produção animal, representando uma pressão seletiva a longo prazo que poderá levar à co-seleção de bactérias tolerantes a metais e/ou resistentes a antibióticos. Neste contexto, a **hipótese** desta tese foi de que a tolerância a metais/biocidas contribui para a sobrevivência e emergência/persistência de determinados serótipos e clones de *Salmonella* MDR em diversos ambientes e hospedeiros. O **principal objetivo** consistiu em avaliar a ocorrência de genes adquiridos de tolerância a metais-Me^T, caracterizar as plataformas genéticas em que se inserem e o fenótipo de tolerância em diferentes serótipos de *Salmonella* e clones emergentes. Foram incluídos 406 isolados de *Salmonella* (2000-2014) de 50 serótipos e diversos clones com taxas de frequência e de emergência variáveis nos últimos 15 anos. Estes isolados foram selecionados de uma coleção de ~2500 *Salmonella* de diversas regiões de Portugal com base na origem (humanos, alimentos, animais de produção e ambiente), fenótipos/genótipos de suscetibilidade aos antibióticos, perfis de PFGE e *Sequence Types* (ST).

A pesquisa de genes adquiridos de Me^T (PCR/sequenciação) mostrou que os que codificam para tolerância ao cobre (*pcoA-pcoD±silA-silE*), prata (*silA-silE*) e mercúrio (*merA*) foram mais frequentes do que os que codificam para tolerância ao arsénio (*arsB*) ou telúrio (*terF*). A ocorrência destes genes em isolados de diversas origens, clones e serótipos sugere um fluxo genético entre diferentes nichos ecológicos e hospedeiros. De destacar os serótipos/clones de *Salmonella* recentemente emergentes e associados a suínos (Clones Europeus/ST34 de *S.* 4,[5],12:i:- e *S.* Typhimurium, *S.* Rissen/ST469, todos MDR), frequentemente isolados na cadeia alimentar e implicados em infeções humanas na Europa/Portugal, mais enriquecidos em genes *sil±pco* do que os associados a outros animais de produção (ex. *S.* Enteritidis-aves/ovos). As elevadas concentrações de cobre permitidas nas rações de suínos comparativamente às de outros animais podem ter contribuído para a seleção e persistência destas *Salmonella* emergentes com *sil±pco*. Estes isolados também apresentaram maior resistência a antibióticos, destacando-se a importância da presença de vários fatores adaptativos para a sua sobrevivência em ambientes contaminados com diversos stresses como metais ou antibióticos.

A caracterização das plataformas genéticas portadoras de genes de Me^T [localização genômica (I-CeuI/S1-PFGE/hibridação), análise de plasmídeos (PL) (PCR-based *replicon typing*/sequenciação) e ensaios de conjugação/transformação] identificou vários elementos associados à co-dispersão de genes Me^T e ABR em diversos serótipos de *Salmonella*. Detetaram-se plasmídeos de elevado tamanho e de diferentes famílias (ex. Inc A/C, R, FIIA, HI1, HI2, N, P, I1), alguns transferíveis e considerados amplamente distribuídos em *Enterobacteriaceae*, como sendo portadores de genes de tolerância a biocidas (ex. *oqxAB*-compostos de amônio quaternário), Me^T (ex. *sil±pco*, *merA*, *arsB*, *terF*) e ABR [ex. *tet(A)*, integrões de classe 1-*sul1/sul3*], incluindo clinicamente relevantes (ex. *bla*_{CTX-M-9}, *aac(6')-Ib-cr/oqxAB*). Embora alguns desses plasmídeos tenham sido já descritos como portadores de genes Me^T, neste trabalho evidenciou-se a relevância dos IncFIIA e IncN na dispersão destes genes. A co-localização de genes Me^T e ABR nos mesmos plasmídeos reforça também a importância de eventos únicos de transferência horizontal para a sua co-dispersão entre bactérias que partilham os mesmos ecossistemas e sujeitas a diversas pressões seletivas. Vários genes Me^T (*sil±pco*, *merA*, *terF*) e ABR [ex. *bla*_{TEM}, *tet(B)*, *sul2*] foram também detetados em regiões cromossômicas não transferíveis em vários serótipos de *Salmonella*, mas principalmente nos clones Europeus de *S.* 4,[5],12:i:- e *S.* Typhimurium e *S.* Rissen, todos MDR e de emergência recente. Estes dados salientam a importância da adaptação dos clones a determinadas condições ambientais não só para a sua melhor propagação, mas também para a manutenção de genes Me^T. Os dados deste estudo juntamente com uma análise genômica *in silico* mostraram que os *clusters pco+sil* estão frequentemente adjacentes e inseridos em elementos Tn7-like (Tn6230), com diferentes níveis de recombinação possivelmente implicados na sua fixação genômica, particularmente em serótipos/clones de *Salmonella* de emergência recente.

Os ensaios de suscetibilidade ao cobre e à prata (CMI_{CuSO₄} e CMI_{AgNO₃} em aerobiose e anaerobiose pelo método standard de diluição em agar) mostraram que os isolados de *Salmonella* (diferentes serótipos/clones; selvagens e transconjugantes) portadores do sistema de efluxo *sil* (cromossoma ou plasmídeos) foram associados a uma maior tolerância a CuSO₄ em anaerobiose e a AgNO₃ após exposição prévia à prata em aerobiose. Estes dados contribuem para reforçar o papel do *cluster sil* na adaptação de *Salmonella* a diferentes metais presentes no ambiente de produção animal, particularmente para superar a forma mais tóxica Cu⁺ presente em ambientes reduzidos (ex. intestino animal, estrume). A modificação metodológica destes ensaios de tolerância ao Cu/Ag foi crítica para identificar isolados com *sil±pco* e para propor valores *cut-off* de tolerância, permitindo a comparação criteriosa de dados quando utilizados em estudos futuros.

Os resultados obtidos permitem inferir como **conclusão geral** que a aquisição de genes Me^T, particularmente os que codificam para tolerância ao Cu/Ag, conferem uma vantagem a determinados serótipos/clones clinicamente relevantes de *Salmonella*, facilitando a sua melhor sobrevivência e persistência em ambientes contaminados com esses metais ao longo da cadeia alimentar. Este conhecimento poderá suportar a implementação de medidas de intervenção mais eficientes relativas à utilização e/ou acumulação de metais em diversos ambientes, prevenindo assim a expansão e/ou emergência de patógenos zoonóticos como serótipos/clones de *Salmonella* MDR.

Palavras chave: *Salmonella*, Clones, Resistência a múltiplos antibióticos, Metais, Biocidas

CONTENTS

CHAPTER 1 – INTRODUCTION.....	1
1.1. The foodborne zoonotic <i>Salmonella</i>	3
1.1.1. Clinical relevance, transmission dynamics and epidemiology.....	3
1.1.2. Non-typhoidal <i>Salmonella</i> serotypes in poultry/eggs and pig products	6
1.1.3. <i>Salmonella</i> adaptive features to food-animal setting	14
1.2. Antibiotic-resistant non-typhoidal <i>Salmonella</i> infections and the food-animal link.....	15
1.2.1. Use of antibiotics and the growing burden of antibiotic resistance.....	15
1.2.2. The contribution of food-animals for the global burden of antibiotic-resistant <i>Salmonella</i>	20
1.2.2.1. Resistance to extended-spectrum β -lactams in <i>Salmonella</i>	24
1.2.2.2. Resistance to quinolones in <i>Salmonella</i>	33
1.2.3. Acquisition and dissemination of antimicrobial resistance in <i>Salmonella</i>	39
1.2.3.1. Clonal spread.....	40
1.2.3.2. Horizontal gene transfer	41
1.3. Bacterial tolerance to biocides, particularly to metals	43
1.3.1. Definition, classification, application and accumulation of metals and biocides... ..	44
1.3.2. Mechanisms of action of the biocide activity of metals.....	46
1.3.3. Mechanisms of tolerance to metals, particularly in <i>Salmonella</i>	49
1.3.3.1. Copper homeostasis and tolerance mechanisms	51
1.3.3.2. Silver tolerance mechanisms	54
1.3.3.3. Mercuric ion tolerance mechanisms.....	55
1.3.3.4. Tolerance mechanisms to metalloids arsenic and tellurite.....	57
1.3.4. Bacterial tolerance to metals and biocides and the possible link with the selection and persistence of antibiotic-resistant <i>Salmonella</i>	60
REFERENCES	65
CHAPTER 2 – OBJECTIVES AND OUTLINE OF THE STUDY	101
2.1. Statement of objectives	103
2.2. Outline of the thesis	105

CHAPTER 3 - RESULTS AND DISCUSSION.....107

3.1. Insights on the population structure and antimicrobial resistance of clinically-relevant <i>Salmonella</i> serotypes from clinical and non-clinical settings	109
3.1.1. Leakage of emerging clinically relevant multidrug-resistant <i>Salmonella</i> clones from pig farms	111
3.1.2. Characterization of the emerging clinically-relevant multidrug-resistant <i>Salmonella enterica</i> serotype 4,[5],12:i:- (monophasic variant of <i>S. Typhimurium</i>) clones.....	117
3.2. Analysis of the dispersion of metals/biocides tolerance genes and tolerance phenotype as well as their associated genetic platforms in diverse non-typhoidal <i>Salmonella</i> serotypes and clones	129
3.2.1. Metal tolerance in emerging clinically relevant multidrug-resistant <i>Salmonella enterica</i> serotype 4,[5],12:i:- clones circulating in Europe.....	131
3.2.2. Tolerance to multiple metal stressors in emerging non-typhoidal MDR <i>Salmonella</i> serotypes: a relevant role for copper in anaerobic conditions.....	141
3.2.3. <i>Salmonella enterica</i> serotype Bovismorbificans, a new host for CTX-M-9	185
3.2.4. Clinical <i>Salmonella</i> Typhimurium ST34 with metal tolerance genes and an IncHI2 plasmid carrying <i>oqxAB-aac(6')-Ib-cr</i> from Europe	189

CHAPTER 4 - CONCLUSIONS.....195

APPENDIX.....203

I- Review paper - "Salmonellosis: the role of poultry meat"	205
II- Protocols optimized for assessment of metal and biocide susceptibility by genotypic and/or phenotypic methods.....	221
III- Permissions for inclusion of papers, figures and tables	239

LIST OF FIGURES

Chapter 1 – Introduction

1.1.1. FIGURE 1 – Classification of <i>Salmonella</i> by different typing methods and disease syndrome	5
1.1.1. FIGURE 2 – MLST sequential scheme for obtaining <i>Salmonella</i> ST (http://mlst.warwick.ac.uk/mlst/dbs/Senterica)	6
1.1.2. FIGURE 3 – Infection routes and dissemination of <i>Salmonella</i> within the ecosystem.....	7
1.1.2. FIGURE 4 – Distribution of the major serotypes of non-typhoidal <i>Salmonella</i> associated with human cases (salmonellosis), poultry and pig meat in EU, 2011 to 2013	9
1.2.1. FIGURE 5 – Potential routes for dissemination of antimicrobial-resistant bacteria and resistance genes.....	19
1.2.1. FIGURE 6 – Clinical and epidemiological breakpoints of susceptibility to antibiotics	20
1.2.2. FIGURE 7 – Examples of antimicrobial resistance mechanisms in bacteria	23
1.2.2.1. FIGURE 8 – β -lactams target interactions and associated cell death mechanisms	24
1.2.2.2. FIGURE 9 – Quinolones target interactions and associated cell death mechanisms	33
1.2.3. FIGURE 10 – Genetic events responsible for horizontal and vertical transmission of genes that can favour the selection of antibiotic-resistant bacteria.....	39
1.2.3.2. FIGURE 11 – Horizontal gene transfer of antimicrobial resistance: (1) – transduction, (2) – conjugation and (3) - transposition	41
1.3.1. FIGURE 12 – Main sites of application of biocides in the food-animal setting as part of biosecurity measures	45
1.3.2. FIGURE 13 – Mechanisms of action and bacterial targets of biocides	47
1.3.2. FIGURE 14 – Antimicrobial mechanisms of metal toxicity.....	48
1.3.2. FIGURE 15 – Copper mechanisms of action within macrophages of mammalian host, following phagocytosis of bacteria	49
1.3.3. FIGURE 16 – Intrinsic or acquired metal tolerance mechanisms.....	51
1.3.3.1. FIGURE 17 – Copper homeostasis mechanisms based in Cue and Gol systems in <i>Salmonella</i>	52
1.3.3.1. FIGURE 18 – Proposed copper (<i>pco</i> cluster) tolerance mechanism	53

1.3.3.2. FIGURE 19 – Proposed silver (<i>sil</i> cluster) tolerance mechanism	54
1.3.3.3. FIGURE 20 – Proposed mercury (<i>mer</i> operon) tolerance mechanism.....	57
1.3.3.4. FIGURE 21 – Proposed arsenic (<i>ars</i> operon) tolerance mechanism.....	58
1.3.4. FIGURE 22 – Representation of the main types of multidrug-resistance efflux pumps involved in extrusion of QACs and several other antibiotics in Gram-negative bacteria	61

Chapter 3 – Results and discussion

3.1.1. FIGURE 1 – PFGE patterns of <i>Salmonella</i> isolates from piggeries and others previously characterized	115
3.1.2. FIGURE S1 – <i>Xba</i> I profiles of <i>Salmonella enterica</i> 4,[5],12:i:- isolates	127
3.2.3. FIGURE 1 – (a) Genetic context of <i>bla</i> _{CTX-M-9} within an Inc60 variant identified in a <i>Salmonella enterica</i> serotype Bovismorbificans isolate (HP507391). (b) Antimicrobial susceptibility profiles and resistance genotypes of <i>S. Bovismorbificans</i> carrying <i>bla</i> _{CTX-M-9} (HP507391) and the corresponding transconjugant (HP507391 T).	187
3.2.4. FIGURE 1 – Genetic environment (12 110 bp) of <i>oqxAB-aac(6')-Ib-cr</i> genes in an IncHI2 plasmid from an ST34 <i>Salmonella</i> Typhimurium strain	191
3.2.4. FIGURE S1 – PFGE patterns of clinical ciprofloxacin-resistant <i>S. enterica</i> isolates carrying <i>oqxAB</i> and <i>aac(6')-Ib-cr</i> genes.....	193

LIST OF TABLES

Chapter 1 – Introduction

1.1.2. TABLE 1 – <i>Salmonella</i> outbreaks and emerging clones linked to poultry (including eggs), pig and products thereof.....	11
1.2.1. TABLE 2 – Antimicrobials approved for use in food-producing animals.....	16
1.2.2.1. TABLE 3 - <i>Salmonella</i> serotypes/clones carrying extended-spectrum (ESBLs) and plasmid-encoded AmpC β -lactamases recovered from poultry, pigs and products thereof.....	27
1.2.2.2. TABLE 4 – <i>Salmonella</i> serotypes/clones carrying plasmid-mediated quinolone resistance (PMQR) genes recovered from poultry, pigs and products thereof.....	35
1.3.1. TABLE 5 – Classification of biocides based on the bacterial target	44

Chapter 3 – Results and discussion

3.1.1. TABLE 1 – Characterization of the <i>Salmonella</i> clones from Portuguese piggeries.	114
3.1.2. TABLE 1 – Genotypic and phenotypic characterization of <i>Salmonella enterica</i> serotype 4,[5],12:i:- clones from Portugal	121
3.2.1. TABLE 1 – Epidemiological and antibiotic and metal tolerance genetic features of <i>Salmonella enterica</i> subsp. <i>enterica</i> serotype 4,[5],12:i:- isolates from Portugal	133
3.2.1. TABLE 2 – Minimum inhibitory concentrations (MICs) of CuSO ₄ and AgNO ₃ for <i>Salmonella enterica</i> subsp. <i>enterica</i> serotype 4,[5],12:i:- isolates	136
3.2.1. TABLE S1 – Primers used in the PCR to search for acquired metal tolerance genes	139
3.2.2. TABLE 1 – Epidemiological features and genotypic/phenotypic characterization of <i>Salmonella enterica</i> isolates carrying acquired copper and silver tolerance genes (<i>sil</i> and <i>pco</i>)	159
3.2.2. TABLE 2 – Minimum inhibitory concentrations of CuSO ₄ for <i>Salmonella enterica</i> isolates	162

3.2.2. TABLE 3 – Minimum inhibitory concentrations of AgNO ₃ for <i>Salmonella enterica</i> isolates	164
3.2.2. TABLE 4 – Co-transfer of copper and silver tolerance with antibiotic resistance	166
3.2.2. TABLE S1 – Epidemiological features and genotypic/phenotypic characterization of <i>Salmonella enterica</i> isolates with absence of acquired copper and silver tolerance genes (<i>sil</i> and <i>pco</i>)	168
3.2.2. TABLE S2 – Primers used in PCR assays to search acquired metal tolerance genes	174
3.2.2. TABLE S3 – Dispersion of <i>sil±pco</i> gene clusters and their genetic environments in <i>Salmonella enterica</i> genomes available at GenBank database	176
3.2.4. TABLE 1 – Antimicrobial susceptibility phenotypes and genotypes of <i>Salmonella</i> Typhimurium isolates (234 and 248), the selected transformant (248_T) and the recipient (<i>Escherichia coli</i> DH5α)	191

LIST OF BOXES

Chapter 1 – Introduction

1.1.1. BOX 1 - Phenotyping methods – serotyping and phage typing	5
1.1.1. BOX 2 - Molecular typing methods – PFGE and MLST	5
1.2.1. BOX 3 - Antimicrobial resistance: concepts	20
1.2.2. BOX 4 - Antibiotic resistance mechanisms in <i>Salmonella</i>	23
1.2.2.1. BOX 5 - Classification and mechanism of action of β -lactams	24
1.2.2.2. BOX 6 - Classification and mechanisms of action of quinolones.....	33

LIST OF ABBREVIATIONS

	Description
ABR	Antibiotic resistance
Ag	Silver chemical symbol
As	Arsenic chemical symbol
CDC	Centers for Disease Control and Prevention
CLSI	Clinical and Laboratory Standard Institute
Cu	Copper chemical symbol
DNA	Deoxyribonucleic Acid
dNTP	Deoxynucleoside triphosphates
eBG	eBurstGroups
ECDC	European Center for Disease Prevention and Control
ECOFFs	Epidemiological Cut-Off Values
EDTA	Ethylenediaminetetraacetic acid
EEA	European Economic Area
EFSA	European Food Safety Authority
EMA	European Medicines Agency
ESBL	Extended-Spectrum β -Lactamases
ESC	Extended-Spectrum Cephalosporins
EU	European Union
EUCAST	European Committee on Antimicrobial Susceptibility Testing
GI	Genomic Island
Hg	Mercury chemical symbol
HGT	Horizontal Gene Transfer
ICE	Integrative Conjugative Element
IR	Inverted Repeat
IS	Insertion Sequence
Kb	Kilobase
LGT	Lateral Gene Transfer
MCO	Multicopper oxidase
MDR	Multidrug Resistance
MH	Mueller-Hinton

MIC	Minimum Inhibitory Concentration
MGE	Mobile Genetic Element
MLST	Multilocus Sequence Typing
mM	Milimolar
NCBI	National Center for Biotechnology Information
PAI	Pathogenicity Island
PBP_s	Penicillin Binding Proteins
PCR	Polymerase Chain Reaction
PFGE	Pulsed-Field Gel Electrophoresis
PMQR	Plasmid-Mediated Quinolone Resistance
QACs	Quaternary Ammonium Compounds
QRDR	Quinolone-Resistance Determining Region
RFLP	Restriction Fragment Length Polymorphism
Rep	Plasmid-specific replication initiator protein
RNA	Ribonucleic Acid
RND	Resistance-nodulation-cell division
SCENIHR	Scientific Committee on Emerging and Newly Identified Health Risks
ST	Sequence Type
Te	Tellurium chemical symbol
USA	United States of America
WHO	World Health Organization



Chapter | 1

Introduction

“It’s not that I’m smart, it’s just I stay with problems longer.”

Albert Einstein

1.1. The foodborne zoonotic *Salmonella*

1.1.1. Clinical relevance, transmission dynamics and epidemiology

Salmonella enterica (henceforward *S. enterica*) infections are a worldwide major public health concern, being the foodborne illness salmonellosis caused by non-typhoidal *Salmonella* (serotypes other than *S. Typhi* and *S. Paratyphi*) (See box 1 for details). Currently, there are over 2500 identified *Salmonella* serotypes, although *S. enterica* subspecies I *enterica* includes all of the major serotypes that are pathogenic to humans (See box 1 for details) (1, 2). Salmonellosis is typically characterized by a self-limiting gastroenteritis syndrome (manifested as diarrhoea as main symptom, although fever, vomiting and abdominal pain or cramps can occur), with an incubation period between 4-72h, being death rare (<1%) (3-5). In healthy humans, the infectious dose is generally 10^6 to 10^8 bacterial cells, but lower number can cause disease in predisposing people, including infants and the elderly (4). Although uncommon, life-threatening invasive infections with bacteraemia (5-10% of infected persons) and/or other extra-intestinal infections may occur, affecting especially the risk groups (infants, young children, older people and the immunocompromised patients) (3-5). In severe cases, effective antimicrobial agents are essential, being the emergence of *Salmonella* resistant to critical antibiotics of concern (3, 4).

In industrialized countries, the main reservoir of non-typhoidal *Salmonella* is the gastrointestinal tract of warm-blooded animals, including food-producing animals from which it is readily transmitted to farm surrounding environments and the food chain (5-7). Therefore, although other sources (e.g. contact with animals/reptiles, environment or person-to-person) are recognized, the most relevant transmission to humans occurs mainly through the ingestion of contaminated food (e.g. eggs, poultry, pork, beef, dairy products and fresh produce), leading to a high global impact in human health (7). It was estimated that non-typhoidal *Salmonella* causes around 93.8 million of illnesses and 155.000 deaths each year worldwide (8). In the USA, the CDC estimated 1 million annual cases of foodborne salmonellosis, being associated with the largest number of hospitalizations (35%) and deaths (28%) compared with other foodborne microorganisms (9). In Europe, in the last year's salmonellosis has been the second most common zoonosis (82.694 confirmed cases and 20.4 cases per 100,000 populations in 2013) and the most frequent cause of foodborne outbreaks (22.5% of all cases), in spite of the reported decreasing trend scenario resulting from *Salmonella* control programmes (10).

Although different serotypes have been associated with salmonellosis, a limited number is responsible for most human infections, being *S. enterica* serotype Enteritidis (henceforward *S. Enteritidis*) the most frequent one in European Union (EU) (39.5% in 2013), USA (14.5% in 2012) and Canada (44.2% in 2013) followed by *S. enterica* serotype Typhimurium (henceforward *S. Typhimurium*) (20.2% in EU, 2013; 11.6% in USA, 2012; 8.5% in Canada, 2013) (10-12). *S. Enteritidis* is commonly associated with poultry and products thereof, while *S. Typhimurium* has a wider species range, including pigs and cattle as well as poultry (13). Therefore, foods of animal origin, including contaminated poultry products (eggs, egg products, poultry and turkey meat), but also pork have been considered the main vehicles of *Salmonella* infection (10, 14, 15).

In recent years, with the implementation of *Salmonella* control programmes mainly along poultry/egg production (e.g. in the EU and USA), changing trends in foodborne salmonellosis and associated serotypes were observed, with the expansion of previously less common serotypes (e.g. increase in reported cases of *S. enterica* Typhimurium monophasic variant – henceforward *S. 4,[5],12:i:-*) causing human infections through food chain and with origin in different food-animal species. Importantly, the increasing occurrence of these serotypes has been associated with certain subtypes, designated clonal lineages or clones (characterized by typing methods of reference as Pulsed-Field Gel Electrophoresis or Multilocus Sequence typing) (See box 2 for details) (16), which in turn exhibit resistance to multiple antimicrobial agents (2, 3, 17, 18). With the increasing globalization of foodstuffs like poultry and pig meat, which are two of the most consumed and increasing globally traded meat products (19, 20), and changes in food-animal production and in consumer behaviours (e.g. increased consumption of fresh produce) new problems might arise regarding salmonellosis control. An overview of the role of food-animals, mainly poultry/eggs and pig (the most common sources of human infection) on salmonellosis at a global scale and in the persistence/emergence of specific *Salmonella* serotypes/clones is provided on the next topic.

BOX 1 - Serotyping and Phage Typing

Salmonella spp. is classified in ≈2500 serotypes based on the classical Kauffmann–White–Le Minor scheme, with the majority of them (≈1500) belonging to *S. enterica* subsp. I *enterica*, a group of greatest clinical relevance due to its common association with humans and warm-blooded animals. In this technique, specific agglutination reactions are observed between adsorbed antisera that contain specific antibodies (“factors”), and the corresponding antigens when present within either surface lipopolysaccharides (LPS, O-antigens) or two alternatively expressed phases of the flagellar antigens (proteins, H-antigens) (Figure 1) (1, 21). Based on the type of antigens detected each serotype is represented by an antigenic formulae written as O:H1:H2 (e.g. antigenic formulae 1,4,[5],12:i:1,2 defines *S. Typhimurium*) (1). In addition to classical/conventional serotyping, others methods such as PCR or genoserotyping (a microarray assay) can also be used for the identification of the most common reported *Salmonella* serotypes through detection of the genes coding surface O and H antigens (16). Phage typing is carried out by infecting a specific *Salmonella* serotype (the most common Typhimurium and Enteritidis), with a number of specific bacteriophages, leading to a typing scheme based on the susceptibility to lysis (22).




FIGURE 1 – Classification of *Salmonella* by classical/conventional typing methods and disease syndrome [adapted from reference (21)].

BOX 2 - Molecular typing methods – PFGE, MLST and others

Pulsed-Field Gel Electrophoresis (PFGE) has a higher discriminatory power comparing for example with serotyping and phagotyping and is widely used for characterizing *Salmonella*, particularly in outbreak investigations. This method uses restriction enzymes (e.g. *XbaI* as primary and *BlnI* or *SpeI* as additional enzymes are currently used) to cut bacterial DNA in specific restriction sites, being these fragments separated through a gel matrix in a size-dependent manner by carefully orientated electrical pulses (16). The analysis of *Salmonella* DNA fingerprint on the gel is then used to discriminate PFGE-profiles and clones.

Multilocus Sequence Typing (MLST) is based on sequencing of internal fragments of seven housekeeping genes, usually distributed around the chromosome. For each of the seven genes, different sequences (alleles) occur. To obtain an MLST type, the seven gene sequences are deposited and forwarded to the online database (<http://mlst.warwick.ac.uk/mlst/>) and an allelic profile or Sequence Type (ST) is returned (Figure 2). Additionally, two or more STs that share six of the seven alleles that defined the ST were further grouped in genetically closely related clusters, named eBurstGroups (eBGs) (21).

BOX 2 - Molecular typing methods – PFGE, MLST and others (cont.)

Other molecular typing methods are also recently used for the characterization of *Salmonella* outbreaks, such as Multiple-Locus Variable number tandem repeat Analysis (MLVA) or Whole Genome Sequencing (WGS) (16). The MLVA might have an improved discriminatory power compared with PFGE, thus can be used as a complement of this technique (16). It is based on measurement of length of variable number of tandem repeat (VNTR) loci by PCR amplification and electrophoresis, using this information to create a genotype to distinguish between isolates of the same serotype (16). WGS is based on the full sequencing of a bacterial genome, thus achieving the highest discriminatory power between the several methods. Nevertheless, some challenges have been reported for WGS such as the storage, interpretation, annotation and harmonization of the huge amount of information (16).




FIGURE 2 – MLST sequential scheme for obtaining *Salmonella* ST (<http://mlst.warwick.ac.uk/mlst/dbs/Senterica>). An example is given for ST19 achievement, which is characteristic of *S. Typhimurium* and *S. 4,[5],12:i:-*.

1.1.2. Non-typhoidal *Salmonella* serotypes in poultry/eggs and pig products

Animal populations, in particular chicken, but also turkey and pigs, are frequently colonized with *Salmonella* without manifestation of detectable symptoms (sub-clinical infections/healthy carriers). They are colonized by horizontal (e.g. contaminated feed and water) and vertical transmission at primary production level (6, 7, 23). Additionally, in early stages of the production line (e.g. slaughter process of the animals, egg collecting or milking) cross-contamination can also occur (24, 25). The presence of *Salmonella* in asymptomatic healthy animals, that may excrete *Salmonella* continuously or intermittently, is suggested as the main risk factor for allowing bacteria to be easily

transmitted vertically in table eggs (through infected chickens), and in animal meat to humans. Animal faeces may also contaminate farmland, surface water flow and vegetables if they are fertilized with untreated animal manure (Figure 3) (13, 23, 26).




FIGURE 3 – Infection routes and dissemination of *Salmonella* within the ecosystem [reprinted with permission from reference (26)].

In Europe, it is assumed that the observed reduction in salmonellosis cases (32% between 2008 and 2012) is mainly due to implemented successful *Salmonella* control measures (e.g. surveillance, biosecurity, sanitation/disinfection, feed and water additives, vaccination) (25, 27, 28), targeting poultry (including breeding flocks of *Gallus gallus*, broilers and breeding and fattening flocks of turkeys) and eggs production (including laying hens and eggs) (10, 29-32). Those measures are focused in particular serotypes, such as *S. Enteritidis*, *S. Typhimurium* (including the monophasic variant *S. 4,[5],12:i:-*), *S. Infantis*, *S. Virchow* and *S. Hadar*, considered of public health significance (6, 10, 13, 27, 32). This scenario led to the achievement of reduction of *Salmonella* rates for poultry populations in most EU countries and lower non-compliance regarding *Salmonella* in poultry/eggs products (6, 10). Moreover, these decreasing contamination rates in European raw poultry products are in agreement with those recently observed in industrialized countries from other geographical regions applying pathogen reduction programmes, such as the USA (27, 33, 34). It is worth to mention, that by 2000's high incidence of *Salmonella* in poultry products was reported in the EU, with rates higher than 50% for several countries (35). In the 2013 zoonosis EFSA/ECDC report involving

data from European countries, as in previous years, *Salmonella* was most frequently reported, although at low levels, in fresh turkey (5.4%) and fresh broiler meat (3.5%), in comparison with eggs (0.1%) or fresh pig meat (0.7%) (10). Despite the highest incidence being detected in poultry meat, consumption of contaminated eggs and pig meat still remain the most important sources of food-borne *Salmonella* outbreaks (10). In fact, using quantitative source attribution models the higher number of human salmonellosis cases in Europe was attributable to eggs (65% in 2011 and 17% in 2012) and pigs (28% in 2011 and 56.8% in 2012) comparing with broilers (2.4% in 2011 and 10.6% in 2012) and turkeys (2.6% in 2011 and 4.5% in 2012) (14, 15). However, diverse surveys targeted to detect *Salmonella* in poultry and pig products in developing countries, some with expansion of the food-animal industry, still detected high percentages of positive samples, ranging from ~13-39%-poultry/~39%-pig in South America (36-38), ~35%-poultry/~14-40%-pigs in Africa (39-42) and ~35-50%-poultry/~30-40%-pig in Asia (43-47). Those differences possibly reflect diverse poultry production husbandry practices and absence of control measures along the food chain, highlighting the importance of the extensive international trade of animals and products thereof for *Salmonella* spread (6).

Worldwide data about *Salmonella* serotypes prevalence in humans and in the diverse range of foodstuffs have contributed to establish an epidemiological link between salmonellosis and poultry or pig products, with diverse serotypes overlapping between humans and poultry (chicken and turkey) or pig meat (Figure 4). In EU, recent changes in the frequency of *Salmonella* serotypes causing human infections were reported, which in some cases were in line with those occurring in poultry and pig meat, reflecting the importance of food chain transmission by these type of products to humans (Figure 4). Nevertheless, interpretation of these data should be cautious, owing to limitations in the number of human and food isolates serotyped each year. Of particular relevance is the decrease in *S. Enteritidis* human cases (19% reduction between 2011 and 2013 in EU, following the observed reduction in human salmonellosis), a serotype typically associated with poultry meat and eggs consumption, in line with a decreasing trend in poultry and poultry products pointed by different studies (11, 27, 48-50). Increasing occurrences of other serotypes implicated in human infections (e.g. *S. Infantis*, *S. Stanley*, *S. Kentucky*) related to poultry meat (chicken and turkey) have been reported in the EU (Figure 4). However, one major difference in the serotype pattern between humans and poultry is related to *S. Typhimurium* and its monophasic variant, both frequently associated with human salmonellosis cases, but less common in poultry meat, being pigs and pig meat the main source (10). In USA and Canada, other serotypes like *S. Heidelberg* and *S. Kentucky* have emerged as predominant serotypes in poultry and have also been implicated in human salmonellosis, beyond *S. Enteritidis* (11, 12, 27, 33, 34, 51-53).




FIGURE 4 – Distribution of the major serotypes of non-typhoidal *Salmonella* associated with human cases (salmonellosis), poultry and pig meat in EU, 2011 to 2013. The data were obtained from EFSA reports: humans (2011 to 2013) and turkey/broiler/pig meat (2013) (10); turkey/broiler/pig meat (2011 to 2012) (54, 55) (Last accessed 1 November 2015). The percentages were calculated based on the total number of serotyped isolates per type of meat or human salmonellosis cases.

The shift in *Salmonella* serotypes related to poultry and poultry production has been associated with the spread of certain clones, with most of them being responsible for large outbreaks (Table 1). For instance, *S. Infantis* increase (26.5% in human cases between 2011 and 2013 accounting for the fourth most common serotype and the most reported in broilers and broiler meat) (10), has been associated with the spread of several clones of broiler origin in diverse European countries, including the dominant Hungarian clone (56-58). Also, *S. Stanley* showed an increase since 2011 with a peak in 2012, being the sixth most common human serotype and one of the three most common in turkey meat, together with *S. Derby* and *S. Kentucky* in 2013 (10). A large *S. Stanley* outbreak caused by a new clone (novel PFGE-type), was linked with the consumption of turkey meat, and is still circulating in the European food market (at least since 2011), with a considerable risk of becoming endemic in the poultry production chain in Europe (59-61). In the USA, *S. Heidelberg* in particular has been identified as one of the top human and

poultry serotypes, with several clones implicated in diverse large multistate outbreaks resulting from the consumption of contaminated chicken or turkey products (27, 53). The spread and the global persistence of serotype *S. Kentucky* reflect other particular situation related to the increased globalization of travel and food/animal trade in different geographical regions. This serotype has been associated with a worldwide (Europe, Africa, and Asia) spread of a particular epidemic clone (*S. Kentucky* ST198-X1), recovered from several livestock reservoirs, particularly poultry farms, with chicken and turkey implicated as the potential major human infection vehicles (18, 62-65).

TABLE 1 – *Salmonella* outbreaks and emerging clones linked to poultry (including eggs), pig and products thereof.

Serotype (clone designation when applied)	Source	Year(s) ^a	Country	No. of ^b :		Reference(s)
				Cases	Deaths	
4:[5],12:i:- (European clone) 4:[5],12:i:-	Pork products	2006	Luxembourg	133	1	(66)
	Banquet pot pies (chicken or turkey)	2007	USA	272	0	CDC, 2015 ^c
	Dried pork sausage	2010	France	69	0	(67)
	Dried pork sausage	2011	France	337	0	(68)
	Pork	2015	USA	188	0	CDC, 2015 ^c
Derby	Raw fermented pork	2013-2014	Germany	145	NS	(69)
Enteritidis	Eggs, scrambled	2007	USA	81	0	CDC, 2015 ^c
	Eggnog	2008	USA	18	0	CDC, 2015 ^c
	Eggs, scrambled	2009	USA	59	0	CDC, 2015 ^c
	Eggs and poultry	2009	United Kingdom	152	2	(70)
	Shell eggs	2010	USA	1939	NS	CDC, 2015 ^c
	Raw eggs	2014	Europe (multi-country)	287	1	(71)
	Chicken	2015-	USA	5	0	CDC, 2015 ^c
	Chicken	2015-	USA	15	0	CDC, 2015 ^c
	Turkey burgers	2010-2011	USA	12	0	CDC, 2015 ^c
	Live poultry	2012	USA	46	0	CDC, 2015 ^c
Hadar	Broiled chicken livers	2011	USA	190	0	CDC, 2015 ^c
	Ground turkey	2011	USA	136	1	CDC, 2015 ^c
	Chicken	2012-2013	USA	134	0	CDC, 2015 ^c
Heidelberg	Chicken	2013-2014	USA, Puerto Rico	634	0	CDC, 2015 ^c
	Chicken	2014	USA	9	0	CDC, 2015 ^c
	Broilers and chicken	2004-2009	Europe	NA	NS	(56-58)
Infantis (Hungarian clone)	Broiler chickens	2007-2009	Israel	NA	NS	(50)
Infantis (Israel clone)	Raw pork products	2013	Germany	267	1	(72)
Infantis	Raw poultry products	2015-	Canada	98	0	PHAC, 2015 ^f
Kentucky (Sequence Type 198-X1 clone)	Chicken and turkey	2002-2013	Europe, Africa, Asia	NA	NS	(62-64)
Montevideo	Live poultry	2012	USA	93	1	CDC, 2015 ^c
Stanley	Turkey	2011-2013	Europe	710	NS	(59, 61)
Typhimurium	Raw egg mayonnaise	2009	Australia	71	0	AGDH, 2015 ^d
	Salami	2010	Denmark	20	0	(73)
	Duck eggs	2010	Ireland	24	0	FSAI, 2015 ^e
	Live poultry	2013	USA	356	0	CDC, 2015 ^c

^a “year-” include ongoing reported outbreaks.^b Estimated number of cases only when outbreaks were reported. NA, not applicable in emerging clones. NS, not specified.^c CDC, 2015. *Salmonella*. Reports of selected *Salmonella* Outbreak Investigations. Available at: <http://www.cdc.gov/salmonella/outbreaks.html> [accessed December 2015].^d AGDH, 2015. Annual report of the OzFoodNet network, 2009. Available at: <http://www.health.gov.au/internet/main/publishing.nsf/Content/cda-cdi3404b.htm> [accessed December 2015].^e FSAI, 2015. Large outbreak in recent years linked to *Salmonella* in duck eggs. Available at: <https://www.fsai.ie/14092010.html> [accessed December 2015].^f PHAC, 2015. Outbreak of *Salmonella* infections under investigation. Available at: <http://www.phac-aspc.gc.ca/phn-asp/2015/salmonella-infantis-eng.php> [accessed December 2015].

With the implementation of control programs in poultry/eggs production and with the subsequent *Salmonella* prevalence reduction, the role of pork products has been enhanced, being nowadays one of the most important sources of human salmonellosis at global or the EU level (10, 74). In fact, using quantitative source attribution models, EFSA estimated that around 28% (14) to 56.8% (15) of the human salmonellosis cases are attributable to pigs. Concerning pig production, control programs are not compulsory at the moment by the Regulation (EC) No. 2160/2003 of the European Parliament and Council, which are primarily focused on *Salmonella* control at poultry production (29). Most of national monitoring programs for *Salmonella* on pig meat and products thereof are based on sampling at the slaughterhouse (e.g. animal faeces, lymph nodes, organ or tissue samples, carcass or nasal swabs) and/or at the processing or cutting plants (10). In fact, EFSA reported a significant percentage of *Salmonella* in pig samples (8.1%) but also at the herd (14.9%) and slaughter (30.0%) batch levels (10). With a similar scenario, the United Kingdom in a baseline survey also reported a high prevalence of *Salmonella* in faecal samples (30.5%), rectal swabs (24.0%) and carcass swabs (9.6%) of slaughter pigs (75). The presence of *Salmonella* in all steps of pig production chain, which may contribute to meat contamination, alert to the need for implementing proper and effective measures to detect and control *Salmonella* serotypes with public health significance, in the pig setting and particularly at the primary farm production level (25, 76).

In Europe, *S. Typhimurium* has been the most frequently reported serotype in pigs (47.8%) and pig meat (30.7%) in line with its frequent association with human salmonellosis (20.2% in human cases in 2013 accounting for the second most common serotype) (Figure 4). On the other hand, the second most common serotype in pig meat, *S. Derby* (14.8% in pigs and 27.1% in pig meat) showed also an increase in human salmonellosis cases (0.9% in 2012 to 1.1% in 2013), although mainly due to an outbreak associated with raw fermented pork (Table 1; Figure 4) (10). Similarly, in a European baseline study reporting a high prevalence (31.8%) of *Salmonella*-positive holdings with breeding pigs, the two most common serotypes in breeding and production holding were *S. Derby* (29.6% and 28.5%) and *S. Typhimurium* (25.4% and 20.1%) (77, 78). These two serotypes were also the most frequently isolated in clinical and non-clinical cases in pigs in USA (27, 79).

Noteworthy, over the last years in several European countries, including Portugal, it has been reported an increase in the prevalence (68.8% between 2011 and 2013) of *S. 4,[5],12:i:-*, a pig-related serotype, ranking nowadays third in frequency (Figure 4) (10, 80, 81). In the USA and Canada, the emergence and spread of this serotype associated with human infections, has also been observed, being the fifth most common in both regions (11, 79, 82, 83). *S. 4,[5],12:i:-* isolates have been responsible for several recent worldwide

large outbreaks (Table 1) (66-68, 84) and food animals, particularly pigs, were pointed out as its reservoir and their derived products as the main infection vehicles (77, 78, 80, 85). This serotype is, since 2011, targeted by EU actions to detect and control *Salmonella* serotypes of concern (32), requiring further surveillance and mitigation strategies, including its epidemiological success characterization. Two major *S.* 4,[5],12:i:- clones were recognised and were previously designated as “European” and “Spanish”. The “European” clone has been circulating in diverse geographical regions of Europe and has been frequently associated with DT120 and DT193 phage types (85-88), whereas the “Spanish” clone, spread in the Iberian Peninsula since 2002, was mostly associated with DT104/U302 phage types (89, 90). As a result of the worldwide increase of *S.* 4,[5],12:i:- clones causing human infections (86) and its increasing expansion in pig populations (85), the monophasic variant is currently considered an emerging epidemic serotype (80, 86).

Other pig-related serotype is *S.* Rissen, recently reported as having an increase in their frequency globally (10). It is now in the TOP20 of the most reported serotypes associated with human salmonellosis in Europe but, in Portugal, this serotype ranked fourth in frequency between 2002 and 2013 (10, 81). *S.* Rissen is usually associated with a predominant PFGE-type clone, resistant to multiple antibiotics and frequently associated with *sul1*-typical or *sul3*-atypical class 1 integrons (24, 91-94). This serotype has been reported in samples from European pigs, pork and surrounding environments. For example, *S.* Rissen was detected in lairage, pig carcasses and intestines samples (17.4%) from Belgium (95), in pig liver and colon content (4.6%) from Italy (92), in pigs and farm environments (7.5% to 15.38%) from Northern Ireland (96) or in pig carcass (10.1%) from The Netherlands (94). Although, in the Iberian Peninsula an increasing frequency (~>20%) of this serotype have been reported (48, 91, 93, 97-99). In fact, in a European baseline survey study, a high incidence of *S.* Rissen in breeding and production pig holdings has been found in Portugal (40% and 22.4%, respectively) and Spain (25% and 29.7%, respectively), being the 1st or 2nd most frequently reported (77, 78). Other non-European regions, such as USA (100-102) and Thailand (101, 103-106) have also been found this serotype in pig abattoirs and retail meat.

These and other examples of multi-country/multistate outbreaks or clonal expansion of *Salmonella* infections linked to poultry and pig meat (Table 1) serves as a reminder of the importance of acting upon any *Salmonella* contamination in the food chain and monitoring to detect the emergence of any serotype or new clone. This scenario also alerts for the need of inclusion of those serotypes or particular clones, considered of public health significance, in control and surveillance programmes (13, 27, 60).

1.1.3. *Salmonella* adaptive features to food-animal setting

Salmonella serotypes and clones associated with human infections and with an enhanced ability to colonize several food animals, able to persist across the food chain (e.g. primary production on-farm, slaughter operations, equipment's and meat handlers, retail meat) with efficient transmission and rapid spread, are of public health relevance (17, 18, 56, 107). Although the exact mechanisms of their persistence and spread in poultry and pig production are still largely unknown, recent studies focusing on emergent poultry and pig-associated *Salmonella* serotypes/clones unveiled specific features (e.g. enhanced colonization/virulence, metal tolerance), other than antibiotic resistance, that could provide a significant advantage both in the environment and in the host (poultry/pig/human) (27, 108, 109).

Recent studies found the presence of diverse virulence genes in *S. Typhimurium* (108, 110) and in specific clones of its monophasic variant *S. 4,[5],12:i:-* strains circulating in Europe (80, 85, 90, 111) associated with a better adaptation to the food-animal host (110). These genes encoded for proteins related to improved intestinal colonization (e.g. *clpB* - chaperone), adhesion (e.g. *csgA*, *fimA*, *stbD* - fimbrial operons) or infection (e.g. *sopA*, *avrA*, *sseI* - effector genes that function to alter host cell physiology and promote bacterial survival in host tissues; *marT* – transcriptional regulator that activates the expression of an extracellular matrix adhesin, the *MisL*; and *fliB* - flagellar operon) (110). Also, a recent study showed that *S. Typhimurium* DT193 and *S. 4,[5],12:i:-*, two serotypes with increased adaptation to faecal pH and organic acid supplementation, were associated with a long-term survival in porcine faeces compared to others such as *S. Derby* and *S. Bredeney* (112). In fact, the persistence of these two pig-associated *Salmonella* serotypes may be important for a better transmission by faecal-oral routes in pig herds (112).

In poultry associated serotypes other examples could be cited. In Israel, a comparison between the pre-emergent and the emergent *S. Infantis* clones showed in the latter the presence of a megaplasmid (hybrid plasmid pESI belonging to IncI1 and IncP-1 α families of ~280Kb) associated with an increased tolerance to stress factors (e.g. mercury and oxidative stress) and virulence/pathogenicity (e.g. enhanced biofilm formation, adhesion and invasion into avian and mammalian host cells) (113). Also, a genomic study of several predominant *Salmonella* serotypes from Canadian broiler chickens showed the presence of multiple features related to pathogenicity (e.g. genes encoding adhesins, flagellar proteins, iron acquisition systems, type III secretion system-T3SS) and stress tolerance (e.g. metal and antiseptic tolerance genes; better acid-stress response) (51). *S. Heidelberg*, including ground turkey outbreak isolates, carried phages and plasmids with diverse virulence factors (e.g. P2-like phage-*sopE1* gene, IncX-T4SS), which could play a

role in their virulence (a serotype highly associated with invasive infections), colonization and persistence (a poultry-associated serotype) (27, 53). In *S. Kentucky*, the acquisition of an *E. coli* ColV virulence plasmid (IncFIB/FIIA families) with genes encoding particular virulence features (e.g. afimbrial adhesins and colicins), was also associated with enhanced colonization of the chicken cecum and persistence in the avian extraintestinal environment, particularly in a dominant avian clonal type (27, 114).

These features can play a role in the successful spread of emergent and virulent serotypes/clones that could contribute in short time to replacing other *Salmonella*. Moreover, those emergent *Salmonella* serotypes are usually enriched with antimicrobial (antibiotics, metals or biocides) resistance determinants conferring multidrug-resistance/tolerance (27, 51, 53, 112, 113), which is currently one of the major public health concerns (discussed in the next topics).

1.2. Antibiotic-resistant non-typhoidal *Salmonella* infections and the food-animal link

1.2.1. Use of antibiotics and the growing burden of antibiotic resistance

Several national and global entities, such as World Health Organization (WHO), have identified antibiotic resistance as one of the major threats to human health, also linked to food safety via food ingestion, direct contact with animals or environmental spread (115, 116). This grim scenario results from the extraordinary adaptation of bacteria to selective pressures caused by the intensive use of antibiotics, mainly in non-human applications. In Europe, antibiotics are mainly used in livestock production for animal health in veterinary medicine as, since 2006, they were banned as growth promoters. In other regions, the real contribution of the use of antibiotics for animal health *versus* the use as growth promoters continues largely unknown (Table 2) (18, 117-119).

TABLE 2 – Antimicrobials approved for use in food-producing animals [adapted with permission from reference (120)].

Antimicrobial families/classes	Antimicrobial	USA		Europe		Portugal ^c	
		Used in C, P, S ^b	Available for growth promotion	Human use	Used in C, P, S ^b		Used in C, P, S ^b
Penicillins	Amoxicillin	C, P, S		X	C, S	X	C, P, S
	Ampicillin	C, P, S		X	C, S	X	C, P, S
	Cloxacillin	C		X	NL	X	C
	Penicillin (Procaine)	C, S	X	X	C, S	X	C, S
1 st generation cephalosporins ^a	Cephalexin	C, P, S		X	C, P, S	X	C
	Cefalonium	C, P			C, P		C
	Cephapirin	C			C		C
	Cefazolin	C, P		X	C, P	X	C
	Cefacetrile	C			C		NA
	Cefuroxime	C		X	C	X	NA
2 nd generation cephalosporins ^a	Cefoperazone	C, P		X	C, P	X	C
3 rd generation cephalosporins ^a	Ceftiofur	C, P, S			C, S		C, S
	Ceftriaxone	C, P, S		X	C, P, S	X	NA
4 th generation cephalosporins ^a	Cefquinome	C, P, S			C, S		C, S
	Danofloxacin	C			C		C
Quinolones	Enrofloxacin	C, S			C, P, S		C, P, S
	Apramycin	S	X		NL		C, P, S
Aminoglycosides	Gentamicin	C, P, S		X	P, S	X	C, P, S
	Neomycin	C, P, S	X	X	C, P, S	X	C, P, S
Tetracyclines	Chlortetracycline	C, P, S	X		C, P, S		C, P, S
	Oxytetracycline	C, P, S	X	X	C, P, S	X	C, P, S
	Tetracycline	C, P, S		X	C, P	X	C, P
	Oleandomycin	C			P, S	X	NA
Macrolides	Tilmicosin	P, S	X		NL		C, P, S
	Tylosin	C, P, S	X		C, S		C, P, S
Polypeptides	Erythromycin	C, P, S		X	C, P, S	X	C, S
	Bacitracin	C, P, S	X	X	Withdrawn	X	Withdrawn
Flavofosfolipid	Bambermycin	C, P, S	X		Withdrawn		Withdrawn
	Flavomycin	P	X		Withdrawn		Withdrawn
Quinoxaline	Carbadox	P, S	X		Withdrawn		Withdrawn

TABLE 2 - Continued

Antimicrobial families/classes	Antimicrobial	USA		Europe			Portugal ^c
		Used in C, P, S ^b	Available for growth promotion	Human use	Used in C, P, S ^b	Human use	
Polypeptides	Colistin/Polymyxin B	C, P	X	X	C, P, S	X	C, P, S
Phenicol	Florfenicol	C, S			C, S		C, S
Lincosamines	Lincomycin	C, P, S	X	X	C, P, S		C, P, S
	Pirlimycin	C			NL		C
Novobiocin	Novobiocin	C, P		X	C, P	X	NA
Aminocyclitol	Spectinomycin	C, P, S		X	NL	X	C, P, S
Diterpene	Tiamulin	S	X		P, S		P, S
Triamidine	Tulathromycin	C, S			NL		C, S
Streptogramin	Virginiamycin	P, S	X		Withdrawn		Withdrawn
Sulfonamides	Sulfadimethoxine	C, P, S	X		P		C, P, S
	Sulfamethazine	C, P, S	X		S		C, P

^a, Although extensively implemented in the past, FDA and EU have now controlled the off-label and unapproved use of cephalosporins (especially, extended-spectrum cephalosporins) in poultry, cattle and pigs (120).

^b, C, cattle; P, poultry; S, swine; NA, not available; NL, not listed in the references used by Seiffert et al (120).

^c, All the approved antimicrobials are considered veterinary medicinal products subject to medical prescription (121).

In fact, in 2012, the quantities of active substances of antimicrobials sold in 26 EU/European Economic Area (EEA) countries for animals (7982 tonnes or average 144.0 mg/kg), specially in chickens and pigs, were higher than that applied in humans (3400 tonnes or average 116.4 mg/kg) (122). Similarly, in the USA about 13.569,037 kilograms of antimicrobial active ingredient were sold in 2011 for food-animals comparing with the 3.289,176 kilograms reported for human medicine (123, 124). However, the greatest uncertainty about current antibiotic use in livestock is in the low- and middle-income regions/countries, such as Africa or the Asiatic continent, where a rise in antibiotic consumption is estimated (125, 126). A recent global study indicated that some of those countries (e.g. China, Brazil, Mexico and India) were the largest consumers of antibiotics for livestock, and after the application of statistical models a projection estimates that this will rise by two-thirds until 2030 (62.300 tonnes in 2010 to 105.600 tonnes in 2030), mainly due to an increase in livestock product consumption and the shift from extensive (i.e. small-scale) to large-scale intensive farming systems (i.e. industrial-scale) (125, 126).

The most frequently used antibiotic classes in food-producing animals were tetracycline's, sulphonamides, and penicillin's and, although 3rd-, 4th-generation cephalosporins and fluoroquinolones consumption was much lower for animals than for humans a small increase has been observed over the last years (18, 120, 122, 124, 127-129). Several classes of antimicrobials may not be fully metabolised by the human and animal bodies, and a large amount may be excreted in faeces or urine and enter wastewater treatment facilities (118). The slow rate of degradation of some antibiotics and the use of treated and untreated sewage as fertiliser can also lead to the accumulation of antimicrobials in diverse environments/hosts, including in food-animals and consequently in the food-chain production (Figure 5) (118). These subinhibitory concentrations may provide a constant selection pressure for maintenance of particular pathogenic bacteria, such as non-typhoidal *Salmonella* serotypes or clones carrying antibiotic resistance genes and other adaptive features associated with survival or improved host and environmental colonization (18, 118).




FIGURE 5 – Potential routes for dissemination of antimicrobial-resistant bacteria and resistance genes. Blue dotted arrows represent the selective pressures introduced by the use of antimicrobial agents [adapted from references (118, 130)].

Currently, the emergence and spread of *Salmonella* isolates presenting antimicrobial resistance (R) (see box 3 for details) or multidrug resistance¹ (MDR) profiles, especially to clinically relevant antibiotics for treating invasive infections, are of concern (3, 17). In fact, the raise of antimicrobial resistance represents a huge economic burden both within the EU and the USA (e.g. \$365.000.000 in medical costs per year) (79, 131, 132). Since resistance to older antibiotics (e.g. ampicillin, chloramphenicol and trimethoprim-sulfamethoxazole) has been increasing for many years, recommended treatment options for invasive infections included fluoroquinolones (ciprofloxacin) and extended-spectrum cephalosporins (3, 4). However, resistance to those “critically important antibiotics for human health” is emerging, leading to increase severity, morbidity and mortality of the disease and the need for the use of last-line antimicrobials (e.g. carbapenems) in therapy (18, 133). The contribution of food-animals for the global burden of antibiotic-resistant *Salmonella* serotypes/clones, particularly to β -lactams and quinolones are given in the next topics.

¹ Isolates resistant to at least one antibiotic are considered resistant (R) and to three or more antibiotics from different families are considered multidrug-resistant (MDR) (132).

BOX 3 - Antimicrobial resistance: concepts

Susceptibility of a microorganism to a particular antibiotic can be quantified by phenotypic methods such as the determination of the minimum inhibitory concentration (MIC), which is the lowest concentration of an antimicrobial agent that completely inhibits the growth of bacterial cells (117, 134, 135). The use of clinical breakpoints (based on clinical outcome data, MICs distributions, accepted dosing, pharmacokinetic/pharmacodynamic data) are extremely useful for clinicians to access the likelihood of therapeutic failure/success in human patients (135). Based on these clinical breakpoints, the isolate is defined as 1) resistant, 2) intermediate or 3) susceptible, accordingly to annual published guidelines and recommendations such as Clinical & Laboratory Standards Institute (CLSI - <http://clsi.org/>) or European Committee on Antimicrobial Susceptibility Testing (EUCAST - <http://mic.eucast.org/Eucast2/>). Contrarily the use of epidemiological cut-off values (ECOFFs) can be applied to distinguish between organisms without and with phenotypically expressed resistance mechanisms for a species and a drug in a defined test system. In this case microorganism can be categorized in wild type (WT-absence of acquired and mutational resistance mechanisms to the drug in question) and Non-Wild Type (NWT-presence of an acquired or mutational resistance mechanism to the drug in question). ECOFFs are defined as the MIC values that separate the upper limit of the wild type population from those isolates that have a phenotypically expressed resistance (Figure 6) (134, 135).




FIGURE 6 – Clinical and epidemiological breakpoints of susceptibility to antibiotics [adapted with permission from reference (134)].

1.2.2. The contribution of food-animals for the global burden of antibiotic-resistant *Salmonella*

For several decades, the contribution of food-animals as a reservoir of antimicrobial resistance with impacts in human health has been controversial, but accumulating evidences linking particularly the poultry and pig production with human disease have been reported, namely involving non-typhoidal *Salmonella* (136-138). The first evidence is the close association between the use of antimicrobial agents and the occurrence of resistance. In fact, as previously mentioned, regular use of antibiotics (mainly for animal health and in some countries as growth promoters), a practical associated with modern intensive food-animal production, has been considered the main

driver for the development of antibiotic resistance in zoonotic bacteria, such as *Salmonella* (120, 136, 138). For example, licensing of the fluoroquinolone enrofloxacin for animal use, especially in poultry, in the 1990s led to an increase of *S. Typhimurium* DT104 with decreased susceptibility to ciprofloxacin in animal, food and humans (7). On the other hand, an experimental study has shown that administration of tetracycline to pigs colonized with tetracycline-resistant *S. Typhimurium* DT104 was associated with elevated shedding of resistant *Salmonella* compared with untreated pigs (139). Some findings have also suggested a link between resistance to nitrofurans in human *Salmonella* and the food chain, suggesting that its illegal use in poultry industry might have contributed to the selection and persistence of *S. Enteritidis* in poultry, and consequently to human salmonellosis in Portugal (140). More recently, a voluntary withdrawal of ceftiofur by the poultry producers in Canada was correlated with a decrease in the occurrence of ceftiofur resistant *S. Heidelberg* (one of the most common serotypes associated with salmonellosis in this country) from both human infections and retail poultry, with an increase of the resistance levels after reintroduction of use (52).

Further evidence for the impact of food-animal production on human health problems associated with antimicrobial resistance in *Salmonella* is the correlation between different reservoirs (e.g. humans and poultry or pig) obtained from systematic surveillance data. In the last EU report, *Salmonella* resistant or MDR to commonly used antimicrobials (e.g. ampicillin-36.1%, sulphonamides-35.7% and tetracycline-34.5%) was frequently detected in humans (R=50%; MDR=31.8%) and animals, especially poultry [broilers (MDR=56%) and turkeys (MDR=73%), with 5-98% resistance to ampicillin, 5-85% to sulphonamides and 4-85% to tetracycline] and pigs (MDR=37.9%, with 72-91% resistance to ampicillin, sulphonamides and tetracycline), but also in derived meat products (18, 125). Also, in the USA, high levels of resistance and MDR in animal's/retail meat, mainly chicken (R=58.2%; MDR=9.1%/R=59.6%; MDR=26%), turkey (R=64.3%; MDR=28.6%/R=77.4%; MDR=39.6%) and pigs (R=25.6-33.5%; MDR=9-17.7%/R=45.8%; MDR=33.3%), and moderate levels in humans (R=19.2%; MDR=9.8%) were found in the integrated NARMS report (141).

Those data about MDR are extremely worrying due to the possible role of diverse antibiotics in the co-selection of *Salmonella* strains resistant to clinically relevant antibiotics, such as fluoroquinolones and extended-spectrum cephalosporins. In EU, relatively low levels of *Salmonella* non-susceptible ('clinically' resistant and 'intermediate' resistant categories combined) to ciprofloxacin (3.8%) and "microbiological" resistant (non-wild type by epidemiological cut-off values - ECOFFs) to cefotaxime (1.4%) were observed in humans (18). Moreover, the highest levels of "microbiological" resistance to these critically important antimicrobials were detected in broiler meat (68% ciprofloxacin

and 10.1% cefotaxime, with 0.3% of co-resistance to both antibiotics) and turkey meat (73.4% ciprofloxacin and 4.7% cefotaxime) and low levels in pig meat (3.9% ciprofloxacin and 0.9% cefotaxime), suggesting a greater contribution of the poultry production chain to the human burden (18). In fact, it was reported that the highest levels of resistance to ciprofloxacin were more common in *S. Enteritidis*, *S. Infantis* and *S. Kentucky*, three serotypes commonly associated with poultry meat (18). Also, in the USA, in spite of a decreasing trend since 2009, cephalosporin's-ceftriaxone resistance levels remain low in humans (2.5%) and in pig/pork (2.3-2.4%/0%), but still high in turkey/retail ground turkey meat (7.1%/9.4%) and chicken/retail chicken meat (7.3%/19.7%), especially in *S. Heidelberg* a poultry-associated serotype (15% in humans, 29% in retail ground turkey and 17.9% chicken) (141). In middle-income countries such as in the Asiatic continent, high levels of *Salmonella* non-susceptible to ciprofloxacin (15-48%) and cephalosporins (38% ceftriaxone) were observed in humans (142, 143). These data are in agreement with several studies that documented a high prevalence of resistance to fluoroquinolones (>22.5% ciprofloxacin) and cephalosporins (12.5->23.4% ceftriaxone and 26.6% ceftazidime) mainly in poultry meat from South Korea and China (44, 45). In these countries the high rates of antimicrobial resistance detected reflect the levels of antimicrobial consumption in livestock, including for growth promotion, an area that remains largely unregulated (45, 126, 143).

A third indication of the significant impact of antibiotic-resistant *Salmonella* (see box 4 for details) contaminating these foodstuffs on human health are the diverse reports demonstrating the same acquired antibiotic resistance mechanisms, particularly to the clinically important cephalosporins and fluoroquinolones, and/or the transmission of clinically relevant MDR *Salmonella* clones from food-animals to humans (17, 120, 137, 144).

BOX 4 - Antibiotic resistance mechanisms in *Salmonella*

Several antibiotic resistance mechanisms have been described, such as production of enzymes that inactivate antimicrobial agents through degradation or structural modification, reduction of bacterial cell permeability to antibiotics, activation of antimicrobial efflux pumps, and modification of cellular drug targets (145). Some of those mechanisms can be intrinsic (the innate ability of a bacterial species to resist/tolerate to the activity of a particular antimicrobial agent through its inherent structural or functional characteristics) to the bacteria or acquired via mutations in chromosomal genes and by horizontal gene transfer (Figure 7) (145, 146). As previously mentioned, *Salmonella* resistance to older antibiotics (e.g. chloramphenicol and trimethoprim-sulfamethoxazole) presented below, but even of greater concern to those clinically relevant (e.g. cephalosporins and fluoroquinolones) has become a public health issue.

- Phenicol modifying enzymes - Phenicol is a group of bacteriostatic compounds. Among them is chloramphenicol, a potent inhibitor of protein synthesis, able to prevent the formation of peptide bonds upon binding to the 50S ribosomal subunit (147). One of the resistance mechanisms is associated with drug target modification due to the production of acetyltransferases. An example may be that encoded by *cat* genes, which are able to acetylate both hydroxyl radicals of the chloramphenicol molecule, and according to its structure can be of type A or B (145). Phenicol resistance by efflux pumps results from the production of specific transport proteins such as those encoded by *floR* and *cmlA* genes, able to efflux florfenicol and chloramphenicol or only the second one, respectively (145).
- Resistance to trimethoprim and sulphonamides by modification of the cellular drug-binding target - These two agents inhibit the synthesis of nucleic acids by blocking the formation of purine and pyrimidine, interfering in consecutive stages of the folic acid biosynthetic pathway (147, 148). In *Salmonella* the genes responsible for sulphonamide resistance (*sul1*, *sul2* and *sul3*) encode altered dihydropteroate synthases enzymes that have a reduced affinity for the antibiotic. Likewise, trimethoprim resistance is possible due to production of alternative dihydrofolate reductases that have a reduced affinity for the antibiotic. To date, more than 30 genes identified in resistance to trimethoprim are divided according to their structure in *dhfrA* (e.g. *dhfrA12* and *dhfrA17* frequently reported in *Salmonella*) and *dhfrB* types (145, 148).




FIGURE 7 – Examples of antimicrobial resistance mechanisms in bacteria [adapted with permission from reference (149)].

1.2.2.1. Resistance to extended-spectrum β -lactams in *Salmonella*

β -lactams (see box 5 for details), particularly extended-spectrum cephalosporins (ESC), are the most widely used bactericidal antibiotics in clinical practice worldwide, including as a therapeutic option for human salmonellosis (18). However, over the last decade resistance to β -lactams has been extensively reported in *Enterobacteriaceae*, including in *Salmonella*, becoming a serious public health problem that requires full attention and appropriate management (18, 120). The most common mechanism of β -lactams resistance in *Salmonella* is through the production of enzymes, known as β -lactamases, capable of degrading the β -lactam ring, thereby inactivating their antimicrobial activity (145).

BOX 5 - Classification and mechanism of action of β -lactams

This class of antibiotics can be divided in four different groups: penicillin's, cephalosporins, carbapenems and monobactams. Some β -lactams have a narrow antimicrobial spectrum (penicillin's, first- and second-generation cephalosporins) being active mainly in Gram-positive bacteria, while others have a extended-spectrum (third-, fourth- and fifth-generation cephalosporins and carbapenems), acting on both Gram-negative and Gram-positive bacteria (150, 151). They are characterized by the presence of a β -lactam ring in their structure, acting by mimicking and thereby inactivating bacterial transpeptidases, such as penicillin binding proteins (PBPs), which are involved in the biosynthesis of peptidoglycan layer, necessary for cell wall formation (Figure 8) (147).




FIGURE 8 – β -lactams target interactions and associated cell death mechanisms [adapted with permission from reference (147)].

Due to their ability to hydrolyse critical important antibiotics in human and veterinary medicine, such as extended-spectrum β -lactams (especially third- and fourth-generation cephalosporins and carbapenems) three main important groups of β -lactamases are of special interest in *Salmonella*: extended-spectrum β -lactamases (ESBLs), AmpC β -lactamases and more recently carbapenemases (2, 120, 137, 152, 153).

The most frequent group, ESBLs², are generally acquired by horizontal gene transfer, arose by point mutations in the sequences of narrow-spectrum plasmid-mediated β -lactamase genes (e.g. *bla*_{TEM-1} and *bla*_{SHV-1}), and include derivatives of the TEM (Greek patient “Temoneira”) and SHV (“Sulphydryl reagent variable”) enzymes. The most prevalent group CTX-M (Cefotaximase Munich)-type enzymes have been associated with a mobilization process of chromosome-encoded *bla* genes from the environmental *Kluyvera* spp. (137, 150, 151, 154). A wide range of clinically important *Salmonella* serotypes (e.g. Enteritidis, Typhimurium, 4,[5],12:i:-, Heidelberg, Infantis, Kentucky, Virchow) mainly recovered from poultry and pigs (animal/food), have been associated with the worldwide dissemination of ESBLs (Table 3). By far, the most common genes found in poultry and/or poultry meat samples associated with resistance to extended-spectrum cephalosporins were those coding for CTX-M-1, CTX-M-2, CTX-M-9, CTX-M-15 and also TEM-52 enzymes (Table 3) (137). In pig and pork samples stands out the increasing report of CTX-M-15-producing *S. Enteritidis* and *S. Virchow* enzymes mainly in the Asiatic continent, although with a much lower frequency than that found in poultry (155). Also, the emergent *S. 4,[5],12:i:-* European clone has been recently identified as a producer of ESBLs such as CTX-M-1 enzymes mainly in UK and Germany, highlighting the role of this clonal lineage for the maintenance of this serotype in pigs (156, 157). Several studies have demonstrated the presence of some of those *Salmonella* clones carrying ESBLs in both human and food-animal isolates, suggesting an evidence of transmission along the food chain. For example, in several European countries, spread of *S. Virchow* producing CTX-M-2 or CTX-M-9 and *S. Infantis* producing TEM-52 have been found among poultry and humans (158-166).

The second most common group of β -lactamases in *Salmonella*, are the plasmid-encoded AmpC³ cephalosporinases, being enzymes less well reported than ESBLs (137, 167). Within this group the most prevalent enzyme is CMY-2, having the broadest geographic spread and frequently found in diverse *Salmonella* serotypes mainly recovered from poultry and poultry meat, but also pig and pork products (Table 3). Of concern is *S. Heidelberg*, one of the top MDR serotypes in both poultry and human (frequently associated with invasive infections) in North America, presenting increasing resistance to cephalosporins associated with *bla*_{CMY-2} (53, 168). Also MDR *S. Typhimurium*, the most

² The original definition of ESBLs is restricted to functional class 2be/molecular class A β -lactamases, being inhibited by clavulanic acid and possessing a hydrolytic activity against extended-spectrum cephalosporins (150). ESBLs can be classified into different families, according to their amino acid sequence (<http://www.lahey.org/studies/>).

³ AmpC β -lactamases (Ambler class C) hydrolyses efficiently penicillin's and oxyimino-cephalosporins (including cephamycins) and are not inhibited by clavulanic acid (167). These genes encoded originally for chromosomally (cAmpC) inducible (e.g. present in *Citrobacter* spp., *Enterobacter* spp. and *Serratia* spp.) and non-inducible enzymes (e.g. *E. coli* and *Shigella* spp.). Later, they were mobilized to transmissible plasmids (pAmpC) and nowadays have been reported in other Gram-negative bacteria usually poorly expressing or lacking chromosomal *bla*_{AmpC} genes, such as *Salmonella* (167).

prevalent pig-associated serotype and the second most common in human infections, has been increasingly associated with the AmpC *bla*_{CMY-2} mainly in pig and pork from non-EU countries but also from Belgium (169-171).

In addition to the ESBL/AmpC-producing *Salmonella*, production of carbapenemases (e.g. metallo- β -lactamase such as VIM or NDM) present a major challenge for antimicrobial chemotherapy, since these bacteria are usually resistant to almost all β -lactams (including carbapenems) and most of them are not inhibited by classical β -lactamase inhibitors (152, 153). Recently, plasmid-mediated carbapenem-resistant bacteria from food-animals have been reported, alerting for a new public health problem with indefinable risks (120, 153). Although the prevalence of acquired carbapenemases is still low in *Salmonella* (152), some studies reported carbapenemases in strains recovered from humans (e.g. *bla*_{KPC-2}, *bla*_{NDM-1}, *bla*_{OXA-48}, *bla*_{OXA-204}, *bla*_{VIM-1/-2}), with most of them arising from endemic regions (e.g. Algeria, Egypt, India, Libya, Tunisia) (153, 172-179), wild animals (e.g. *bla*_{NDM-1}) (180) and food-animals (181). One of the first reports on livestock farms were on two fattening pig farms and one broiler farm in Germany, where a *bla*_{VIM-1} gene (in an IncHI2 plasmid) was detected on *S. Infantis* strains from the environment and pig faeces (181, 182). In fact, this study demonstrates the ability of this clinically important serotype to be transmitted from farm environments to food-animals and potentially to humans via contaminated food (152, 153).

TABLE 3 - *Salmonella* serotypes/clones carrying extended-spectrum (ESBLs) and plasmid-encoded AmpC β -lactamases recovered from poultry, pigs and products thereof.

<i>Salmonella</i> serotype	ESBL or AmpC ^a enzyme (no. isolates)	Source	Concomitant presence in humans ^b	Country(ies)/year(s)	Non- β -lactam resistance phenotype ^c	Genetic element PL – Inc group	Reference (s)
Agona	CTX-M-1 (n=1)	Poultry (caecum)		The Netherlands/2006	SUL-TET	PL – I1	(165, 166)
	CTX-M-2 (n=2)	Turkey carcass		Brazil/2008	CIP-NAL-STR-SXT-TET	PL – P	(183)
		Poultry		Brazil/2011	NR	PL – NR	(184)
	CTX-M-8 (n=3)	Poultry		Brazil/2008, 2010	NR	PL – NR	(184)
	TEM-52 (n=3)	Poultry		Belgium/2001, 2004	(-)	PL – I1	(162)
	CMY-2 (n=5)	Turkey		Germany/2005	(-) or NAL	PL – I1	(185)
Anatum	CMY-2 (n=4)	Pig		USA/2008-2011	TET	PL – I1- γ	(186)
Bareilly	ACC-1 (n=6)	Poultry	Yes	The Netherlands/2001-02	(-)	PL – NR	(187)
Blockley	TEM-52 (n=6)	Poultry, poultry meat	Yes	The Netherlands/2001-02	(-) or SPT-SUL-TMP	PL – NR	(187)
Bovismorbificans	CTX-M-1 (n=1)	Pig faeces		UK/2009	SUL	PL – I1- γ	(157)
Brackenridge	CTX-M-14 (n=1)	Poultry		Brazil/2011	NR	PL – NR	(184)
Braenderup	ACC-1 (n=2)	Broiler		The Netherlands/2001	(-)	PL – NR	(187)
		Broilers		The Netherlands/2006	(-)	PL – NT	(165)
	SHV-12 (n=1)	Poultry		Italy/2006	(-)	PL – NR	(188)
Bredeney	CTX-M-1 (n=1)	Turkey		USA/2010	NR	PL – N	(189)
	CMY-2 (n=1)	Turkey		Canada/1999	SUL-TET	PL – A/C	(190)
Derby	CTX-M-1 (n=2)	Healthy pigs		Belgium/2009	NR	NR	(170)
	TEM-52(n=1)	Poultry		Belgium/2001	(-)	PL – I1	(162)
Enteritidis	CTX-M-2 (n=1)	Poultry		Brazil/2004	NR	NR	(184)
	CTX-M-9 (n=1)	Broiler		Spain/2000-04	STR-SUL-SXT-NAL-TET	NR	(160)
	CTX-M-14 (n=5)	Imported chicken meat		China/2004	NAL	PL – NR	(191)
		Poultry		China/2012-13	CHL-CIP-ENR-FFC-NAL-OLA-SXT-(AMK-GEN-LVX-TET)	PL – HI2, F, N or B/O	(192)
	CTX-M-15 (n=36)	Chicken meat, faeces or disease	Yes	Korea/2009	STR-SUL-TET-(GEN-NEO)	PL – FII	(193)
		Chicken		South Korea/2009-10	NR or GEN-STR-SUL-TET	PL – FII	(194)
		Pigs		Korea/2012-13	GEN-NAL-NEO-STR-	HI2	(155)

TABLE 3 - Continued

<i>Salmonella</i> serotype	ESBL or AmpC ^a enzyme (no. isolates)	Source	Concomitant presence in humans ^b	Country(ies)/year(s)	Non-β-lactam resistance phenotype ^c	Genetic element PL – Inc group	Reference (s)
Emek Essen	SHV-12 (n=2)	Poultry		Italy/2006	TET	NR	(188)
	CMY-2 (n=1)	Broiler		Portugal/2012-13	CHL-CIP-SUL-NAL-TET	NR	(195)
	CTX-M-15 (n=1)	Poultry		Brazil/2011	NR	NR	(184)
		Diseased chicken		Korea/2009	GEN-STR-SUL-TET	PL – FII	(193)
Gaminara	CTX-M-14 (n=1)	Poultry		Brazil/2009	NR	NR	(184)
Give	CTX-M-2 (n=1)	Poultry		Brazil/2009	NR	NR	(184)
Hadar	CTX-M-14 (n=1)	Poultry		Brazil/2005	NR	NR	(184)
Havana	CTX-M-1 (n=1)	Broiler		Portugal/2012-13	SUL-TET	NR	(195)
	CMY-2 (n=2)	Broiler		Portugal/2012-13	(-)	NR	(195)
Heidelberg	CTX-M-1 (n=1)	Broiler carcass		Portugal/2012-13	SUL-TMP	NR	(195)
	CTX-M-2 (n=14)	Chicken		Venezuela/2005-07, 2008	CIP-GEN	PL – NR	(196)
		Poultry		Brazil/2009, 2011	NR	NR	(184)
	CTX-M-14 (n=1)	Poultry		Brazil/2011	NR	NR	(184)
	SHV-2 (n=1)	Retail chicken meat		Canada/2007	GEN-STR-SUL	PL – I1	(197)
	CMY-2 (n=47)	Chicken or chicken retail	Yes	Canada/2001-04	CHL-GEN-KAN-STR- SUL-SXT-TET	PL – A/C, I1	(168)
Indiana		Porcine		Canada/1999-2007	(CHL-SUL-TET-SXT)	PL – A/C, I1	(169)
		Poultry		Canada/1999-2007	(CHL-KAN-SUL-TET- SXT)	PL – A/C, I1	(169)
		Turkey, chicken		USA/2007	(-) or KAN-TET-(CHL- GEN-STR-SUL)	PL – I1, A/C, N	(198)
		Chicken food		USA/2002, 2004, 2010	(CHL-GEN-KAN-STR- SUL-TET)	PL – A/C, I1	(53)
	CTX-M-14 (n=22)	Poultry		Brazil/2011	NR	NR	(184)
		Poultry		China/2012-13	ENR-NAL-SXT-(AMK- CHL-CIP-FFC-GEN- LVX-OLA-TET)	PL – HI2, A/C, F, PL – N, P-1α, B/O	(192)
		Duck		China/2009-10	CIP-NAL	PL – N	(199)
	CTX-M-24 (n=29)	Chicken		China/2008-09	AMK-CHL-FFC-GEN- NAL-OLA-RIF-STR- SXT-(TET)	PL – HI2	(200)
	CTX-M-27 (n=3) CTX-M-65 (n=17)	Duck Poultry		China/2009-10 China/2012-13	CIP-NAL CHL-NAL-OLA-SXT- (CIP-ENR-FFC-GEN-	PL – HI2, N PL – HI2, F, N	(199) (192)

TABLE 3 - Continued

<i>Salmonella</i> serotype	ESBL or AmpC ^a enzyme (no. isolates)	Source	Concomitant presence in humans ^b	Country(ies)/year(s)	Non-β-lactam resistance phenotype ^c	Genetic element PL – Inc group	Reference (s)
Infantis		Chicken		China/2012	LXX-TET)	PL – NR	(201)
		Chicken		China/2011	GEN-AZI CHL-CIP-GEN-NAL- SXT-TET	PL – NR	(202)
	TEM-52 (n=1) DHA-1 (n=3)	Pork Pig		China/2012 China/2011	GEN-AZI CHL-CIP-GEN-NAL- SXT-TET	PL – NR PL – NR	(201) (202)
		Broilers		The Netherlands/2006	(-)	PL – NT	(165)
		Chicken faeces		South Korea/2006-07	SUL-TMP-(APM-NAL- NEO-STR-TET)	PL – NR	(203)
	CTX-M-1 (n=31)	Poultry (caecum or unknown)	Yes	The Netherlands/2006	SUL-TMP	PL – I1	(165, 166)
	CTX-M-2 (n=3)	Broiler chicken, broiler meat		Italy/2012-14	NAL-CIP-SUL-TET-TRI- (GEN-KAN-STR)	PL – P (pESI-like)	(204)
		Poultry		Brazil/2005	NR	NR	(184)
		Retail Chicken Products		Japan/2002-03	(-)	NR	(205)
	TEM-20 (n=2)	Retail Chicken Products	Yes	Japan/1997, 2003	(-)	NR	(205)
	TEM-52 (n=14)	Poultry		Belgium/2001-05	(-)	PL – I1	(162)
		Broiler		Belgium/2004	(-)	PL – I1	(163)
		Poultry (caecum or unknown)		The Netherlands/2006	(-)	PL – I1	(165, 166)
	CMY-2 (n=50)	Chicken		Japan/2004-06	(-)	PL – NR	(206)
		Retail Chicken Products		Japan/2000, 2003	(-)	NR	(205)
Kentucky	CTX-M- 25/OXA-21 (n=1) SHV-12 (n=3)	Retail Chicken Products		Japan/1997, 1999-03	(-)	PL – NR	(205)
		Healthy pigs		Japan/2007-08	CHL-STR-SUL-TET	PL – NR	(207)
		Turkey		Poland/2009	CIP-NAL	PL – A/C	(64)
		Whole chicken and chicken neck skin		Ireland/2008-09	CHL-SUL-TET	PL – NR	(208)

TABLE 3 - Continued

<i>Salmonella</i> serotype	ESBL or Amp ^{Ca} enzyme (no. isolates)	Source	Concomitant presence in humans ^b	Country(ies)/year(s)	Non- β -lactam resistance phenotype ^c	Genetic element PL – Inc group	Reference (s)
	CMY-2 (n=11)	Chicken meat Whole chicken Poultry		Germany/2005 Ireland/2009 Canada/1999-2007	STR (-) (-)	PL – I1 PL – NR PL – I1	(185) (208) (169)
Kiambu	SHV-2 (n=1)	Porcine Abattoir chicken cecum		Canada/1999-2007 Canada/2006	(-) STR-SUL-SXT	PL – I1 PL – I1	(169) (197)
Livingstone	SHV-12 (n=9)	Poultry or turkey carcass, poultry faeces, broiler faeces		Italy/2005-06	GEN-NAL-(STR-SUL)	NR	(188)
Llandoff	CTX-M-1 (n=1)	Poultry		France/2006	SUL-TET	PL – I1	(209)
Manhattan	CTX-M-2 (n=2)	Retail Chicken Products		Japan/2000, 2003	(-)	NR	(205)
	CTX-M-15 (n=4)	Retail Chicken Products		Japan/2002, 2003	(-)	NR	(205)
	TEM-52 (n=4)	Retail Chicken Products		Japan/2000, 2002-03	(-)	NR	(205)
	SHV-12 (n=2)	Retail Chicken Products		Japan/2002, 2003	(-)	NR	(205)
	CMY-2 (n=2)	Retail Chicken Products		Japan/2002	(-)	PL – NR	(205)
Minnesota	CTX-M-2 (n=1)	Poultry		Brazil/2009	NR	NR	(184)
	CTX-M-8 (n=1)	Poultry		Brazil/2008	NR	NR	(184)
	CTX-M-14 (n=1)	Poultry		Brazil/2010	NR	NR	(184)
	CMY-2 (n=2)	Poultry		Brazil/2011	NR	NR	(184)
Newport	CTX-M-2 (n=1)	Poultry		Brazil/2008	NR	NR	(184)
Ouakam	CTX-M-1 (n=6)	Turkey		USA/2011	NR	PL – N	(189)
Panama	CTX-M-1 (n=1)	Poultry		Brazil/2009	NR	NR	(184)
Poona	CTX-M-2 (n=1)	Poultry		Brazil/2009	NR	NR	(184)
	CTX-M-8+CTX-M-14 (n=1)	Poultry		Brazil/2005	NR	NR	(184)
Rissen	CTX-M-1 (n=1)	Healthy pigs		Belgium/2009	NR	NR	(170)
	SHV-12 (n=1)	Pig		Spain/1999	STR-SUL-TET	NR	(160)
Saintpaul	CMY-2 (n=1)	Poultry		Brazil/2005	NR	NR	(184)
Schwarzengrund	CTX-M-2 (n=1)	Poultry		Brazil/2011	NR	NR	(184)
	CTX-M-2 (n=10)	Turkey, chicken, chicken carcass		Brazil/2008	CIP-NAL-STR-TET	PL – P	(183)

TABLE 3 - Continued

<i>Salmonella</i> serotype	ESBL or AmpC ^a enzyme (no. isolates)	Source	Concomitant presence in humans ^b	Country(ies)/year(s)	Non-β-lactam resistance phenotype ^c	Genetic element PL – Inc group	Reference (s)
Senftenberg	CTX-M-15 (n=1)	Chicken		South Korea/2009	NR	PL – FII	(194)
	CTX-M-1 (n=2)	Poultry meat		Germany/2006	KAN-NEO-STR-SUL-SXT-TET-TMP	PL – I1	(185)
Typhimurium		Swine meat	Yes	Germany/2007	KAN-NEO-STR-SUL-TET-TMP-SXT	PL – N	(185)
	CTX-M-2 (n=10)	Poultry		Brazil/2004-05, 2007	NR	NR	(184)
		Poultry	Yes	Brazil/2003-04	SUL-SXT-TET	PL – NR	(210)
	CTX-M-8 (n=1)	Poultry		Brazil/2011	NR	NR	(184)
	CTX-M-14 (n=1)	Poultry		Brazil/2005	NR	NR	(184)
	CTX-M-15 (n=1)	Broiler		Belgium/2007	(-)	PL – I1	(163)
	TEM-52 (n=6)	Poultry		The Netherlands/2001-02	SPT-SUL-TMP-(NEO-TET)	NR	(187)
		Poultry	Yes	Belgium/2002, France/2004	(-)	PL – I1	(162)
		Healthy pigs		Belgium/2009	NR	NR	(170)
	TEM-52+SHV-12 (n=1)	Poultry meat		The Netherlands/2002	(-)	NR	(211)
	CMY-2 (n=65)	Food from pig Pork and pig		Colombia/UN Mexico/2002-05	OT CHL-STR-SUL-TET-(GEN-KAN-NAL-SXT)	NR NR	(212) (213)
		Retail chicken	Yes	Mexico/2002-04	CHL-STR-SUL-TET-(GEN-KAN-NAL-SXT)	NR	(213)
		Porcine Swine		Canada/1999-2007 USA/2007	CHL-SUL-TET-(GEN) CHL-GEN-KAN-STR-SUL-SXT-TET	PL – A/C PL – A/C, FIB, I1	(169) (198)
		Chicken		USA/2007	STR-SUL-TET	PL – A/C, I1	(198)
Virchow		Healthy pigs Diarrheic pig		Belgium/2009 South Korea/2011-12	NR CHL-FFC-GEN-NAL-SXT-TET	NR PL – A/C, FIB	(170) (171)
	CTX-M-2 (n=12)	Poultry or poultry product		Belgium/2000-01, 2003	NAL-SUL-TET-TMP-(STR)	PL – HI2	(161)
		Poultry		Ireland/UN	SUL-TET-TMP	PL – P	(190)
		Broiler		Belgium/2001, 2004	STR-SUL-TET-TMP	PL – HI2	(163)
		Poultry	Yes	Belgium-France/2000-03	NAL-SUL-TMP-(STR-TET)	PL – NR	(159)
		Broiler		The Netherlands/2002	NAL-SPT-SUL-TET-TMP	NR	(187)
	CTX-M-2+SHV-2+TEM-1 (n=1)						
	CTX-M-9 (n=8)	Chicken faeces		France/2002-03	KAN-NAL-SPT-STR-	PL – NR	(158)

TABLE 3 - Continued

<i>Salmonella</i> serotype	ESBL or AmpC ^a enzyme (no. isolates)	Source	Concomitant presence in humans ^b	Country(ies)/year(s)	Non-β-lactam resistance phenotype ^c	Genetic element PL – Inc group	Reference (s)
		Chicken faeces and retail chicken meat Broiler	Yes	France/2002-03, Spain/2000	SUL-TET-TMP NAL-SPT-STR-SUL- TET-TMP-(KAN)	PL – HI2	(161)
			Yes	Spain/2000-04	STR-SUL-SXT-NAL- (TET)	NR	(160, 164)
	CTX-M-15 (n=5)	Pigs		Korea/2012-13	GEN-NAL-NEO-STR- TET	HI2	(155)
	CTX-M-32 (n=2)	Poultry		Greece/2001	CHL-KAN-STR-SUL- TET-TMP	PL – NR	(214)
Weslaco	TEM-52 (n=1)	Poultry		The Netherlands/2002	SPT-STR-SUL-TMP	NR	(187)
35:c:1,2	CTX-M-8 (n=1)	Poultry		Brazil/2009	NR	NR	(184)
4:[5],12:ii:-	SHV-12 (n=5)	Poultry	Yes	Senegal/2000	CHL-GEN-TET-TOB	NR	(215)
	CTX-M-1 (n=15)	Pig faeces		UK/2009	SUL-(CHL)	PL – I1-γ	(157)
		Swine		Germany/2007, 2009-10	SUL-TET-(STR)	PL – N, I1,	(156)
	CTX-M-2 (n=1)	Avian		Germany/2010	GEN-NAL-SPT-STR- SUL-TET	PL – NT	(156)
	CMY-2 (n=3)	Poultry		Canada/1999-2007	(-)	PL – I1	(169)
		Chicken		USA/2007	KAN-SUL-TET	PL – A/C, N	(198)

Abbreviations: AMK, amikacin; APM, apramycin; AZI, azithromycin; CHL, chloramphenicol; CIP, ciprofloxacin; ENR, enrofloxacin; FFC, florfenicol; GEN, gentamicin; KAN, kanamycin; LVX, levofloxacin; NAL, nalidixic acid; NEO, neomycin; OLA, olaquinox; OT, oxytetracycline; RIF, rifampicin; SPT, spectinomycin; STR, streptomycin; SUL, sulphonamides compound; SXT, trimethoprim/sulfamethoxazole; TET, tetracycline; TMP, trimethoprim; TOB, tobramycin; NR, not reported; NT, plasmids that were not typeable with the scheme used; PL, plasmid. Shaded grey lines represent *Salmonella* isolates from pig and/or pork.

^a, Only references with full-characterized ESBL/AmpC genes were considered.

^b, “Yes” – Clones or serotypes simultaneously detected in poultry or pigs, and humans in the same study.

^c, (-), The non-β-lactam resistance phenotype was not detected using the tested antibiotics and the susceptibility criteria adopted. Variable phenotypes were present between curved brackets.

1.2.2.2. Resistance to quinolones in *Salmonella*

Besides the worldwide resistance to clinically important β -lactams, resistance mechanisms that confer decreased susceptibility to fluoroquinolones (see box 6 for details) such as ciprofloxacin, one of the most important antibiotics for treating salmonellosis, have been widely reported in *Salmonella* from humans and food-animals (144, 145).

BOX 6 - Classification and mechanisms of action of quinolones

First generation quinolones (e.g. nalidixic acid) and 2nd to 4th generation fluoroquinolones (e.g. ciprofloxacin, levofloxacin and gemifloxacin, respectively) are bactericidal antibiotics. They inhibit DNA synthesis in bacterial cells by blocking type II topoisomerases (DNA gyrase and topoisomerase IV) at the DNA cleavage stage, preventing strand rejoining and consequently leading to cell death (Figure 9) (144, 147). Type II topoisomerases are tetrameric enzymes, with two A and two B subunits. The A subunits of DNA gyrase and topoisomerase IV are encoded by the *gyrA* and *parC* genes, respectively, while the B subunits are encoded by *gyrB* and *parE* genes, respectively (144).




FIGURE 9 – Quinolones target interactions and associated cell death mechanisms [adapted with permission from reference (147)].

In *Salmonella*, the most frequent mechanism of quinolone resistance has been mutations in the quinolone-resistance determining region (QRDR) of *gyrA* (DNA gyrase) and/or *parC* (topoisomerase IV) chromosomal genes (144, 145). Moreover, plasmid-mediated resistance determinants to fluoroquinolones have been described, which typically confer decreased susceptibility to ciprofloxacin, designated plasmid-mediated quinolone resistance (PMQRs)⁴. In both cases, the final level of quinolone resistance displayed is related to the number of resistance mechanisms that are present, either mutations in the organism under consideration and/or the concomitant presence with

⁴ The mechanisms of plasmid-mediated quinolone resistance so far described are: i) protection of target enzymes by Qnr proteins blocking their inhibition (e.g. QnrA, QnrB, QnrC, QnrD and QnrS). All the *qnr* gene variants can be accessed online at <http://www.lahey.org/qnrstudies/>; ii) production of acetylases that affect the activity of some fluoroquinolones and aminoglycosides [e.g. AAC(6')-Ib-cr]; and iii) efflux systems that pump fluoroquinolones out of the bacterial cell (e.g. QepA - major facilitator superfamily and OqxAB - resistance nodulation division family) (144).

additional mechanisms such as PMQRs genes [e.g. *qnr*, *aac(6')-Ib-cr*, *qepA* and *oqxAB* genes] (216).

Among these PMQR mechanisms, Qnr proteins have been commonly described in different clinically relevant *Salmonella* serotypes and geographic locations (Table 4). In fact, in poultry-associated serotypes such as *S. Enteritidis*, *S. Heidelberg*, *S. Infantis* or *S. Kentucky* several Qnr proteins (e.g. QnrB2, QnrB19, QnrS1) have been frequently reported (Table 4). As can be seen in an European study from Veldman et al (217), poultry seems to be one of the most important vehicles of non-typhoidal *Salmonella* carrying PMQR-*qnr* genes, alerting for the role of food-animals, including animal/food trade, in its dissemination. Nevertheless, an incidence of OqxAB efflux pump and/or the aminoglycoside acetyltransferase AAC(6')-Ib-cr enzymes in pig-associated serotypes such as *S. Typhimurium* and *S. Derby*, recovered from Asiatic pig and pork, have also been recently found (201, 218). In fact, the increasing presence of OqxAB in Asia could be mainly attributed to the widespread use of olaquinox as growth promoter in pig industry (201).

More worrying is that although still rare in *Salmonella* from food-animals, there have been recently some reports displaying an association of PMQR genes with other plasmid-mediated *bla*_{CTX-M} genes (144, 219). For example, these associations have been found in IncHI2 plasmids of several MDR *Salmonella* serotypes such as Indiana (*oqxAB±aac(6')-Ib-cr±bla*_{CTX-M-14, -24, -27 or -65}) and Enteritidis [*oqxAB±aac(6')-Ib-cr+bla*_{CTX-M-14}] mainly recovered from chickens (192, 199-202). These studies highlight the possibility of clinical treatment failure and the additional co-selection potential by two critically important antibiotics, quinolones and β-lactams.

The increasing prevalence of ESBLs, AmpC β-lactamases and/or PMQR genes in MDR *Salmonella* from food-animals and humans is demonstrated as being in a close association with highly adapted clones and/or favoured by horizontal transfer of mobile genetic elements (e.g. plasmids, transposons, genomic islands), two issues that will be addressed in the next topic.

TABLE 4 – *Salmonella* serotypes/clones carrying plasmid-mediated quinolone resistance (PMQR) genes recovered from poultry, pigs and products thereof.

<i>Salmonella</i> serotype	PMQR ^a mechanism (no. Isolates)	Source ^b	Country(ies)/year(s)	Antibiotic resistance phenotype ^c	Genetic element PL – Inc group	Reference(s)
Agona	QnrB2 (n=1)	Turkeys	Germany/NR	NR	NR	(217)
Braenderup	QnrD (n=2)	Fowls	Spain/NR	NR	NR	(217)
Concord	QnrB2 (n=1)	Pigs	Czech Republic/NR	NR	NR	(217)
Dabou	QnrD (n=1)	Fowls	Spain/NR	NR	NR	(217)
Derby	QnrB2 (n=56)	Fowls or turkeys	Spain/NR	NR	NR	(217)
	QnrS1 (n=1)	Chicken	China/2013	AMP-CHL-CIP-GEN-KAN-NAL-OLA-STR-SUL-TET	NR	(201)
	OqxAB + QnrS8 + QnrB (n=1)	Chicken	China/2013	AMP-CHL-CIP-KAN-NAL-OLA-STR-SUL-TET	NR	(201)
	OqxAB (n=1)	Pork	China/2013	AMP-CHL-CIP-GEN-KAN-NAL-OLA-STR-SUL-TET	NR	(201)
	OqxAB + QnrS1 + QnrB (n=1)	Pork	China/2013	AMP-CHL-CIP-GEN-KAN-NAL-OLA-STR-SUL-TET-(AZI)	NR	(201)
	OqxAB + AAC(6')-Ib-cr + QnrS2 (n=11)	Pork	China/2012-13	AMP-CHL-CIP-GEN-KAN-NAL-OLA-STR-SUL-TET-(AZI)	NR	(201)
	OqxAB + AAC(6')-Ib-cr + QnrS2 + QnrB8 (n=1)	Pork	China/2012	AMP-CHL-CIP-GEN-KAN-NAL-OLA-STR-SUL-TET	NR	(201)
Enteritidis	QnrB2 (n=1)	Fowls or turkeys	Spain/NR	NR	NR	(217)
	QnrB10/B19 (n=1)	Laying hen flock	Poland/2009	CIP-NAL	NR	(64)
	QnrD (n=3)	Fowls	Spain/NR	NR	NR	(217)
	QnrS1/S3 (n=3)	Broiler meat, broiler flock - faeces	Poland/2008-09	AMP-CIP-(STR-TET)	NR	(220)
		Swine flock - faeces	Poland/2008	AMP-CIP	NR	(220)
	OqxAB (n=3)	Chickens	China/2012-13	AMP-CHL-CIP-CIP-ENR-FFC-NAL-OLA-SXT-(GEN-LVX-TET)	PL - HI2, F, N, B/O	(192)
	OqxAB + AAC(6')-Ib-cr (n=1)	Chickens	China/2012-13	AMK-AMP-CHL-CIP-CTX-ENR-FFC-GEN-LVX-NAL-OLA	PL - HI2	(192)

TABLE 4 - Continued

<i>Salmonella</i> serotype	PMRQ ^a mechanism (no. Isolates)	Source ^b	Country(ies)/year(s)	Antibiotic resistance phenotype ^c	Genetic element PL – Inc group	Reference(s)
Give	QnrB19 (n=1)	Imported turkey meat from Brazil	Finland/NR	SXT-TET	NR	(217)
Goldcoast	QnrS1 (n=1)	Pigs	Belgium/NR	NR	NR	(217)
Hadar	QnrB2 (n=2)	Fowls or turkeys	Spain/NR	NR	NR	(217)
	QnrB5 (n=4)	Imported turkey meat	Germany/2007	STR-TET	NR	(221)
	QnrB19 (n=15)	Fowls, turkeys	Germany, Denmark/NR	NR	NR	(217)
Havana	QnrB2 (n=3)	Poultry	Portugal/2009-10	(-) or NAL	PL – L/M	(222)
	QnrB19 (n=1)	Poultry	Portugal/2009-10	(-)	PL – HI2	(222)
Heidelberg	QnrB19 (n=2)	Chicken	Venezuela/2005-07, 2008	CIP-GEN	PL – NR	(196)
Indiana	OqxAB (n=52)	Chickens	China/2012-13	AMP-CIF-NAL-SXT-(AMK-CAZ-CHL-CIP-CTX-ENR-FFC-GEN-LVX-OLA-TET)	PL – HI2, F, N, P-PL – 1α, B/O	(192)
		Chicken	China/2008-09	AMK-CHL-FFC-GEN-NAL-OLA-RIF-STR-SXT-(TET)	PL – HI2	(200)
		Duck	China/2009-10	AMP-CAZ-CIP-CRO-CTX-NAL	PL – HI2, N	(199)
	OqxAB + AAC(6')-Ib-cr (n=18)	Chickens	China/2012-13	AMP-CHL-CIF-NAL-SXT-(CIP-CTX-ENR-FFC-GEN-LVX-OLA-TET)	PL – HI2, A/C, N	(192)
		Duck	China/2009-10	AMP-CAZ-CIP-CRO-CTX-NAL	PL – N	(199)
		Chicken	China/2011	AMP-CHL-CIP-GEN-NAL-SXT-TET-(CAZ-CTX)	PL – NR	(202)
		Pig	China/2011	AMP-CAZ-CHL-CIP-CTX-GEN-NAL-SXT-TET	PL – NR	(202)
		Pork	China/2012	AMP-AZI-CHL-CIP-CRO-GEN-KAN- NAL-OLA-STR-SUL-TET	NR	(201)
	AAC(6')-Ib-cr (n=5)	Chicken	China/2011	AMP-CHL-CIP-GEN-NAL-SXT-TET-(CAZ-CTX)	PL – NR	(202)
		Duck	China/2009-10	AMP-CAZ-CIP-CRO-CTX-NAL	PL – N	(199)
	OqxAB + AAC(6')-Ib-cr + QnrB (n=1)	Chicken	China/2012	AMP-AZI-CHL-CIP-CRO-GEN-KAN- NAL-OLA-STR-SUL-TET	NR	(201)

TABLE 4 - Continued

<i>Salmonella</i> serotype	PMRQ ^a mechanism (no. Isolates)	Source ^b	Country(ies)/year(s)	Antibiotic resistance phenotype ^c	Genetic element PL – Inc group	Reference(s)
Infantis	QnrS1 (n=1)	Chicken	Germany/2004	AMP	NR	(223)
	QnrB19 (n=1)	Retail chicken	Colombia/2004	KAN-NAL-NEO-STR-TET	PL – ColE like	(224)
Kentucky	QnrS1 (n=1)	Chicken	The Netherlands/NR	NR	PL – N	(225)
London	QnrB2 (n=22)	Fowls or turkeys	Spain/NR	NR	NR	(217)
Mbandaka	QnrB19 (n=1)	Poultry	Portugal/2009-10	NAL	PL – HI2	(222)
Montevideo	QnrB2 (n=3)	Fowls or turkeys	Spain/NR	NR	NR	(217)
	QnrD (n=6)	Fowls, turkeys	Italy/NR	NR	NR	(217)
Newport	QnrB5 (n=3)	Imported turkey meat	Poland/2007	(-)	NR	(221)
	QnrS1/S3 (n=12)	Broiler (meat, flock), turkey meat, goose flock - faeces, duck flock - faeces	Poland/2008-11	CIP-(AMP-CHL-FFC-KAN-STR-SUL-TET-NAL)	NR	(64)
Ohio	QnrB19 (n=3)	Turkeys	Denmark/NR	NR	NR	(217)
	QnrD (n=5)	Fowls	Spain/NR	NR	NR	(217)
Rissen	QnrS1 (n=1)	Pig	Korea/2012-13	NAL	NR	(155)
	OqxAB (n=1)	Pork	China/2013	AMP-CHL-CIP-KAN-NAL-OLA-STR-SUL-TET	NR	(201)
Saintpaul	QnrS1 (n=15)	Imported turkey meat	Germany/2007-08	STR-TET-(CPD-CHL)	NR	(221)
		Turkeys	Poland/2008	NR	NR	(217)
			Germany, Denmark/NR	NR	NR	(217)
Typhimurium	QnrA1 (n=1)	Turkeys	Germany/NR	NR	NR	(217)
	QnrD (n=2)	Fowls	Spain/NR	NR	NR	(217)
	OqxAB (n=7)	Chicken	China/2007-09	FFC-OLA-SMX-(AMP-CHL-CIP-GEN-NAL-TET)	PL – HI2, F	(218)
		Pigs	China/2010	AMP-CHL-FFC-NAL-OLA-SLX-TET-(GEN)	PL – F	(218)
		Pork	China/2012-13	AMP-CHL-CIP-KAN-NAL-OLA-STR-SUL-TET-(GEN)	NR	(201)

TABLE 4 - Continued

<i>Salmonella</i> serotype	PMQR ^a mechanism (no. Isolates)	Source ^b	Country(ies)/year(s)	Antibiotic resistance phenotype ^c	Genetic element PL – Inc group	Reference(s)
Virchow	OqxAB + AAC(6')-Ib-cr (n=16)	Chicken, duck	China/2009-10	NAL-OLA-SMX-(AMP-CHL-CIF-ENR-FFC-GEN-TET)	PL – HI2	(218)
		Pigs	China/2008-10	AMP-NAL-OLA-SMX-TET-(CHL-CIF-FFC-GEN)	PL – HI2	(218)
		Pork	China/2012	AMP-CHL-CIP-GEN-KAN-NAL-OLA-STR-SUL-TET	NR	(201)
	OqxAB + AAC(6')-Ib-cr + QnrS1 (n=4)	Pork	China/2012-13	AMP-CHL-CIP-GEN-KAN-NAL-OLA-STR-SUL-TET	NR	(201)
	AAC(6')-Ib-cr + QnrB (n=1)	Chicken	China/2013	AMP-CHL-CIP-GEN-KAN-NAL-OLA-STR-SUL-TET	NR	(201)
	OqxAB + AAC(6')-Ib-cr + QnrS1 (n=1)	Duck	China/2010	AMP-CHL-CIF-FFC-GEN-NAL-OLA-SMX-TET	PL – HI2	(218)
	QnrS1	Chicken	UK/2004-05	AMP-CIP	N	(226)
		Chicken	Turkey/2005	AMP-NAL	PL – NT	(227)
		Chicken carcass	Korea/2002	AMP-CEF-STR-SUL-TET-TMP	N	(228)
	OqxAB + AAC(6')-Ib-cr + QnrS8 + QnrD (n=1)	Pork	China/2013	AMP-CHL-CIP-GEN-KAN-NAL-OLA-STR-SUL-TET	NR	(201)

Abbreviations: AMC, amoxicillin/clavulanate; AMK, amikacin; AMP, ampicillin; AZI, azithromycin; CAZ, ceftazidime; CEF, cefalotin; CHL, chloramphenicol; CIF, ceftiofur; CIP, ciprofloxacin; CPD, cefpodoxime; CRO, ceftriaxone; CTX, cefotaxime; ENR, enrofloxacin; FFC, florfenicol; GEN, gentamicin; KAN, kanamycin; LVX, levofloxacin; NAL, nalidixic acid; NEO, neomycin; OLA, olaquinox; RIF, rifampicin; STR, streptomycin; SUL, sulphonamides compound; SXT, trimethoprim/sulfamethoxazole; TET, tetracycline; TMP, trimethoprim; NR, not reported; NT, plasmids that were not typeable with the scheme used; PL, plasmid. Shaded grey lines represent *Salmonella* isolates from pig and/or pork.

^a, Only references with full-characterized PMQR genes were considered.

^b, No studies were found simultaneously reporting the same *Salmonella* clones or serotypes both in poultry/pigs and humans in the same study.

^c, (-), Antibiotic resistance phenotype was not detected using the tested antibiotics and according to the susceptibility criteria adopted. Variable phenotypes were present between curved brackets.

1.2.3. Acquisition and dissemination of antimicrobial resistance in *Salmonella*

The long-term persistence and dissemination of bacteria in antimicrobial selective environments, including the establishment of well-adapted *Salmonella* serotypes/clones⁵, is in part favoured by the vertical (e.g. clonal spread) or horizontal acquisition of genetic units (e.g. plasmids, transposons, integrons, genomic islands) (Figure 10) carrying genes encoding for several adaptive features (e.g. antibiotic resistance and others such as metal/biocide tolerance or virulence) (118). Additionally, the enormous ability of bacteria to undergo different events such as mutations or recombination with foreign DNA, are also major contributors for the evolution of antimicrobial resistant bacterial populations (Figure 10) (118).




FIGURE 10 – Genetic events responsible for horizontal and vertical transmission of genes that can favour the selection of antibiotic-resistant bacteria. Small filled red circles represent antibiotic-resistant bacteria; Small filled blue circles represent antibiotic-susceptible bacteria; Black lines represent vertical transmission to daughter cells [adapted from reference (119)].

⁵ Clone is defined as an isolate or group of isolates descending from a common precursor exhibiting identical or closely similar genotypic traits, which are characterized by a specific strain-typing method (e.g. PFGE or MLST).

1.2.3.1. Clonal spread

The spread of MDR *Salmonella* clones has been often implicated in the dissemination of antibiotic resistance, including of clinically relevant genes encoding for ESBL, AmpC enzymes or PMQR genes, among humans and/or food-animals (120, 137, 144). Those MDR adaptive features might have contributed to their persistence in particular ecological niches and to establish an association to human disease, since several studies provide further evidences of transmission from poultry and pigs to humans of those clinically relevant MDR *Salmonella* clones (17, 120, 137, 144). Interestingly, the high-level resistance to ciprofloxacin (due to *gyrA/parC* mutations) and other antibiotics (amoxicillin, streptomycin, spectinomycin, gentamicin, sulfamethoxazole and tetracycline) (18, 62-65) of the previously mentioned *S. Kentucky* ST198-X1-SGI1 strain might also have contributed to its clonal expansion. Additionally, in this particular emerging clone production of carbapenemases (e.g. *bla*_{VIM-2}, *bla*_{OXA-48}) and ESBLs (e.g. *bla*_{CTX-M-25}), has been also reported occasionally (172, 174, 176). In EU, several *S. Stanley* outbreaks linked to turkey meat products consumption have been reported, with the epidemic clone presenting resistance to nalidixic acid and decreased susceptibility to ciprofloxacin (60). Additionally, in the recent outbreak in Austria three strains also presented resistance to gentamicin and cephalosporins (with production of *bla*_{CTX-M-15}) (60, 61). Several *S. Infantis* successful clones of broiler origin are also characterized by MDR profiles, including nalidixic acid, tetracyclines, sulphonamides or nitrofurans (50, 57, 113), antibiotics frequently used in livestock production, which could contribute to their co-selection in the poultry industry and consequently for its transmission to humans. In Belgium and France, particular MDR clones of *S. Infantis* carrying also *bla*_{TEM-52} gene have been found among poultry and human origins (162).

Concerning pig-associated serotypes, antimicrobial resistance acquisition to agents commonly used in livestock production (e.g. tetracycline's, sulphonamides) has also been associated with the spread of particular *Salmonella* serotypes/clones in both food-animals and humans. One example is the dissemination of the epidemic *S. Typhimurium* DT104 clone carrying the typical MDR ACSSuT (A-amoxicillin, C-chloramphenicol, S-streptomycin, Su-sulfamethoxazole and T-tetracycline) profile (17, 91, 107), in some cases associated with ciprofloxacin and/or extended-spectrum cephalosporin resistance (e.g. *bla*_{TEM-52}) (162). More recently in Europe the emergence and persistence of the most important *S. 4,[5],12:i:-* clones have been also related to diverse MDR profiles, particularly the European clone with ASSuT and the Spanish clone with ACGSSuTTm (G-gentamycin, Tm-trimethoprim) (89, 229). In addition, the description of ESBL genes (*bla*_{CTX-M-1}) in *S. 4,[5],12:i:-* isolates from pig sources belonging to the successful European clone is of

concern, due to the possibility of dissemination of these clinically relevant genes (156). Recently, the high incidence of PMQR determinants OqxAB \pm Aac(6')-Ib-cr that has been observed in Asia, appears to be associated with the clonal spread of a MDR *S. Typhimurium*, initially reported in food-animals (218) and later in human isolates (230, 231).

1.2.3.2. Horizontal gene transfer

Several studies have demonstrated that antimicrobial resistance determinants can be mobilized inside or out of the bacterial cells, most likely by intracellular (mobile genetic elements – MGE – such as transposons, gene cassettes and insertion sequences that can move from one genetic location to another in the same cell) or intercellular mobility (plasmids, transposons, genomic islands that can move from one bacteria to another) (148, 232, 233). Intercellular movement of DNA can occur via conjugation (transfers of genetic material from donor to recipient's cells through bacterial pilus), transformation (uptake of free DNA) or transduction (via bacteriophages) (Figure 11) (232).




FIGURE 11 – Horizontal gene transfer of antimicrobial resistance: (1) – transduction, (2) – conjugation and (3) - transposition [adapted with permission from reference (232)].

In *Salmonella* the major conjugative or mobilizable genetic elements that have contributed to its evolution, particularly with regard to the acquisition and dissemination of antimicrobial resistance, were plasmids, genomic islands, transposons and integrons/gene cassettes (234). Plasmids⁶ are the elements considered as the main drivers responsible for the dissemination of antibiotic resistance, including PMQRs, ESBLs and AmpC enzymes, in Gram-negative bacteria, inclusive in *Salmonella* (234-236). These genetic units are typically composed of conserved backbone modules with genes encoding for replication, maintenance and transfer functions as well as variable accessory modules (e.g. ABR, biocides tolerance), which mediate bacterial adaptation (148, 234). They are classified according to different criteria, such as the number of copies, host range (broad or narrow), their ability to transfer (conjugative or mobilizable) between cells and the incompatibility group (Inc) (i.e. two plasmids belonging to the same Inc can not coexist in the same cell) (148, 232). Incompatibility groups have been defined for plasmids of *Enterobacteriaceae*, using a PCR-based replicon-typing scheme, targeting sequences encoding replication (237). In *Salmonella*, transmission of clinically relevant genes have been associated with diverse Inc plasmids, such as the narrow host IncI1 (e.g. *bla*_{CTX-M-1}, *bla*_{TEM-52} and *bla*_{CMY-2}) and IncHI2 (e.g. *bla*_{CTX-M-2}, *bla*_{CTX-M-9}, *oqxAB* and/or *aac(6')-Ib-cr*) or the broad host IncA/C (e.g. *bla*_{CMY-2}) and IncN (e.g. *bla*_{CTX-M-15}), in both human and food-animals suggesting a direct link between the two bacterial populations through food-chain (Table 3 and 4) (161, 162, 168, 193, 204). One recent example was the presence of the same clone of *S. Infantis* carrying a *bla*_{CTX-M-1}-conjugative megaplasmid with other antibiotic resistance genes (~ 280–320Kb), isolated from Italian broiler chickens, broiler meat and humans, suggesting a clonal *Salmonella* transmission from the broiler chicken industry along the food-chain (204). Also the same clone of *S. Enteritidis* harbouring a FII plasmid with CTX-M-15 and several ABR genes was found in chicken, chicken meat and human stool (193). *S. Heidelberg* clones carrying I1 plasmids with the AmpC-*bla*_{CMY-2} enzyme were also found in both human and food-animal samples (e.g. chicken, turkey) (168).

Besides plasmid encoding antimicrobial resistance genes, a number of genomic islands⁷ (GIs) (238, 239), including *Salmonella* GI 1 variants (SGI1) have been associated with the spread of antibiotic resistance mainly to ampicillin, chloramphenicol, streptomycin, sulphonamides and tetracycline (240). One of the best well-known examples is the worldwide dissemination of the epidemic strain *S. Typhimurium* DT104

⁶ Plasmids are usually circular, self-replicating DNA molecules that normally exist in cells as extra-chromosomal replicons.

⁷ Genomic islands (GIs) are large chromosomal regions, usually carrying mobility genes coding for integrases or transposases that are required for chromosomal integration and excision, but also other additional genes encoding for adaptive features. These elements are part of the flexible gene pool, being frequently associated with tRNA genes and flanked by direct repeat structures and insertion sequences (238, 239).

from both humans and from food-producing animals (240). These isolates carried a mobilizable SGI1 (241, 242), consisting of a backbone with a class 1 integrons (In104 of the In4 type) that includes a pentaresistant phenotype [*aadA2*-*bla*_{PSE-1}-*sul1*-*tet*(G)-*floR-cmlA1*] (242). In fact, integrons⁸ (243), mainly class 1 integrons in *Salmonella*, seems to be an important source of acquisition of antibiotic resistance genes (91). The presence of a 2000bp integron carrying *bla*_{OXA-30} and *aadA1* gene cassettes in a MDR *S. Typhimurium* clones was also identified in both animal and human isolates (91, 244).

It is known that successful MDR *Salmonella* serotypes/clones persistence and spread among food-animals and the environment is derived, in part, to the acquisition of resistance mechanisms to commonly used antibiotics in animals and humans. However, several entities, pointed out tolerance to other non-antibiotic compounds with antimicrobial activity (e.g. metals, biocides), widely used in food-animals as a potential adaptive feature to different environments and hosts (animal/human) (245). More worrying is that persistence of these metals/biocides in different environments can contribute for the selection of metal/biocide tolerant and/or antibiotic-resistant bacteria due to mechanisms of cross and co-resistance (245, 246) (more details in the next topics).

1.3. Bacterial tolerance to biocides, particularly to metals

As discussed in the previous topics, antibiotics are considered one of the biggest selective pressures in diverse ecological niches for the emergence of antibiotic-resistant bacteria (118). With the ban of antibiotics as growth promoter's in the EU (247), the use of other non-antibiotic compounds with antimicrobial activity (e.g. metals/biocides) are currently applied in the food-animal setting to control and reduce foodborne zoonotic pathogenic bacteria (245, 248). Recently, the European Commission highlighted that these broad use of metals/biocides (even in subinhibitory concentrations) might contribute to the selection and maintenance of antibiotic-resistant bacteria in some ecological niches (245, 248). In fact, the activity of antibiotics and metals/biocides is often diminished by the same bacterial mechanism (cross-resistance) and/or by different mechanisms codified by genes co-located in common mobile genetic platforms (co-resistance), such as plasmids, genomic islands and transposons (245, 246, 249). These important issues will be addressed in the next topics.

⁸ Integrons are assembly platforms that incorporate exogenous open reading frames by site-specific recombination and convert them to functional genes by ensuring their correct expression. They are composed of three key elements: a gene encoding an integrase (*intI*), a primary recombination site (*attI*), and a promoter (P_c) that directs the transcription of the captured genes. At present, five classes of integrons (defined based on the sequence of the encoded integrases) are known, all physically linked to mobile DNA elements, being class 1 integrons the most widespread in *Enterobacteriaceae* (243).

1.3.1. Definition, classification, application and accumulation of metals and biocides

Biocides are chemical compounds capable of killing or inhibiting microorganisms (250). According to the Directive no. 98/8/EC (251), biocidal products are defined as “active substances and preparations containing one or more active substances, put up in the form in which they are supplied to the user, intended to destroy, deter, render harmless, prevent the action of, or otherwise exert a controlling effect on any harmful organism by chemical or biological means”. They are classified into at least 22 types of biocidal products distributed by 4 main groups (i.e. disinfectants, preservatives, pest control and others) (252). Additional classifications for biocides may be used based either on the functional group it possesses (e.g. alcohols, aldehydes, metals, halogens, quaternary ammonium compounds - QACs) or the bacterial target on which they act (e.g. proteins, membrane, nucleic acids or cell wall) (Table 5) (250).

TABLE 5 – Classification of biocides based on the bacterial target [adapted from reference (250)].

Membrane	Proteins	Nucleic acids	Cell wall
<ul style="list-style-type: none"> • QACs • Biguanides • Phenols • Phenylethers • Aldehydes • Acids • Terpenes • Alcohols • Anilides • Peroxygens • Parabens • Isothiazolones • Anionic surfactant 	<ul style="list-style-type: none"> • Alcohols • Phenols • Phenylethers • Aldehydes • Metals derivatives • Isothiazolones • Acids (parabens) • Peroxygens • Chlorine compounds • Biguanides • Vapor-phase disinfectant 	<ul style="list-style-type: none"> • Alcohols • Acids (parabens) • Antimicrobial dyes • Acridines • Biguanides • Aldehydes • Diamidines • Chlorine compounds • Metals derivatives • Peroxygens • Halogens • Vapor-phase disinfectant 	<ul style="list-style-type: none"> • Alcohols • Phenols • Aldehydes • Chlorine releasing compounds • Metals derivatives (mercurials)

Contrarily to antibiotics, which activity is frequently systemic, biocides are mainly applied as disinfectants, antiseptics or used as preservatives in a variety of settings (245, 253). Those included for example the hospital setting to control nosocomial infections (e.g. disinfectants on medical devices and surfaces, antiseptics used on skin and mucosa), in the food industry (e.g. preservation of food, surface disinfectants) and in the consumer's home within several products (e.g. cosmetics and personal care products, household products and textiles) (245, 253, 254). Additionally, in the food-animal setting biocides also played a key role for controlling microorganisms in animal farming management (biosecurity, hygiene), due to the mandatory need for limiting the presence of pathogens, such as *Salmonella*, in the food supply (29, 255-258). In fact, these

compounds (e.g. chlorhexidine, halogens, silver, copper, QACs, triclosan) have been largely applied for avoiding animal colonization/infection and contamination across the food chain (245, 259), from farms (including in animals) and hatcheries through slaughter facilities and dairy processing plants to food processing, packaging and retail facilities (Figure 12) (245, 253, 254, 259, 260). Additionally, metals can also be used in agriculture as pesticides and in soil fertilizers (e.g. arsenic) (246, 261) and can be applied as antimicrobial coatings and in animal wound dressings (e.g. silver, copper and zinc) (260, 262, 263). Also important is the extensive use of copper and zinc in feed or water (feed additive) as micronutrients and/or for growth promotion of certain farm animals and for enteric disease control (246, 264-267). In Europe, zinc has been usually incorporated in feed at concentrations not exceeding 250 mg/kg, for all animal species (264, 268). Zinc in USA is used between 50-100 ppm for dietary requirements, although higher concentrations can be used for growth promotion (2000-3000 ppm) (269). In the same way, copper has been used in feed at different concentrations according to animal type, ranging from a maximum of 175 mg/kg (~2.8 mM) in piglets to a minimum of 35 mg/kg (~0.6 mM) in species or animal categories other than piglets and ovines (264, 270). In USA copper supplements are not restricted and the typical amounts range from 125 ppm to 250 ppm (~2 to 4 mM) in all animal's diets (271). Other metals, such as arsenic, are still used as growth promoter in some countries (e.g. China) in the form of roxarsone, but in others (e.g. USA) occurred a voluntary withdrawn (261).




FIGURE 12 – Main sites of application of biocides in the food-animal setting as part of biosecurity measures [adapted with permission from reference (272)].

As a consequence of the extensive application of metals, including in feed for food-animals, they can be environmental pollutants and accumulate for example in manure or

waste lagoons in the animal production setting (246, 266, 273), and then be spread by sewage sludge and liquid manure applied in amended soils worldwide or by water bodies (246, 274-276). In fact, high concentrations (e.g. up to 3387.6 mg/kg, mg/kg; 53 mM) of copper have been observed in pig manure and consequently pig manure/sludge-amended soils (275-279). Other metals such as mercury and arsenic, considered as food and/or feed contaminants (273, 280) and tellurium, a contaminant mainly due to industrial pollution (281), could also contribute for these global burden of environmental metal contamination.

In contrast to the monitoring use of antibiotics in human and animal health (e.g. DANMAP⁹-Denmark; EMA/ESAC¹⁰-Europa; FDA¹¹-USA), there are few free publicly available compilations of data on sales or consumption of various metals and biocides (245). According to the last report of the European Commission, in 2009 the estimated annual total volume of sales of biocidal active ingredients (production and imports) in the EU was about 400.000 tonnes, most of which belonged to the class of disinfectants, with a greater incidence on the use of silver salts/nanoparticles (282, 283). In fact, production volumes of many of these compounds are considered to be several orders of magnitude higher than those of antibiotics (245).

Given its importance in the food-animal setting, in the next topics a focus will be given to the metals copper and silver widely applied in the animal welfare and, additionally, to three environmental, food and/or feed contaminants, tellurium, mercury and arsenic.

1.3.2. Mechanisms of action of the biocide activity of metals

Based on studies of the mechanisms of action of antimicrobial agents it is well known that antibiotics act in specific bacterial cell targets, resulting in the inhibition of essential physiological processes (e.g. nucleic acid synthesis, folate biosynthesis) or alteration of cellular structures (146, 147). Contrarily, biocides at in use-concentrations (i.e. concentrations, usually high, of the biocidal product applied in a real context) may have multiple and unspecific target sites within the cell (e.g. interaction with outer cellular components, cytoplasmic membrane, proteins, DNA, RNA and/or other cytosolic components), acting through physicochemical interactions or chemical reactions (Figure 13) (250, 260, 284, 285). For example, the main antimicrobial mechanism of action of

⁹ DANMAP is the Danish Programme for surveillance of antimicrobial consumption and resistance in bacteria from animals, food and humans.

¹⁰ EMA is the European Medicines Agency; ESAC is the European Surveillance of Antimicrobial Consumption Network.

¹¹ FDA is the United States Food and Drug Administration.

QACs (e.g. benzalkonium chloride, cetrимide) is primarily related to their cationic surfactant (detergent) properties (286). In fact, the association of QACs at in-use concentrations, with the phospholipids on the bacterial cytoplasmic membrane, leads to physical disruption and partial solubilisation of membrane, allowing vital intracellular components to be released (250, 253, 286, 287).




FIGURE 13 – Mechanisms of action and bacterial targets of biocides [reprinted from reference (259)].

Until now, several mechanisms of toxicity in bacteria have been described for metals (Figure 14), such as formation of reactive oxygen species and antioxidant depletion [e.g. As(III), Cu(II), Te(IV)], protein dysfunction [e.g. Ag(I), Cu(I), Hg(II), Te(IV)], impaired membrane function [e.g. Ag(I), Cu(II)] (e.g. compromising the cytoplasmic membrane integrity, disruption the activity of the bacterial electron transport chain) or genotoxicity [e.g. As(III), some organomercury complexes] (262, 263, 288, 289). Concerning some of non-essential metals for the cell, such as silver (Ag), mercury (Hg) or tellurium (Te), it is known that they are extremely toxic to most bacteria, having microbicide activity even at very low concentrations (288).




FIGURE 14 – Antimicrobial mechanisms of metal toxicity; Solid arrows represent pathways in which the underlying biochemistry has been elucidated, whereas dashed arrows represent a route of toxicity for which the underlying biochemical mechanism is unclear; ROS - reactive oxygen species [adapted with permission from reference (288)].

Contrarily, metals such as copper (Cu), are essential for eukaryotic and prokaryotic cellular functions (290). In fact, the essentiality of Cu, lies in its capacity to undergo redox cycling between the oxidized cupric form Cu^{2+} (borderline Lewis acid) and, the more toxic reduced cuprous form Cu^+ (soft Lewis acid) (291, 292). Because of this property, the Cu^+/Cu^{2+} couple with a high redox potential can act as electron donor/acceptor of many cupro-proteins (291, 292). Many of these enzymes are essential for diverse processes, such as oxidative phosphorylation (e.g. enzyme cytochrome c oxidase), iron homeostasis (e.g. ceruloplasmin) or superoxide dismutation (e.g. superoxide dismutase) (290, 292, 293). Nevertheless, at toxic concentrations, Cu can damage bacterial cells by multifactorial mechanisms of action (Figure 14). In fact, an important role for copper in phagosomal killing of bacteria has been reported, especially for *Salmonella*, a pathogen with an intracellular lifecycle, since during copper deficiency this ability seems to be diminished (Figure 15) (294, 295). Under aerobic condition, Cu can take part in Fenton-like reactions leading to oxidative damage by the generation of reactive oxygen species (ROS) (e.g. hydroxyl radicals, hydrogen peroxide and superoxide), which in turn damage lipids, proteins and DNA (263, 295). In the absence of oxygen, disruption of iron sulphur clusters in dehydratases involved in processes such as

branched-chain amino acid synthesis seems to be the most important mechanism (291, 295, 296).




FIGURE 15 – Copper mechanisms of action within macrophages of mammalian host, following phagocytosis of bacteria. CTR1 is a Cu^+ transport protein; ATOX1 is an antioxidant 1 copper metallochaperone and ATP7A is a copper-transporting P-type ATPase [reprinted with permission from reference (295)].

Apart from the several mechanisms of metal/biocide toxicity previously described, there are a number of intrinsic¹² (e.g. concentration, contact time and stability of the biocide) and extrinsic¹³ (e.g. temperature, pH, presence of organic matter mainly at the food-animal setting, presence of biofilms or microorganisms that are able to inactivate the biocide molecule) factors that could have a great influence in the efficacy of a biocide (297). Consequently, depending on the biocidal concentrations reached in a particular environment, the growth, metabolism/physiology and division cycle of the bacterial cell may be affected at different extents, which might contribute to the selection of tolerant bacteria (298). Tolerance mechanisms of metals included in the experimental work of this thesis will be discussed in the next topics.

1.3.3. Mechanisms of tolerance to metals, particularly in *Salmonella*

In the field of metals/biocides, it is known that terms such as “resistance” and “tolerance” have acquired recently specific technical meanings, although they have been

¹² The impact of intrinsic factors on the biocide activity depends on the biocide compound specific chemical features and the way it is applied by the user.

¹³ The impact of extrinsic factors on the biocide activity depends on the environmental conditions that are present during application of biocide.

applied rather loosely in several studies. The term bacterial resistance should be used when the phenomenon being studied is bacterial cell killing (297). According to Maillard and colleague's (299) biocide resistance is considered when a microbicide is ineffective against a microorganism that was previously susceptible to that microbicide. Contrarily, the term bacterial "tolerance" should be used while discussing adaptation to inhibitory concentrations of the biocide compared to wild type populations (297). The tolerance of a microorganism to a biocide comparing to the wild type populations might occur to concentrations well below those used in products applied in hospitals, domestic and industrial practice. Thus, these more tolerant microorganisms apparently do not compromise the biocide effectiveness in daily applications (297, 299). As in this thesis studies we evaluated the phenotypic changes, usually through the determination of MIC values compared to wild type bacterial populations henceforth we will use the term "tolerance".

Unlike antibiotics, a major problem when assessing tolerance to this type of compounds is the lack of available standard methodologies, which in turn leads to poor levels of comparability between studies and a lack of reliable data (245, 299). Indeed, in the case of antibiotics, standardized test methods allowed to define clinical breakpoints and epidemiological cut-off values (ECOFFs) based in MICs (134), which in the case of metals/biocides are currently scarce (300, 301). Development of standardized techniques is currently important for application in metal/biocide tolerance surveillance programmes, for providing informative data in biocidal product development or usage, and for helping to establish regulatory policies.

To persist and survive in metal contaminated environments, bacteria have evolved mechanisms to handle with toxic concentrations of these compounds (263). Metal tolerance in bacteria is not usually limited to a single mechanism but rather a combination of them, since these compounds have multiple targets inside the cell (288). There are several mechanisms of tolerance that bacteria use to withstand metal toxicity, such as, reduced uptake (e.g. regulation of transporters involved in metal uptake), efflux (e.g. membrane transporters, driven by ATP hydrolysis or chemiosmotic potential, able to transport specific toxic metals out of the cell), extracellular sequestration (e.g. up regulation of the expression of extracellular polymers or siderophores), intracellular sequestration (e.g. metal precipitation or use of cytoplasmic proteins, such as bacterioferritin and metallothioneins), repair, metabolic by-pass (e.g. production of alternative proteins with modified catalytic cores) or chemical modification (e.g. reactions that alter the chemical reactivity of the metal atoms) (Figure 16) (263, 288). Additionally,

physiological adapted bacterial communities such as in biofilms, were shown to withstand metal toxicity (302) as also previously demonstrated for antibiotics.

Some of these metal tolerance genes/mechanisms, also present in *Salmonella*, are centralized in an online platform (<http://bacmet.biomedicine.gu.se/>) (303) and will be detailed in the next topics.




FIGURE 16 – Intrinsic or acquired metal tolerance mechanisms [adapted with permission from reference (288)].

1.3.3.1. Copper homeostasis and tolerance mechanisms

Copper is an essential metal required in many biological processes (292). However, toxic Cu concentrations for microbes can occur during cellular immunological or predation strategies (e.g. macrophages, amoebas) and anthropogenic activities (e.g. animal feed additives, agriculture microbicides) in diverse environments (e.g. manure, soil) (246, 292, 304). To cope with both the essentiality and toxicity of Cu, *Enterobacteriaceae*, including *Salmonella*, have evolved by elaborating interwoven mechanisms of tolerance, homeostasis and delivery pathways, which act to tightly control intracellular Cu pools (305). In general, these mechanisms include Cu efflux (e.g. chromosomally or plasmid encoded efflux systems), Cu sequestration (e.g. by chaperones or siderophores) and copper oxidation (e.g. mixed copper oxidases and superoxide dismutase mimics) (293).

In *Salmonella*, intracellular copper homeostasis both in aerobic and anaerobic conditions is maintained mainly as the result of the activity of Cue (Cu efflux) system (306, 307). This system consists of the copper sensing MerR-family transcriptional regulator CueR that mediates copper-responsive expression of the cytoplasmic PIB1-type ATPase Cu⁺ efflux protein CopA, the periplasmic multi-copper oxidase CuiD (re-oxidation of Cu⁺ to Cu²⁺; also involved in *Salmonella* virulence) (308) and the periplasmic copper binding protein CueP (chaperone; particularly relevant in anaerobic conditions) (Figure

17) (294, 305-307, 309, 310). Besides the Cue system, the cytoplasmic copper-handling regulatory circuit is partially duplicated in *Salmonella* (305). In fact, an additional Gol system has been identified, consisting of a second CueR-like sensor GolS (primarily associated with gold sensing) and a cytoplasmic Atx1/CopZ copper chaperone-like protein GolB (311, 312). Also, a PIB1-type ATPase GolT, that together with CopA are involved in efflux as part of a Cu detoxification and a supply route to the periplasmic superoxide dismutase SodCII by the CueP chaperone, is also present (Figure 17) (305). Since *Salmonella* does not have the CusCBA HME-RND¹⁴ type efflux system, such as *E. coli* to transport Cu from the periplasm to outside the cell, a TolC protein was recently proposed as a potential outer membrane copper exporter in aerobic conditions (313).




FIGURE 17 – Copper homeostasis mechanisms based in Cue and Gol systems in *Salmonella* [adapted from reference (314)].

In environments where copper concentrations would overcome chromosomally encoded copper homeostatic systems, bacteria would be able to tolerate these higher copper values if they contain additional plasmid-encoded copper tolerance mechanisms (293). These extrachromosomal tolerance systems are usually metal oxidation state selective (293). The best known Cu tolerance mechanism although not yet fully characterized is the *pcoABCDRE* cluster (Figure 18). This cluster was firstly described in a *E. coli*

¹⁴ HME-RND is a Heavy Metal Efflux Resistance-Nodulation-Division system.

pRJ1004 plasmid (with MIC_{CuSO₄} up to 12-20 mM) recovered from a piggery effluent (315), and later in other *Enterobacteriaceae* such as *Salmonella* (with MIC_{CuSO₄} up to 18 mM) (316).




FIGURE 18 – Proposed copper (*pco* cluster) tolerance mechanism. The bottom line indicates the genes and their transcriptional and translational directions, with the open circles representing potential promoter regions/transcript start sites [adapted with permission from reference (304)].

The transcription of PcoABCD proteins appears to be regulated by PcoRS and *cusRS* seems to regulate PcoE protein (263, 304, 317). PcoA is a multi-copper oxidase and may have a similar function to CueO, oxidizing Cu^+ to less toxic Cu^{2+} . PcoB possibly functions as the outer membrane transporter (263, 304, 317). Sitting in the inner membrane, PcoD drives the transport of Cu^+ from the periplasm to the cytoplasm, with periplasmic PcoC chaperoning/delivering the Cu^+ to PcoD (263, 304, 317). PcoE is an additional chaperone that binds $\text{Cu}^+/\text{Cu}^{2+}$ but also Ag^+ and in the periplasm and probably shuttles it to PcoA and/or PcoS (317). The roles of PcoFG have not been elucidated. This *pco* cluster (only *pcoA* searched) has already been detected in a collection of specific *Salmonella* isolates from pig and related environments (318). However, the MIC_{CuSO₄} (16-24 mM) values detected were higher in aerobic conditions both in isolates carrying *pco* or not (318), which indicate that these genes might not lead to an increase in copper tolerance in such environmental conditions. Although similar MIC_{CuSO₄} values (20-28 mM) were previously found in *Salmonella* isolates from pig farms, in this case no

genotyping studies were performed to detect the presence of *pco* cluster (319). More recently, Hao and colleagues (304) have discussed the possibility of *pco* gene cluster be part of a genetic cluster comprising other metal tolerance determinants, such as the *sil* gene cluster associated with Ag tolerance. These authors proposed that those gene clusters might be a possible “copper pathogenicity island”, associated or not with Tn7-like elements (263, 304, 320-322).

1.3.3.2. Silver tolerance mechanisms

Silver is a non-essential metal, being extremely toxic to most bacteria, thus some tolerance mechanisms to overcome high concentrations of Ag have been described. The first described, and to date best-characterized Ag⁺ tolerance mechanism, is the *silCFBAGPRSE* gene cluster. It was firstly described in a 180kb IncHI2 plasmid, pMG101, of a *S. Typhimurium* from a burn unit (323). The region that is responsible for Ag tolerance ($\text{MIC}_{\text{AgNO}_3} > 256 \text{ mg/L}$; $> 1.5 \text{ mM}$), the inducible *silCFBAGPRSE*, comprises 9 genes (Figure 19) (304). In fact, 8 of these genes were primarily characterized based on homologies with other known metal tolerance determinants, such as the *cus* system of *E. coli* (324).




FIGURE 19 – Proposed silver (*sil* cluster) tolerance mechanism. The bottom line indicates the genes and their transcriptional and translational directions, with the open circles representing potential promoter regions/transcript start sites [adapted with permission from reference (304)].

SilE is predicted to bind Ag^+ and Cu^{2+} in aerobic conditions (325), transporting them to the HME-RND efflux system (SilCBA), responsible for exporting the metal ions out of the cell. Additionally, it was also proposed that this *silE* gene may have an additional role in Cu tolerance under more reducing environments (e.g. anaerobic conditions found in animal gut/manure/waste lagoons) by binding to the more toxic form Cu^+ (260, 304, 325). SilF is predicted to act as a metal chaperone to SilCBA too. The other putative efflux pump present in *sil* cluster is SilP, a P-type cation ATPase, which transports Ag^+ ions from the cytoplasm to the periplasm (262, 323, 326). While the expression of *silCFBAP* is thought to be controlled by the two component membrane sensor and transcriptional responder SilRS, the expression of *silE* is thought to be regulated/co-regulated by the Cus system (304, 317).

The *sil* genes have been reported as scattered in different bacterial genera, although most studies have been focused in non-*Salmonella* *Enterobacteriaceae* isolates (e.g. *Enterobacter cloacae* and *E. coli*), particularly recovered from nosocomial and animal/human clinical settings (325, 327-330) or environmental bacteria (326). The *sil* gene cluster, alone or associated with *pco* or other metal tolerance genes, has also been identified in *E. coli* pAPEC-02-R (331), *Serratia marcescens* pR478 (332), *Klebsiella pneumonia* pK2044 and pLVPK plasmids (333, 334). In *Salmonella*, *sil* associated with *pco* cluster has been found in sequenced chromosomes and plasmids of some serotypes (304). Although silver tolerance genes have been already reported, an association between the presence of the *sil* cluster and the corresponding phenotypic behaviour, obtained by different susceptibility methods, has only been established in few *Enterobacteriaceae* isolates (327-330, 335), remaining unexplored in comprehensive *Salmonella* collections.

1.3.3.3. Mercuric ion tolerance mechanisms

Mercury (Hg) is toxic to humans and with no known positive role in cellular function. However, it has been widely used in chemical, agriculture and pharmaceutical industry (e.g. thimerosal, merbromin, nitromersol) since the XIX century (336). Nowadays, due to its toxicity, the usage of antimicrobial mercury compounds is in decline and is likely to be replaced (263). Elemental Hg^0 (non-toxic) present in the air is able to settle into the water or land, oxidized to Hg^{2+} and transformed into the highly toxic form of methyl mercury (336). This cycle regularly exposes microorganisms, humans, animals and plants to different Hg forms (inorganic and organic), by different routes (e.g. water, soil, food chain) (337).

Bacterial tolerance to Hg, including in *Salmonella*, is mainly due to the presence of the *mer* operon, that comprise a group of proteins involved in the detection, scavenging,

transport and reduction of Hg (336, 338). Transcription of *mer* operon is induced (or repressed) by the product of *merR* gene in the presence (or absence) of Hg^{2+} ion, and its expression is further regulated by *merD* (339). Two types of *mer* operons were mainly described including those conferring tolerance only to inorganic Hg salts (narrow type) or both to organomercurials and inorganic Hg salts (broad type) (336, 338, 339). All types of *mer* operons contain a gene encoding a mercuric reductase (MerA) responsible for the enzymatic reduction in the cytosol of Hg^{2+} to the volatile Hg^0 (Figure 20) (336, 339). The periplasmic Hg^{2+} scavenging protein (MerP) and one or more inner membrane spanning proteins (MerT, MerC, MerE, MerF, MerG) that transport Hg^{2+} to the cytoplasm are also reported as part of the *mer* operon (Figure 20) (336, 339). Broad types of *mer* operons also contain a organomercurial lyase MerB that hydrolyses organic radicals before the action of *merA* (339).

Most of *mer* genes are described to be widespread among diverse bacterial hosts, genetic platforms and environments (339, 340). Among *Salmonella*, *mer* operons or *merA* gene were recently found downstream of the chromosomal RR1–RR2 antibiotic resistance region in *S. 4,[5],12:i:-* (229), similar to those reported in several *Salmonella* genomic island 1 (SGI1) variants (240, 341, 342). Additionally, this operon was found in transposons, particularly of the Tn21-like family (340), widespread in different *Enterobacteriaceae*, including in the recent *E. coli* O104:H4 mass food poisoning outbreak isolates from 2011 (343, 344) and in *Salmonella* isolates from diverse serotypes (321, 341, 342). *Salmonella* conjugative plasmids of IncA/C, IncF and IncHI2 families from human and food-animal samples (e.g. chicken, pig, cattle) carrying *mer* genes have also been reported (182, 321, 345-353). In fact, a recent study demonstrated that a *S. Infantis* clone associated with several outbreaks, carrying a plasmid harbouring the *mer* operon was able to grow in the presence of 25 mM mercury compared to the pre-emergent strain (without the *mer* operon), highlighting the role of *mer* operon for Hg tolerance in contaminated environments (113).




FIGURE 20 – Proposed mercury (*mer* operon) tolerance mechanism; a) the generic *mer* operon with curve brackets representing variable genes, b) The cellular *mer*-encoded mercury detoxification mechanisms, L=ligand with subscripts denoting the ligand type [adapted from reference (339)].

1.3.3.4. Tolerance mechanisms to metalloids arsenic and tellurite

During several years' organic arsenic (As) derived compounds, such as carbarsone, nitarosone and roxarsone have been used as feed additives for poultry, in the USA and China, acting as growth promoters and in controlling coccilobacillosis disease (263, 354). A recent study suggested that organoarsenic feed additives in the animal wastes are an important source of arsenate and arsenite pollution in Chinese fertilized agricultural lands, where organoarsenic compounds are still widely used (355). Only very recently, in June 2011, the US Food and Drug Administration announced the voluntary withdraw of the sale of roxarsone due to the presence of highly toxic inorganic arsenic residues in chicken meat from chickens fed on roxarsone-supplemented feeds (354). Arsenics can be found in animals and plants combined with carbon and oxygen to form organic arsenic compounds or in the environment combined with oxygen, chlorine or sulphur to form highly toxic inorganic As compounds [e.g reduced form As(III) (arsenite) and oxidised As(V) (arsenate)] (263, 356).

In *Enterobacteriaceae*, including *Salmonella*, the most common As tolerance mechanism, usually conferring tolerance to As(III), As(V) and antimonial compounds, is associated with the presence of a 5-gene system *ars* operon usually plasmid located (Figure 21) (356). Although the minimum *ars* operon consist always of *arsR* [As(III)-

responsive *trans*-acting transcriptional repressor protein], *arsB* [As(OH)₃/H⁺ antiporter protein that could be an ATP-dependent process or require the help of ArsA] and *arsC* [arsenate reductase that reduces As(V) to As(III)], additional genes such as *arsA* [ATPase that provides energy to ArsB for the efflux of As(III), which is more efficient than ArsB alone] and *arsD* [encoding a negative regulatory protein that binds to the promoter region of the operon and represses the transcription but also a metallochaperone for As(III) efflux via ArsAB] could also be present (Figure 21) (263, 356, 357).




FIGURE 21 – Proposed arsenic (*ars* operon) tolerance mechanism. Arsenic enters the cell as either As(III) by the aquaglyceroporin, GlpF or As(V) by *pst* and *pho* phosphate permeases [reprinted from reference (357)].

In *Salmonella*, arsenic tolerance mechanisms are less described than in other *Enterobacteriaceae*, such as *Klebsiella pneumoniae* carrying IncFII_K plasmids with *ars* genes (e.g. GenBank accession nos. CP000648, JX430448, CP002474, KF719971) (358, 359). However, these genes have already been reported in *Salmonella* chromosome (360) as well as in few plasmids from IncH (161, 348) and IncI1 families (361). In addition, very few studies in *Enterobacteriaceae*, including *Salmonella*, have demonstrated an association between the *ars* operon and arsenic tolerance and, when available, the majority of susceptibility techniques performed are quite different between studies, hindering results comparison. Joerger and colleagues (360) have demonstrated a higher incidence (in all studied isolates) of *arsB* and *arsD* genes in *S. Kentucky*, a USA poultry-associated serotype, comparing with other *Salmonella* serotypes (26/58 positive strains). In this study, with few exceptions, the acquisition of *ars* genes was correlated with a certain level of arsenate tolerance [*ars*⁺= 0.5-10 mM *versus* *ars*⁻≤ 0.1 mM As(V)] and arsenite [*ars*⁺= 0.1-1 mM *versus* *ars*⁻≤ 0.05 mM As(III)], highlighting the role of *ars*

operon for survival in arsenic contaminated environments (e.g. animal cecum or litter sources) (360).

Tellurite [TeO_3^{2-} ion or Te(IV)], is the oxyanion of the chalcogen tellurium (Te), that can be reduced to the insoluble and less toxic Te^0 (362, 363). Tellurite is highly toxic to bacteria (concentrations as low as 1 mg/mL) and therefore, was used for many years as an antimicrobial and therapeutic agent in medicine (364). Currently several forms of tellurium are used in industry (e.g. cadmium telluride solar panels, steel manufacturing industry, pesticide in agriculture) and as a selective agent in diverse microbiological culture media for the isolation of pathogens (362, 364). Until now six bacterial tellurite tolerance mechanisms have been proposed (coded by *ter*, *teh*, *tel*, *tpm*, *cysK*, and *ars* genes), although the genetic, biochemical and/or physiological bases underlying their tolerance have not been defined (363). Apparently, tellurite-tolerant bacteria can be associated with mechanisms that cause oxyanion extrusion or biochemical modifications different from reduction (e.g. generation of methylated forms of Te) (362). One of the most important mechanisms in *Enterobacteriaceae*, and that usually confers higher levels of TeO_3^{2-} tolerance (512-1024 $\mu\text{g K}_2\text{TeO}_3/\text{ml}$), is that encoded by the *ter* operon (common *terZABCDE* \pm *terF* or *terW*) (363). Currently little is known about the function of *ter* operon genes, including the tolerance mechanism that remains unknown (363).

In *Salmonella* this *ter* operon has been frequently associated with *ars* operon, the two ubiquitously present in IncH plasmids (161, 182, 321, 348). This determinant was also found in IncH (e.g. GenBank accession nos. AP006726, BX664015, DQ517526, JX182975) and IncF (e.g. GenBank accession nos. CP001919) plasmids from other *Enterobacteriaceae*, being also associated with pathogenicity islands (e.g. tellurite genomic islands-TER-GIs) of *Escherichia coli* O157:H7 (365, 366), and with hyper virulent clonal groups of *Klebsiella pneumonia* (367).

There is no doubt that metals and biocides are invaluable compounds that provide society with a number of benefits, contributing for the general human and animal health. However, the widespread use of these compounds, including in the food-animal setting, led to the emergence of some concerns by the European Union Scientific Committee for New and Emerging Health Risks, including the emergence of metal and biocide tolerance and their potential role for the selection and persistence of antibiotic-resistant bacteria (discussed in the next topic).

1.3.4. Bacterial tolerance to metals and biocides and the possible link with the selection and persistence of antibiotic-resistant *Salmonella*

The serious concern that exposure of bacteria to metals/biocides may boost the spread of antibiotic resistance has contributed to the establishment of a new European Biocidal Product Regulation (252). This new document requires that biocidal products manufacturers, besides the evaluation of safety and efficacy of the substances, ensure that they will not contribute to antimicrobial resistance development (252). Additionally, extensive metals/biocides environmental contamination, even at low-level concentrations, is a topic of increasing concern recognized by diverse authorities and authors (246, 265, 266, 270, 368, 369). Since some of those compounds, including metals, may not be completely degraded, they can persist in the environment, contributing for the selection of metal tolerant and/or antibiotic-resistant bacteria by mechanisms of cross or co-resistance (246, 249, 267). Metals can also potentially participate as catalysts of gene transfer among diverse microorganisms sharing the same ecologic niche (246, 369). In fact, recent environmental studies in metal contaminated soils/animal manure, particularly with Cu due to anthropogenic-derived sources (e.g. animal feed additives, agriculture microbicides), have shown that metal availability (e.g. 22.4-3172 mg/Kg; ~0.4-50 mM of Cu), even at low level concentrations, was correlated with a high prevalence of antibiotic resistance in diverse bacteria (277-279, 370, 371). Although the previous studies have shown the prevalence of antibiotic-resistant bacteria in the presence of metals/biocides, the factors behind the selection and maintenance of these antimicrobial resistant bacteria were not fully elucidated.

In *Salmonella* tested in laboratory conditions, exposure of *S. Typhimurium* to different concentrations of biocides (e.g. QACs, triclosan, oxidizing compound, aldehyde, oil phenol) was able to select for mutants with decreased biocide susceptibility, with most of them also often displaying decreased susceptibility to various antibiotics, mainly by overexpression of AcrAB-TolC efflux pump, indicating that biocides can mediate co-selection of antibiotic resistance by cross-resistance mechanisms (372-375). As previously mentioned co-selection of antibiotic-resistant bacteria can be due to cross-resistance, which describes one mechanism that counteracts two or more antimicrobial agents, such as antibiotics and metals/biocides (249). Probably one of the most important mechanisms in *Enterobacteriaceae*, including *Salmonella*, is related to the wide spectrum of substrates

(e.g. QACs¹⁵, antibiotics-fluoroquinolones) recognized by different chromosomally or plasmid encoded efflux systems (Figure 22) (253). In fact, overexpression of *Salmonella* and *E. coli* AcrAB-TolC efflux pumps is hypothesized to mediate cross-resistance to fluoroquinolones, a clinically important antibiotic, and QACs (253, 376, 377). Additionally, the transferable plasmid-encoded RND multidrug efflux pump, OqxAB, recently dispersed in *S. Typhimurium* (201, 218), showed to confer reduced susceptibility to biocides-QACs (benzalkonium chloride and to some extent cetrимide) and to diverse antibiotics (fluoroquinolones, chloramphenicol, trimethoprim and olaquinox) (378), which is of special concern due to the current use of fluoroquinolones for treating invasive salmonellosis.




FIGURE 22 – Representation of the main types of multidrug-resistance efflux pumps involved in extrusion of QACs and several other antibiotics in Gram-negative bacteria [reprinted from reference (253)].

Besides cross-resistance, probably one of the most important co-selection mechanisms in *Enterobacteriaceae* is co-resistance (246, 249). This can result from the acquisition of two or more resistance/tolerance genes linked together in the same genetic unit (such as a plasmid, transposon, genomic island or integron) (246, 249). This physical linkage results in the co-selection of all the genes in the same element even in the presence of a single environmental selective pressure (e.g. antibiotics or metals/biocides) (246, 249). In fact, Pal and colleagues (379) have recently demonstrated the co-occurrence of

¹⁵ The primary mechanisms of QACs tolerance in Gram-negative bacteria, including *Salmonella*, are due to the acquisition of genes for specialized QAC efflux pumps or overexpression of those chromosomally encoded systems (e.g. AcrAB-TolC proteins from RND family) (287). Although, other mechanisms such as reduced expression of porins (e.g. *S. Typhimurium* exposed to sub lethal concentrations of three QACs, where found with reduced levels of the outer membrane porins OmpC, OmpF and OmpA) could also be present (374).

metals/biocides tolerance and antibiotic resistance genes in fully sequenced bacterial genomes and plasmids from several bacteria taxa. These authors have found metals/biocides tolerance genes in 86% of bacterial genomes, being 17% co-located with antibiotic resistance genes, although these datasets are not representative of the full microbial diversity (379). Independently of the chromosomal or plasmid location, bacteria with metals/biocides tolerance genes carried more frequently antibiotic resistance genes (379). Additionally, the metals/biocides tolerance genes commonly co-occurring with antibiotic resistance genes were mercury tolerance and the *qacEA1* genes (379). This data is not surprising since for example class 1 integrons, widely distributed in *Enterobacteriaceae*, including *Salmonella* from human and food-animal origins, are usually associated with *qacEA1* genes plus sulphonamide resistance (*sul1* ou *sul3*) and other antibiotic resistance gene cassettes (e.g. trimethoprim - *dfrA1* and *dfrA12* and aminoglycosides - *aadA2*) (91, 380-382). In fact, it has been described that class 1 integrons are more prevalent in bacteria exposed to detergents/biocides and that these compounds could co-select for antibiotic resistance due to the co-location of both genes in the same genetic elements (383). Similarly, the widespread of mercury resistance genes has been associated with Tn21 family, with most of these transposons also carrying class 1 integrons and antibiotic resistance genes (263, 340)). In *Salmonella*, *mer* operons have been found increasingly associated with multidrug resistance genes, either associated with chromosomal resistance region (ASSuT phenotypes) as in the case of *S. 4,[5],12:i:-* European clone (229) or with SGI1 variants (e.g. SGI1-K1 AGSSuT profile; SGI1-K2 ASSuT; SGI1-L1 ACGSSuTTr) in diverse *Salmonella* serotypes (240, 341, 342, 384). Although widely reported in *Salmonella*, a correlation between environmental mercury exposure and increased antibiotic resistance has never been established for this foodborne pathogen.

Pal et al (379) also found that clinically important genera mainly recovered from human and domestic animals, including *Salmonella*, tended to carry both metal/biocide tolerance and antibiotic resistance genes on the same plasmids. Notably, the frequency of antibiotic resistance genes was significantly higher on metal/biocide tolerance genes-carrying plasmids, than plasmids without metal/biocide tolerance genes (379). This is not uncommon since a number of plasmids from different families (e.g. IncH, IncA/C), also present in *Salmonella*, have been reported as carrying both metals (*sil/pco* clusters, *mer*, *ars*, *ter* operons) and multiple antibiotic resistance genes (304, 320, 321, 349). In fact, recently, an *in vitro* assay showed that the presence of sub lethal levels of Cu or As were sufficient to co-select for the maintenance of *E. coli* and *Klebsiella pneumoniae* MDR plasmids (385), highlighting the role of these compounds for selection and persistence of bacteria carrying these genetic elements. Also, Cavaco and colleagues (386, 387) argued

that the use of zinc in feed could be an additional factor that may have contributed to the emergence of Meticillin-Resistant *Staphylococcus aureus* (MRSA), although in this case the antibiotic resistance and metal tolerance genes did not have a plasmid location. These authors have found that several MRSA strains carried *czrC* gene, associated with reduced zinc and cadmium susceptibility, in the *SCCmec* genomic island characteristic of these strains either from human or food-animals (386-388).

Some of the previously reported evidences highlight that a variety of biocides and metals (e.g. widely used as disinfectants, antiseptics/medicines, preservatives, growth promoters) found in the food-animal setting and other diverse polluted environments could have the ability to co-select for antibiotic-resistant bacteria and/or resistance plasmids (246, 249). Nevertheless, a comprehensive analysis of metal tolerance, including acquired metal gene dispersion and their genetic context within different *Enterobacteriaceae*, including *Salmonella*, from diverse ecological niches or belonging to frequent or emergent MDR clones, remains poorly explored. Additionally, we urgently need to be aware of the simultaneously action of biocides/metals with other stressors (e.g. antibiotics; reducing environments) for an enhanced survival, selection and persistence of particular metal/biocide tolerant-MDR strains.

REFERENCES

1. Grimont PA, Weill FX, eds. Antigenic formulae of the *Salmonella* serovars, 9th edn. Paris, France: WHO Collaborating Centre for Reference and Research on *Salmonella*, Institut Pasteur, 2007.
2. Hur J, Jawale C, Lee JH. Antimicrobial resistance of *Salmonella* isolated from food animals: A review. Food Research International. 2012;45(2):819-30.
3. Parry CM, Threlfall EJ. Antimicrobial resistance in typhoidal and nontyphoidal salmonellae. Curr Opin Infect Dis. 2008;21(5):531-8.
4. Chen HM, Wang Y, Su LH, Chiu CH. Nontyphoid *Salmonella* infection: microbiology, clinical features, and antimicrobial therapy. Pediatr Neonatol. 2013;54(3):147-52.
5. Crump JA, Sjolund-Karlsson M, Gordon MA, Parry CM. Epidemiology, Clinical Presentation, Laboratory Diagnosis, Antimicrobial Resistance, and Antimicrobial Management of Invasive *Salmonella* Infections. Clin Microbiol Rev. 2015;28(4):901-37.
6. Barrow PA, Jones MA, Smith AL, Wigley P. The long view: *Salmonella*-the last forty years. Avian Pathol. 2012;41(5):413-20.
7. Cosby DE, Cox NA, Harrison MA, Wilson JL, Buhr RJ, Fedorka-Cray PJ. *Salmonella* and antimicrobial resistance in broilers: A review. The Journal of Applied Poultry Research. 2015;24(3):408-26.
8. Majowicz SE, Musto J, Scallan E, Angulo FJ, Kirk M, O'Brien SJ, *et al*. The global burden of nontyphoidal *Salmonella* gastroenteritis. Clin Infect Dis. 2010;50(6):882-9.
9. Scallan E, Hoekstra RM, Angulo FJ, Tauxe RV, Widdowson MA, Roy SL, *et al*. Foodborne illness acquired in the United States-major pathogens. Emerg Infect Dis. 2011;17(1):7-15.
10. European Food Safety Authority (EFSA); The european union summary report on trends and sources of zoonoses, zoonotic agents and food-borne outbreaks in 2013. EFSA J. 2015;13:165. Available at: <http://www.efsa.europa.eu/en/efsajournal/pub/3991> (last accessed 1 December 2015).
11. BCCDC BC Center for Disease Control; BC Integrated Surveillance of Foodborne Pathogens (BCISFP): *Salmonella* findings 2013. 2014. Available at: http://www.bccdc.ca/NR/rdonlyres/3872B775-8B56-459B-9EE9-4DB7774C8D5B/o/2013BC_IntegratedSalmAnnualReport.pdf- (last accessed 1 December 2015).
12. Centers for Disease Control and Prevention (CDC); National Enteric Disease Surveillance: *Salmonella* Annual Report, 2012. 2014. Available at:

<http://www.cdc.gov/ncezid/dfwed/pdfs/salmonella-annual-report-2012-508c.pdf> (last accessed 1 December 2015).

13. Hugas M, Beloeil P. Controlling *Salmonella* along the food chain in the European Union - progress over the last ten years. *Euro Surveill.* 2014;19(19).

14. European Food Safety Authority (EFSA) Panel on Biological Hazards (BIOHAZ); Scientific Opinion on a quantitative estimation of the public health impact of setting a new target for the reduction of *Salmonella* in broilers. *EFSA J.* 2011;9:2106. Available at: <http://www.efsa.europa.eu/en/efsajournal/pub/2106> (last accessed 1 December 2015).

15. European Food Safety Authority (EFSA) Panel on Biological Hazards (BIOHAZ); Scientific Opinion on an estimation of the public health impact of setting a new target for the reduction of *Salmonella* in turkeys. *EFSA J.* 2012;10:2616. Available at: <http://www.efsa.europa.eu/en/efsajournal/pub/2616> (last accessed 1 December 2015). 2012.

16. European Food Safety Authority (EFSA) Panel on Biological Hazards (BIOHAZ); Scientific Opinion on the evaluation of molecular typing methods for major food-borne microbiological hazards and their use for attribution modelling, outbreak investigation and scanning surveillance: Part 1 (evaluation of methods and applications). *EFSA J.* 2013;11:3502. Available at: <http://www.efsa.europa.eu/en/efsajournal/pub/3502> (last accessed 1 December 2015).

17. Butaye P, Michael GB, Schwarz S, Barrett TJ, Brisabois A, White DG. The clonal spread of multidrug-resistant non-typhi *Salmonella* serotypes. *Microbes Infect.* 2006;8(7):1891-7.

18. European Food Safety Authority (EFSA); EU Summary Report on antimicrobial resistance in zoonotic and indicator bacteria from humans, animals and food in 2013. *EFSA J.* 2015b;13:4036. Available at: <http://www.efsa.europa.eu/en/efsajournal/pub/4036> (last accessed 1 December 2015).

19. European Feed Manufacturers' Federation (FEFAC); Statistics. 2012. Available at: <http://www.fefac.eu/files/47239.pdf> (last accessed 1 December 2015).

20. Food and Agriculture Organization (FAO) of the United Nations; Meat consumption. 2014. Available at: <http://www.fao.org/ag/againfo/themes/en/meat/background.html> (last accessed 1 December 2015).

21. Achtman M, Wain J, Weill FX, Nair S, Zhou Z, Sangal V, *et al.* Multilocus sequence typing as a replacement for serotyping in *Salmonella enterica*. *PLoS Pathog.* 2012;8(6):e1002776.

22. Switt AI, Sulakvelidze A, Wiedmann M, Kropinski AM, Wishart DS, Poppe C, *et al.* *Salmonella* Phages and Prophages: Genomics, Taxonomy, and Applied Aspects. In:

Schatten H, Eisenstark A, editors. *Salmonella*: Methods and Protocols. 2nd ed. New York: Humana Press; 2015.

23. Gopinath S, Carden S, Monack D. Shedding light on *Salmonella* carriers. Trends Microbiol. 2012;20(7):320-7.

24. Gomes-Neves E, Antunes P, Tavares A, Themudo P, Cardoso MF, Gartner F, *et al.* *Salmonella* cross-contamination in swine abattoirs in Portugal: Carcasses, meat and meat handlers. Int J Food Microbiol. 2012;157(1):82-7.

25. Andres VM, Davies RH. Biosecurity Measures to Control *Salmonella* and Other Infectious Agents in Pig Farms: A Review. Comprehensive Reviews in Food Science and Food Safety. 2015;14(4):317-35.

26. Hernandez-Reyes C, Schikora A. *Salmonella*, a cross-kingdom pathogen infecting humans and plants. FEMS Microbiol Lett. 2013;343(1):1-7.

27. Foley SL, Nayak R, Hanning IB, Johnson TJ, Han J, Ricke SC. Population dynamics of *Salmonella enterica* serotypes in commercial egg and poultry production. Appl Environ Microbiol. 2011;77(13):4273-9.

28. Ricke SC. Application of Molecular Approaches for Understanding Foodborne *Salmonella* Establishment in Poultry Production. Advances in Biology. 2014;2014:25.

29. Commission Regulation (EC) No 2160/2003 of the European Parliament and of the Council of 17 November 2003 on the control of *Salmonella* and other specified food-borne zoonotic agents [2003]. OJ L325/1.

30. Commission Regulation (EC) No 1168/2006 of 31 July 2006 implementing Regulation (EC) No 2160/2003 as regards a Community target for the reduction of the prevalence of certain *Salmonella* serotypes in laying hens of *Gallus gallus* and amending Regulation (EC) No 1003/2005 [2006]. OJ L211/4.

31. Commission Regulation (EC) No 200/2010 of 10 March 2010 implementing Regulation (EC) No 2160/2003 of the European Parliament and of the Council as regards a Union target for the reduction of the prevalence of *Salmonella* serotypes in adult breeding flocks of *Gallus gallus* [2010]. OJ L61/1.

32. Commission Regulation (EC) No 517/2011 of 25 May 2011 implementing Regulation (EC) No 2160/2003 of the European Parliament and of the Council as regards a Union target for the reduction of the prevalence of certain *Salmonella* serotypes in laying hens of *Gallus gallus* and amending Regulation (EC) No 2160/2003 and Commission Regulation (EU) No 200/2010 [2010]. OJ L138/45.

33. Mazengia E, Samadpour M, Hill HW, Greeson K, Tenney K, Liao G, *et al.* Prevalence, concentrations, and antibiotic sensitivities of *Salmonella* serovars in poultry from retail establishments in Seattle, Washington. J Food Prot. 2014;77(6):885-93.

-
- 34.** United States Department of Agriculture/Food Safety and Inspection Service (USDA/FSIS). Progress report on *Salmonella* and *Campylobacter* testing of raw meat and poultry products, 1998–2013. 2014. Available at: <http://www.fsis.usda.gov/wps/wcm/connect/885647f4-2568-48bf-ae5c-4a0d8279f435/Progress-Report-Salmonella-Campylobacter-CY20113.pdf?MOD=AJPERES> (last accessed 1 December 2015).
- 35.** Antunes P, Réu C, Sousa JC, Peixe L, Pestana N. Incidence of *Salmonella* from poultry products and their susceptibility to antimicrobial agents. *Int J Food Microbiol.* 2003;82(2):97-103.
- 36.** Donado-Godoy P, Byrne BA, Leon M, Castellanos R, Vanegas C, Coral A, *et al.* Prevalence, resistance patterns, and risk factors for antimicrobial resistance in bacteria from retail chicken meat in Colombia. *J Food Prot.* 2015;78(4):751-9.
- 37.** Ribeiro A, Kellermann A, Bessa M, dos Santos L, do Nascimento V. *Salmonella* spp. in raw broiler parts: occurrence, antimicrobial resistance profile and phage typing of the *Salmonella* Enteritidis isolates. *Braz J Microbiol.* 2007;38(2):296-9.
- 38.** Kich JD, Coldebella A, Mores N, Nogueira MG, Cardoso M, Fratamico PM, *et al.* Prevalence, distribution, and molecular characterization of *Salmonella* recovered from swine finishing herds and a slaughter facility in Santa Catarina, Brazil. *Int J Food Microbiol.* 2011;151(3):307-13.
- 39.** Abd-Elghany SM, Sallam KI, Abd-Elkhalek A, Tamura T. Occurrence, genetic characterization and antimicrobial resistance of *Salmonella* isolated from chicken meat and giblets. *Epidemiol Infect.* 2015;143(5):997-1003.
- 40.** Adeyanju GT, Ishola O. *Salmonella* and *Escherichia coli* contamination of poultry meat from a processing plant and retail markets in Ibadan, Oyo State, Nigeria. *Springerplus.* 2014;3:139.
- 41.** Akoachere JF, Tanih NF, Ndip LM, Ndip RN. Phenotypic characterization of *Salmonella* Typhimurium isolates from food-animals and abattoir drains in Buea, Cameroon. *J Health Popul Nutr.* 2009;27(5):612-8.
- 42.** Kikui GM, Ombui JN, Mitema ES. Serotypes and antimicrobial resistance profiles of *Salmonella* isolates from pigs at slaughter in Kenya. *J Infect Dev Ctries.* 2010;4(4):243-8.
- 43.** Ta YT, Nguyen TT, To PB, Pham da X, Le HT, Thi GN, *et al.* Quantification, serovars, and antibiotic resistance of *Salmonella* isolated from retail raw chicken meat in Vietnam. *J Food Prot.* 2014;77(1):57-66.
- 44.** Yang B, Cui Y, Shi C, Wang J, Xia X, Xi M, *et al.* Counts, serotypes, and antimicrobial resistance of *Salmonella* isolates on retail raw poultry in the People's Republic of China. *J Food Prot.* 2014;77(6):894-902.

-
45. Yoon RH, Cha SY, Wei B, Roh JH, Seo HS, Oh JY, *et al.* Prevalence of *Salmonella* isolates and antimicrobial resistance in poultry meat from South Korea. *J Food Prot.* 2014;77(9):1579-82.
46. Thai TH, Hirai T, Lan NT, Yamaguchi R. Antibiotic resistance profiles of *Salmonella* serovars isolated from retail pork and chicken meat in North Vietnam. *Int J Food Microbiol.* 2012;156(2):147-51.
47. Yang B, Qu D, Zhang X, Shen J, Cui S, Shi Y, *et al.* Prevalence and characterization of *Salmonella* serovars in retail meats of marketplace in Shaanxi, China. *Int J Food Microbiol.* 2010;141(1-2):63-72.
48. Clemente L, Manageiro V, Ferreira E, Jones-Dias D, Correia I, Themudo P, *et al.* Occurrence of extended-spectrum β -lactamases among isolates of *Salmonella enterica* subsp. *enterica* from food-producing animals and food products, in Portugal. *Int J Food Microbiol.* 2013;167(2):221-8.
49. Figueiredo R, Henriques A, Sereno R, Mendonca N, da Silva GJ. Antimicrobial resistance and extended-spectrum β -lactamases of *Salmonella enterica* serotypes isolated from livestock and processed food in Portugal: an update. *Foodborne Pathog Dis.* 2015;12(2):110-7.
50. Gal-Mor O, Valinsky L, Weinberger M, Guy S, Jaffe J, Schorr YI, *et al.* Multidrug-resistant *Salmonella enterica* serovar Infantis, Israel. *Emerg Infect Dis.* 2010;16(11):1754-7.
51. Dhanani AS, Block G, Dewar K, Forgetta V, Topp E, Beiko RG, *et al.* Genomic Comparison of Non-Typhoidal *Salmonella enterica* Serovars Typhimurium, Enteritidis, Heidelberg, Hadar and Kentucky Isolates from Broiler Chickens. *PLoS One.* 2015;10(9):e0137697.
52. Dutil L, Irwin R, Finley R, Ng LK, Avery B, Boerlin P, *et al.* Ceftiofur resistance in *Salmonella enterica* serovar Heidelberg from chicken meat and humans, Canada. *Emerg Infect Dis.* 2010;16(1):48-54.
53. Hoffmann M, Zhao S, Pettengill J, Luo Y, Monday SR, Abbott J, *et al.* Comparative genomic analysis and virulence differences in closely related *Salmonella enterica* serotype Heidelberg isolates from humans, retail meats, and animals. *Genome Biol Evol.* 2014;6(5):1046-68.
54. European Food Safety Authority (EFSA); The European Union Summary Report on Trends and Sources of Zoonoses, Zoonotic Agents and Food-borne Outbreaks in 2011. *EFSA J.* 2013;11:3129. Available at: <http://www.efsa.europa.eu/en/efsajournal/pub/3129> (last accessed 1 December 2015).
55. European Food Safety Authority (EFSA); The European Union Summary Report on Trends and Sources of Zoonoses, Zoonotic Agents and Food-borne Outbreaks in 2012.

EFSA J. 2014;12:3547. Available at: <http://www.efsa.europa.eu/en/efsajournal/pub/3547> (last accessed 1 December 2015).

56. Nogrady N, Kardos G, Bistyak A, Turcsanyi I, Meszaros J, Galantai Z, *et al.* Prevalence and characterization of *Salmonella* Infantis isolates originating from different points of the broiler chicken-human food chain in Hungary. *Int J Food Microbiol.* 2008;127(1-2):162-7.

57. Nogrady N, Kiraly M, Davies R, Nagy B. Multidrug resistant clones of *Salmonella* Infantis of broiler origin in Europe. *Int J Food Microbiol.* 2012;157(1):108-12.

58. Nogrady N, Toth A, Kostyak A, Paszti J, Nagy B. Emergence of multidrug-resistant clones of *Salmonella* Infantis in broiler chickens and humans in Hungary. *J Antimicrob Chemother.* 2007;60(3):645-8.

59. Kinross P, van Alphen L, Martinez Urtaza J, Struelens M, Takkinen J, Coulombier D, *et al.* Multidisciplinary investigation of a multicountry outbreak of *Salmonella* Stanley infections associated with turkey meat in the European Union, August 2011 to January 2013. *Euro Surveill.* 2014;19(19).

60. Springer B, Allerberger F, Kornschöber C. Letter to the editor: *Salmonella* Stanley outbreaks-a prompt to reevaluate existing food regulations. *Euro Surveill.* 2014;19(22).

61. European Centre for Disease Prevention and Control/European Food Safety Authority (ECDC/EFSA); Multi-country outbreak of *Salmonella* Stanley infections – Third update, 8 May 2014. Stockholm and Parma; [2014]. Available at: <http://ecdc.europa.eu/en/publications/Publications/salmonella-stanley-multi-country-outbreak-assessment-8-May-2014.pdf> (last accessed 1 December 2015).

62. Le Hello S, Bekhit A, Granier SA, Barua H, Beutlich J, Zajac M, *et al.* The global establishment of a highly-fluoroquinolone resistant *Salmonella enterica* serotype Kentucky ST198 strain. *Front Microbiol.* 2013;4:395.

63. Le Hello S, Hendriksen RS, Doublet B, Fisher I, Nielsen EM, Whichard JM, *et al.* International spread of an epidemic population of *Salmonella enterica* serotype Kentucky ST198 resistant to ciprofloxacin. *J Infect Dis.* 2011;204(5):675-84.

64. Wasyl D, Kern-Zdanowicz I, Domanska-Blicharz K, Zajac M, Hosiowski A. High-level fluoroquinolone resistant *Salmonella enterica* serovar Kentucky ST198 epidemic clone with IncA/C conjugative plasmid carrying *bla*_(CTX-M-25) gene. *Vet Microbiol.* 2015;175(1):85-91.

65. Westrell T, Monnet DL, Gossner C, Heuer O, Takkinen J. Drug-resistant *Salmonella enterica* serotype Kentucky in Europe. *Lancet Infect Dis.* 2014;14(4):270-1.

66. Mossong J, Marques P, Ragimbeau C, Huberty-Krau P, Losch S, Meyer G, *et al.* Outbreaks of monophasic *Salmonella enterica* serovar 4,[5],12:i:- in Luxembourg, 2006. *Euro Surveill.* 2007;12(6):E11-2.

- 67.** Bone A, Noel H, Le Hello S, Pihier N, Danan C, Raguenaud ME, *et al.* Nationwide outbreak of *Salmonella enterica* serotype 4,12:i:- infections in France, linked to dried pork sausage, March-May 2010. *Euro Surveill.* 2010;15(24).
- 68.** Gossner CM, van Cauteren D, Le Hello S, Weill FX, Terrien E, Tessier S, *et al.* Nationwide outbreak of *Salmonella enterica* serotype 4,[5],12:i:- infection associated with consumption of dried pork sausage, France, November to December 2011. *Euro Surveill.* 2012;17(5).
- 69.** Frank C, Werber D, Askar M, Bluemel B, Rabsch W, Simon C, *et al.* Catering risky food to those at-risk: *Salmonella* Derby outbreak among the elderly in Berlin, December 2013 to January 2014. In abstract book of ESCAIDE European Scientific Conference on Applied Infectious Disease Epidemiology. 2014. Available at: <http://ecdc.europa.eu/en/ESCAIDE/programme/abstract-book/Documents/ESCAIDE-2014-abstracts.PDF> (last accessed 1 December 2015).
- 70.** Janmohamed K, Zenner D, Little C, Lane C, Wain J, Charlett A, *et al.* National outbreak of *Salmonella* Enteritidis phage type 14b in England, September to December 2009: case-control study. *Euro Surveill.* 2011;16(15).
- 71.** Inns T, Lane C, Peters T, Dallman T, Chatt C, McFarland N, *et al.* A multi-country *Salmonella* Enteritidis phage type 14b outbreak associated with eggs from a German producer: 'near real-time' application of whole genome sequencing and food chain investigations, United Kingdom, May to September 2014. *Euro Surveill.* 2015;20(16).
- 72.** Schroeder S, Harries M, Prager R, Rabsch W, Rimek D. A prolonged outbreak of *Salmonella* Infantis associated with pork products in central Germany, April to October 2013. In abstract book of ESCAIDE European Scientific Conference on Applied Infectious Disease Epidemiology. 2014. Available at: <http://ecdc.europa.eu/en/ESCAIDE/programme/abstract-book/Documents/ESCAIDE-2014-abstracts.PDF> (last accessed 1 December 2015).
- 73.** Kuhn K, Torpdahl M, Frank C, Sigsgaard K, Ethelberg S. An outbreak of *Salmonella* Typhimurium traced back to salami, Denmark, April to June 2010. *Euro Surveill.* 2011;16(19).
- 74.** Pires SM, de Knecht L, Hald T. SCIENTIFIC/TECHNICAL REPORT submitted to EFSA Estimation of the relative contribution of different food and animal sources to human *Salmonella* infections in the European Union. 2011. Available at: http://www.efsa.europa.eu/sites/default/files/scientific_output/files/main_documents/184e.pdf (last accessed 1 September 2015).
- 75.** Department for Environment, Food and Rural Affairs (DEFRA); The 2013 UK Zoonoses Report. 2015. Available at:

- https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/447771/pb13987-zoonoses-report-2013.pdf (last accessed 1 September 2015).
- 76.** Argüello H, Rubio P, Carvajal A. *Salmonella* Control Measures at Farm in Swine Production. 2012. Available at: <http://www.intechopen.com/books/salmonella-distribution-adaptation-control-measures-and-molecular-technologies/salmonella-control-measures-at-farm-in-swine-production> (last accessed 1 December 2015). In: Annous BA, Gurtler JB, editors. *Salmonella* - Distribution, Adaptation, Control Measures and Molecular Technologies. InTech ed.
- 77.** European Food Safety Authority (EFSA); Analysis of the baseline survey on the prevalence of *Salmonella* in holdings with breeding pigs in the EU, 2008 - Part A: *Salmonella* prevalence estimates. EFSA J. 2009;7:1377. Available at: <http://www.efsa.europa.eu/en/efsajournal/pub/1377> (last accessed 1 December 2015).
- 78.** European Food Safety Authority (EFSA); Analysis of the baseline survey on the prevalence of *Salmonella* in holdings with breeding pigs, in the EU, 2008 - Part B: factors associated with *Salmonella* pen positivity. EFSA J. 2011;9:2329. Available at: <http://www.efsa.europa.eu/en/efsajournal/pub/2329> (last accessed 1 September 2015).
- 79.** Centers for Disease Control and Prevention (CDC); An atlas of *Salmonella* in the United States, 1968-2011. 2013. Available at: <http://www.cdc.gov/salmonella/pdf/salmonella-atlas-508c.pdf> (last accessed 1 December 2015).
- 80.** European Food Safety Authority (EFSA) Panel on Biological Hazards (BIOHAZ); Scientific Opinion on monitoring and assessment of the public health risk of “*Salmonella* Typhimurium-like” strains. EFSA J. 2010;8:1826. Available at: <http://www.efsa.europa.eu/en/efsajournal/pub/1826> (last accessed 1 September 2015).
- 81.** Silveira L, Marques A, Machado J. INSA Instituto Nacional de Saúde Dr Ricardo Jorge; Infecções por *Salmonella enterica* no período entre 2000-2012. Boletim Epidemiológico 1 - 2ª série. 2013. Available at: [http://www.insa.pt/sites/INSA/Portugues/PublicacoesRepositorio/Documents/observa%C3%A7%C3%B5es N%C2%BA Especial 1 2013_artigo6.pdf](http://www.insa.pt/sites/INSA/Portugues/PublicacoesRepositorio/Documents/observa%C3%A7%C3%B5es%20N%C2%BA%20Especial%201%202013_artigo6.pdf) (last accessed 1 September 2015).
- 82.** Mulvey MR, Finley R, Allen V, Ang L, Bekal S, El Bailey S, *et al.* Emergence of multidrug-resistant *Salmonella enterica* serotype 4,[5],12:i:- involving human cases in Canada: results from the Canadian Integrated Program on Antimicrobial Resistance Surveillance (CIPARS), 2003-10. J Antimicrob Chemother. 2013;68(9):1982-6.
- 83.** Switt AI, Soyer Y, Warnick LD, Wiedmann M. Emergence, distribution, and molecular and phenotypic characteristics of *Salmonella enterica* serotype 4,5,12:i:-. Foodborne Pathog Dis. 2009;6(4):407-15.

- 84.** Raguenaud ME, Le Hello S, Salah S, Weill FX, Brisabois A, Delmas G, *et al.* Epidemiological and microbiological investigation of a large outbreak of monophasic *Salmonella* Typhimurium 4,5,12:i:- in schools associated with imported beef in Poitiers, France, October 2010. *Euro Surveill.* 2012;17(40):20289.
- 85.** Hauser E, Tietze E, Helmuth R, Junker E, Blank K, Prager R, *et al.* Pork contaminated with *Salmonella enterica* serovar 4,[5],12:i:-, an emerging health risk for humans. *Appl Environ Microbiol.* 2010;76(14):4601-10.
- 86.** Hopkins KL, Kirchner M, Guerra B, Granier SA, Lucarelli C, Porrero MC, *et al.* Multiresistant *Salmonella enterica* serovar 4,[5],12:i:- in Europe: a new pandemic strain? *Euro Surveill.* 2010;15(22):19580.
- 87.** Lucarelli C, Dionisi AM, Torpdahl M, Villa L, Graziani C, Hopkins K, *et al.* Evidence for a second genomic island conferring multidrug resistance in a clonal group of strains of *Salmonella enterica* serovar Typhimurium and its monophasic variant circulating in Italy, Denmark, and the United Kingdom. *J Clin Microbiol.* 2010;48(6):2103-9.
- 88.** Mandilara G, Lambiri M, Polemis M, Passiotou M, Vatopoulos A. Phenotypic and molecular characterisation of multiresistant monophasic *Salmonella* Typhimurium (1,4,[5],12:i:-) in Greece, 2006 to 2011. *Euro Surveill.* 2013;18(22).
- 89.** Garcia P, Guerra B, Bances M, Mendoza MC, Rodicio MR. IncA/C plasmids mediate antimicrobial resistance linked to virulence genes in the Spanish clone of the emerging *Salmonella enterica* serotype 4,[5],12:i:-. *J Antimicrob Chemother.* 2011;66(3):543-9.
- 90.** Garcia P, Malorny B, Hauser E, Mendoza MC, Rodicio MR. Genetic types, gene repertoire, and evolution of isolates of the *Salmonella enterica* serovar 4,5,12:i:- Spanish clone assigned to different phage types. *J Clin Microbiol.* 2013;51(3):973-8.
- 91.** Antunes P, Machado J, Peixe L. Characterization of antimicrobial resistance and class 1 and 2 integrons in *Salmonella enterica* isolates from different sources in Portugal. *J Antimicrob Chemother.* 2006;58(2):297-304.
- 92.** Piras F, Brown DJ, Meloni D, Mureddu A, Mazzette R. Investigation of *Salmonella enterica* in Sardinian slaughter pigs: prevalence, serotype and genotype characterization. *Int J Food Microbiol.* 2011;151(2):201-9.
- 93.** Garcia-Fierro R, Montero I, Bances M, Gonzalez-Hevia MA, Rodicio MR. Antimicrobial Drug Resistance and Molecular Typing of *Salmonella enterica* Serovar Rissen from Different Sources. *Microb Drug Resist.* 2015. [Epub ahead of print].
- 94.** van Hoek AH, de Jonge R, van Overbeek WM, Bouw E, Pielaat A, Smid JH, *et al.* A quantitative approach towards a better understanding of the dynamics of *Salmonella* spp. in a pork slaughter-line. *Int J Food Microbiol.* 2012;153(1-2):45-52.

- 95.** De Busser EV, Maes D, Houf K, Dewulf J, Imberechts H, Bertrand S, *et al.* Detection and characterization of *Salmonella* in lairage, on pig carcasses and intestines in five slaughterhouses. *Int J Food Microbiol.* 2011;145(1):279-86.
- 96.** Belsue JB, Alujas AM, Porter R. Detection of high serological prevalence and comparison of different tests for *Salmonella* in pigs in Northern Ireland. *Vet Rec.* 2011;169(6):153.
- 97.** Argüello H, Carvajal A, Collazos JA, García-Feliz C, Rubio P. Prevalence and serovars of *Salmonella enterica* on pig carcasses, slaughtered pigs and the environment of four Spanish slaughterhouses. *Food Research International.* 2012b;45(2):905-12.
- 98.** Argüello H, Rubio P, Carvajal A. Occurrence and epidemiology of *Salmonella enterica* in two slaughterhouses and cutting plants in Spain. In abstract book of 40th International Conference on the Epidemiology and Control of Biological, Chemical and Physical Hazards in Pigs and Pork. 2011. Available at: <http://lib.dr.iastate.edu/cgi/viewcontent.cgi?article=1408&context=safepork> (last accessed 1 September 2015).
- 99.** Campos J, Mourão J, Silveira L, Machado J, Novais C, Peixe L, *et al.* Clonal dynamics of diverse *Salmonella enterica* serotypes from traded food products in Portugal. 2015. In abstract book of IMMEX XI International Meeting on Microbial Epidemiological Markers. Available at: <https://www.dropbox.com/s/3y15j3i6t6wfmjm/Refer%C3%Aancia%2099-%20Campos%20J.pdf?dl=0>.
- 100.** Keelara S, Scott HM, Morrow WM, Hartley CS, Griffin DL, Gebreyes WA, *et al.* Comparative phenotypic and genotypic characterization of temporally related nontyphoidal *Salmonella* isolated from human clinical cases, pigs, and the environment in North Carolina. *Foodborne Pathog Dis.* 2014;11(2):156-64.
- 101.** Pornsukarom S, Patchanee P, Erdman M, Cray PF, Wittum T, Lee J, *et al.* Comparative phenotypic and genotypic analyses of *Salmonella* Rissen that originated from food animals in Thailand and United States. *Zoonoses Public Health.* 2015;62(2):151-8.
- 102.** Schmidt JW, Brichta-Harhay DM, Kalchayanand N, Bosilevac JM, Shackelford SD, Wheeler TL, *et al.* Prevalence, enumeration, serotypes, and antimicrobial resistance phenotypes of *Salmonella enterica* isolates from carcasses at two large United States pork processing plants. *Appl Environ Microbiol.* 2012;78(8):2716-26.
- 103.** Dorn-In S, Fries R, Padungtod P, Kyule MN, Baumann MP, Srikitjakarn L, *et al.* A cross-sectional study of *Salmonella* in pre-slaughter pigs in a production compartment of northern Thailand. *Prev Vet Med.* 2009;88(1):15-23.

-
- 104.** Hendriksen RS, Vieira AR, Karlsmose S, Lo Fo Wong DM, Jensen AB, Wegener HC, *et al.* Global monitoring of *Salmonella* serovar distribution from the World Health Organization Global Foodborne Infections Network Country Data Bank: results of quality assured laboratories from 2001 to 2007. *Foodborne Pathog Dis.* 2011;8(8):887-900.
- 105.** Sanguankiat A, Pinthong R, Padungtod P, Baumann MP, Zessin KH, Srikitjakarn L, *et al.* A cross-sectional study of *Salmonella* in pork products in Chiang Mai, Thailand. *Foodborne Pathog Dis.* 2010;7(8):873-8.
- 106.** Tadee P, Boonkhot P, Pornruangwong S, Patchanee P. Comparative phenotypic and genotypic characterization of *Salmonella* spp. in pig farms and slaughterhouses in two provinces in northern Thailand. *PLoS One.* 2015;10(2):e0116581.
- 107.** Gomes-Neves E, Antunes P, Manageiro V, Gartner F, Canica M, da Costa JM, *et al.* Clinically relevant multidrug resistant *Salmonella enterica* in swine and meat handlers at the abattoir. *Vet Microbiol.* 2014;168(1):229-33.
- 108.** Fabrega A, Vila J. *Salmonella enterica* serovar Typhimurium skills to succeed in the host: virulence and regulation. *Clin Microbiol Rev.* 2013;26(2):308-41.
- 109.** Foley SL, Johnson TJ, Ricke SC, Nayak R, Danzeisen J. *Salmonella* pathogenicity and host adaptation in chicken-associated serovars. *Microbiol Mol Biol Rev.* 2013;77(4):582-607.
- 110.** Chaudhuri RR, Morgan E, Peters SE, Pleasance SJ, Hudson DL, Davies HM, *et al.* Comprehensive assignment of roles for *Salmonella* Typhimurium genes in intestinal colonization of food-producing animals. *PLoS Genet.* 2013;9(4):e1003456.
- 111.** Garaizar J, Porwollik S, Echeita A, Rementeria A, Herrera S, Wong RM, *et al.* DNA microarray-based typing of an atypical monophasic *Salmonella enterica* serovar. *J Clin Microbiol.* 2002;40(6):2074-8.
- 112.** Rajtak U, Boland F, Leonard N, Bolton D, Fanning S. Roles of diet and the acid tolerance response in survival of common *Salmonella* serotypes in feces of finishing pigs. *Appl Environ Microbiol.* 2012;78(1):110-9.
- 113.** Aviv G, Tsyba K, Steck N, Salmon-Divon M, Cornelius A, Rahav G, *et al.* A unique megaplasmid contributes to stress tolerance and pathogenicity of an emergent *Salmonella enterica* serovar Infantis strain. *Environ Microbiol.* 2014;16(4):977-94.
- 114.** Johnson TJ, Thorsness JL, Anderson CP, Lynne AM, Foley SL, Han J, *et al.* Horizontal gene transfer of a ColV plasmid has resulted in a dominant avian clonal type of *Salmonella enterica* serovar Kentucky. *PLoS One.* 2010;5(12):e15524.
- 115.** Walsh C, Fanning S. Antimicrobial resistance in foodborne pathogens-a cause for concern? *Curr Drug Targets.* 2008;9(9):808-15.
- 116.** WHO. World Health Organization (WHO); Tackling antibiotic resistance from a food safety perspective in Europe. 2011. Available at:

http://www.euro.who.int/__data/assets/pdf_file/0005/136454/e94889.pdf (last accessed 1 December 2015).

117. Cantón R, Morosini MI. Emergence and spread of antibiotic resistance following exposure to antibiotics. *FEMS Microbiol Rev.* 2011;35(5):977-91.

118. Davies J, Davies D. Origins and evolution of antibiotic resistance. *Microbiol Mol Biol Rev.* 2010;74(3):417-33.

119. Meek RW, Vyas H, Piddock LJ. Nonmedical Uses of Antibiotics: Time to Restrict Their Use? *PLoS Biol.* 2015;13(10):e1002266.

120. Seiffert SN, Hilty M, Perreten V, Endimiani A. Extended-spectrum cephalosporin-resistant Gram-negative organisms in livestock: an emerging problem for human health? *Drug Resist Updat.* 2013;16(1-2):22-45.

121. Direção Geral de Alimentação e Veterinária (DGAV); Database of Portuguese medicines, products and biocides of veterinary use. 2015. Available at: <http://medvet.dgav.pt/Listagem?classATC=QA> (last accessed 1 December 2015).

122. European Centre for Disease Prevention and Control/European Food Safety Authority/European Medicines Agency (ECDC/EFSA/EMA); First joint report on the integrated analysis of the consumption of antimicrobial agents and occurrence of antimicrobial resistance in bacteria from humans and food-producing animals. *EFSA J.* 2015;11:4006. Available at: <http://www.efsa.europa.eu/en/efsajournal/pub/4006> (last accessed 1 December 2015).

123. United States Food and Drug Administration (FDA); Estimate of human antibacterial drug sales. 2012. Available at: <http://www.fda.gov/downloads/Drugs/DrugSafety/InformationbyDrugClass/UCM319435.pdf> (last accessed 1 December 2015).

124. United States Food and Drug Administration (FDA); Annual Summary Report on Antimicrobials Sold or Distributed in 2014 for Use in Food-Producing Animals. 2015. Available at: <http://www.fda.gov/downloads/ForIndustry/UserFees/AnimalDrugUserFeeActADUFA/UCM476258.pdf> (last accessed 1 December 2015).

125. The Center for Disease Dynamics, Economics & Policy (CDDEP); The State of the World's Antibiotics, 2015. Available at: http://cddep.org/sites/default/files/swa_2015_final.pdf (last accessed 1 December 2015).

126. Van Boeckel TP, Brower C, Gilbert M, Grenfell BT, Levin SA, Robinson TP, *et al.* Global trends in antimicrobial use in food animals. *Proceedings of the National Academy of Sciences.* 2015;112(18):5649-54.

127. European Medicines Agency (EMA); Sales of veterinary antimicrobial agents in 25 EU/EEA countries in 2011 - Third ESVAC report. 2013. Available at:

- http://www.ema.europa.eu/docs/en_GB/document_library/Report/2013/10/WC500152311.pdf (last accessed 1 December 2015).
- 128.** European Medicines Agency (EMA); Sales of veterinary antimicrobial agents in 26 EU/EEA countries in 2012 - Fourth ESVAC report. 2014. Available at: http://www.ema.europa.eu/docs/en_GB/document_library/Report/2014/10/WC500175671.pdf (last accessed 1 December 2015).
- 129.** European Medicines Agency (EMA); Sales of veterinary antimicrobial agents in 26 EU/EEA countries in 2013 - Fifth ESVAC report. 2015. Available at: http://www.ema.europa.eu/docs/en_GB/document_library/Report/2015/10/WC500195687.pdf (last accessed 1 December 2015).
- 130.** Lina Maria C, Frank MLA. Resistance in bacteria of the food chain: epidemiology and control strategies. Microbial Drug Resistance: Future Medicine Ltd. 2013;136-58.
- 131.** World Health Organization (WHO); Antimicrobial Resistance Global Report on Surveillance. 2014. Available at: http://apps.who.int/iris/bitstream/10665/112642/1/9789241564748_eng.pdf (last accessed 1 December 2015).
- 132.** Magiorakos AP, Srinivasan A, Carey RB, Carmeli Y, Falagas ME, Giske CG, *et al.* Multidrug-resistant, extensively drug-resistant and pandrug-resistant bacteria: an international expert proposal for interim standard definitions for acquired resistance. Clin Microbiol Infect. 2012;18(3):268-81.
- 133.** World Health Organization (WHO); Critically important antimicrobials for human medicine, 3rd revision. 2012. Available at: http://apps.who.int/iris/bitstream/10665/77376/1/9789241504485_eng.pdf (last accessed 1 December 2015).
- 134.** Martinez JL, Coque TM, Baquero F. What is a resistance gene? Ranking risk in resistomes. Nat Rev Microbiol. 2015;13(2):116-23.
- 135.** Kahlmeter G. Defining antibiotic resistance-towards international harmonization. Ups J Med Sci. 2014;119(2):78-86.
- 136.** European Food Safety Authority (EFSA) Panel on Biological Hazards (BIOHAZ); Joint Opinion on antimicrobial resistance (AMR) focused on zoonotic infections. EFSA J. 2009;7:1372. Available at: <http://www.efsa.europa.eu/en/efsajournal/pub/1372> (last accessed 1 December 2015).
- 137.** European Food Safety Authority (EFSA) Panel on Biological Hazards (BIOHAZ); Scientific Opinion on the public health risks of bacterial strains producing extended-spectrum β -lactamases and/or AMP β -lactamases in food and food-producing animals. EFSA J. 2011;9:2322. Available at: <http://www.efsa.europa.eu/en/efsajournal/pub/2322> (last accessed 1 December 2015).

- 138.** Aarestrup FM. The livestock reservoir for antimicrobial resistance: a personal view on changing patterns of risks, effects of interventions and the way forward. *Philos Trans R Soc Lond B Biol Sci.* 2015;370(1670):20140085.
- 139.** Delsol AA, Anjum M, Woodward MJ, Sunderland J, Roe JM. The effect of chlortetracycline treatment and its subsequent withdrawal on multi-resistant *Salmonella enterica* serovar Typhimurium DT104 and commensal *Escherichia coli* in the pig. *J Appl Microbiol.* 2003;95(6):1226-34.
- 140.** Antunes P, Machado J, Peixe L. Illegal use of nitrofurans in food animals: Contribution to human salmonellosis? *Clin Microbiol Infect.* 2006;12(11):1047-9.
- 141.** Centers for Disease Control and Prevention/U.S. Department of Agriculture/Food and Drug Administration (CDC/USDA/FDA); The National Antimicrobial Resistance Monitoring System (NARMS): The 2012-2013 Integrated NARMS Report. 2015. Available at: <http://www.fda.gov/downloads/AnimalVeterinary/SafetyHealth/AntimicrobialResistance/NationalAntimicrobialResistanceMonitoringSystem/UCM453398.pdf> (last accessed 1 December 2015).
- 142.** Lee HY, Su LH, Tsai MH, Kim SW, Chang HH, Jung SI, *et al.* High rate of reduced susceptibility to ciprofloxacin and ceftriaxone among nontyphoid *Salmonella* clinical isolates in Asia. *Antimicrob Agents Chemother.* 2009;53(6):2696-9.
- 143.** Van TT, Nguyen HN, Smooker PM, Coloe PJ. The antibiotic resistance characteristics of non-typhoidal *Salmonella enterica* isolated from food-producing animals, retail meat and humans in South East Asia. *Int J Food Microbiol.* 2012;154(3):98-106.
- 144.** Rodriguez-Martinez JM, Cano ME, Velasco C, Martinez-Martinez L, Pascual A. Plasmid-mediated quinolone resistance: an update. *J Infect Chemother.* 2011;17(2):149-82.
- 145.** Abatcha MG, Zakaria Z, Kaur DG, Thong KL. Review Article: A trends of *Salmonella* and antibiotic resistance. *Advances in Life Science and Technology* 2014;17:9-21.
- 146.** Blair JM, Webber MA, Baylay AJ, Ogbolu DO, Piddock LJ. Molecular mechanisms of antibiotic resistance. *Nat Rev Microbiol.* 2015;13(1):42-51.
- 147.** Kohanski MA, Dwyer DJ, Collins JJ. How antibiotics kill bacteria: from targets to networks. *Nat Rev Microbiol.* 2010;8(6):423-35.
- 148.** Partridge SR. Resistance mechanisms in *Enterobacteriaceae*. *Pathology.* 2015;47(3):276-84.
- 149.** Lewis K. Platforms for antibiotic discovery. *Nat Rev Drug Discov.* 2013;12(5):371-87.

- 150.** Bush K. The ABCD's of β -lactamase nomenclature. *J Infect Chemother.* 2013;19(4):549-59.
- 151.** Bush K. Proliferation and significance of clinically relevant β -lactamases. *Ann N Y Acad Sci.* 2013;1277:84-90.
- 152.** European Food Safety Authority (EFSA) Panel on Biological Hazards (BIOHAZ); Scientific Opinion on Carbapenem resistance in food animal ecosystems. *EFSA J.* 2013;11:3501. Available at: <http://www.efsa.europa.eu/en/efsajournal/pub/3501> (last accessed 1 December 2015).
- 153.** Guerra B, Fischer J, Helmuth R. An emerging public health problem: acquired carbapenemase-producing microorganisms are present in food-producing animals, their environment, companion animals and wild birds. *Vet Microbiol.* 2014;171(3-4):290-7.
- 154.** Cantón R, Gonzalez-Alba JM, Galan JC. CTX-M Enzymes: Origin and Diffusion. *Front Microbiol.* 2012;3:110.
- 155.** Tamang MD, Gurung M, Nam HM, Moon DC, Kim SR, Jang GC, *et al.* Prevalence and characterization of *Salmonella* in pigs from conventional and organic farms and first report of *S. serovar* 1,4,[5],12:i:- from Korea. *Vet Microbiol.* 2015;178(1-2):119-24.
- 156.** Rodriguez I, Jahn S, Schroeter A, Malorny B, Helmuth R, Guerra B. Extended-spectrum β -lactamases in German isolates belonging to the emerging monophasic *Salmonella enterica* subsp. *enterica* serovar Typhimurium 4,[5],12:i:- European clone. *J Antimicrob Chemother.* 2012;67(2):505-8.
- 157.** Freire Martin I, AbuOun M, Reichel R, La Ragione RM, Woodward MJ. Sequence analysis of a CTX-M-1 IncI1 plasmid found in *Salmonella* 4,5,12:i:-, *Escherichia coli* and *Klebsiella pneumoniae* on a UK pig farm. *J Antimicrob Chemother.* 2014;69(8):2098-101.
- 158.** Weill FX, Lailier R, Praud K, Kerouanton A, Fabre L, Brisabois A, *et al.* Emergence of extended-spectrum- β -lactamase (CTX-M-9)-producing multiresistant strains of *Salmonella enterica* serotype Virchow in poultry and humans in France. *J Clin Microbiol.* 2004;42(12):5767-73.
- 159.** Bertrand S, Weill FX, Cloeckaert A, Vrints M, Mairiaux E, Praud K, *et al.* Clonal emergence of extended-spectrum β -lactamase (CTX-M-2)-producing *Salmonella enterica* serovar Virchow isolates with reduced susceptibilities to ciprofloxacin among poultry and humans in Belgium and France (2000 to 2003). *J Clin Microbiol.* 2006;44(8):2897-903.
- 160.** Riano I, Moreno MA, Teshager T, Saenz Y, Dominguez L, Torres C. Detection and characterization of extended-spectrum β -lactamases in *Salmonella enterica* strains of healthy food animals in Spain. *J Antimicrob Chemother.* 2006;58(4):844-7.
- 161.** Garcia Fernandez A, Cloeckaert A, Bertini A, Praud K, Doublet B, Weill FX, *et al.* Comparative analysis of IncHI2 plasmids carrying *bla*_{CTX-M-2} or *bla*_{CTX-M-9} from *Escherichia*

coli and *Salmonella enterica* strains isolated from poultry and humans. Antimicrob Agents Chemother. 2007;51(11):4177-80.

162. Cloeckaert A, Praud K, Doublet B, Bertini A, Carattoli A, Butaye P, *et al.* Dissemination of an extended-spectrum- β -lactamase *bla*_{TEM-52} gene-carrying IncI1 plasmid in various *Salmonella enterica* serovars isolated from poultry and humans in Belgium and France between 2001 and 2005. Antimicrob Agents Chemother. 2007;51(5):1872-5.

163. Smet A, Martel A, Persoons D, Dewulf J, Heyndrickx M, Cloeckaert A, *et al.* Comparative analysis of extended-spectrum- β -lactamase-carrying plasmids from different members of *Enterobacteriaceae* isolated from poultry, pigs and humans: evidence for a shared β -lactam resistance gene pool? J Antimicrob Chemother. 2009;63(6):1286-8.

164. Riano I, Garcia-Campello M, Saenz Y, Alvarez P, Vinue L, Lantero M, *et al.* Occurrence of extended-spectrum β -lactamase-producing *Salmonella enterica* in northern Spain with evidence of CTX-M-9 clonal spread among animals and humans. Clin Microbiol Infect. 2009;15(3):292-5.

165. Dierikx C, van Essen-Zandbergen A, Veldman K, Smith H, Mevius D. Increased detection of extended spectrum β -lactamase producing *Salmonella enterica* and *Escherichia coli* isolates from poultry. Vet Microbiol. 2010;145(3-4):273-8.

166. Leverstein-van Hall MA, Dierikx CM, Cohen Stuart J, Voets GM, van den Munckhof MP, van Essen-Zandbergen A, *et al.* Dutch patients, retail chicken meat and poultry share the same ESBL genes, plasmids and strains. Clin Microbiol Infect. 2011;17(6):873-80.

167. Jacoby GA. AmpC β -lactamases. Clin Microbiol Rev. 2009;22(1):161-82.

168. Andrysiak AK, Olson AB, Tracz DM, Dore K, Irwin R, Ng LK, *et al.* Genetic characterization of clinical and agri-food isolates of multi drug resistant *Salmonella enterica* serovar Heidelberg from Canada. BMC Microbiol. 2008;8:89.

169. Martin LC, Weir EK, Poppe C, Reid-Smith RJ, Boerlin P. Characterization of *bla*_{CMY-2} plasmids in *Salmonella* and *Escherichia coli* isolates from food animals in Canada. Appl Environ Microbiol. 2012;78(4):1285-7.

170. de Jong A, Smet A, Ludwig C, Stephan B, De Graef E, Vanrobaeys M, *et al.* Antimicrobial susceptibility of *Salmonella* isolates from healthy pigs and chickens (2008-2011). Vet Microbiol. 2014;171(3-4):298-306.

171. Lee KE, Lim SI, Choi HW, Lim SK, Song JY, An DJ. Plasmid-mediated AmpC β -lactamase (CMY-2) gene in *Salmonella* Typhimurium isolated from diarrheic pigs in South Korea. BMC Res Notes. 2014;7:329.

- 172.** Le Hello S, Harrois D, Bouchrif B, Sontag L, Elhani D, Guibert V, *et al.* Highly drug-resistant *Salmonella enterica* serotype Kentucky ST198-X1: a microbiological study. *Lancet Infect Dis.* 2013;13(8):672-9.
- 173.** Irfan S, Khan E, Jabeen K, Bhawan P, Hopkins KL, Day M, *et al.* Clinical isolates of *Salmonella enterica* serovar Agona producing NDM-1 metallo- β -lactamase: first report from Pakistan. *J Clin Microbiol.* 2015;53(1):346-8.
- 174.** Seiffert SN, Perreten V, Johannes S, Droz S, Bodmer T, Endimiani A. OXA-48 carbapenemase-producing *Salmonella enterica* serovar Kentucky isolate of sequence type 198 in a patient transferred from Libya to Switzerland. *Antimicrob Agents Chemother.* 2014;58(4):2446-9.
- 175.** Rodriguez E, Bautista A, Barrero L. First report of a *Salmonella enterica* serovar Typhimurium isolate with carbapenemase (KPC-2) in Colombia. *Antimicrob Agents Chemother.* 2014;58(2):1263-4.
- 176.** Ktari S, Le Hello S, Ksibi B, Courdavault L, Mnif B, Maalej S, *et al.* Carbapenemase-producing *Salmonella enterica* serotype Kentucky ST198, North Africa. *J Antimicrob Chemother.* 2015;70(12):3405-7.
- 177.** Jure MA, Duprilot M, Musa HE, Lopez C, de Castillo MC, Weill FX, *et al.* Emergence of KPC-2-Producing *Salmonella enterica* Serotype Schwarzengrund in Argentina. *Antimicrob Agents Chemother.* 2014;58(10):6335-6.
- 178.** Sotillo A, Munoz-Velez M, Santamaria ML, Ruiz-Carrascoso G, Garcia-Bujalance S, Gomez-Gil R, *et al.* Emergence of VIM-1-producing *Salmonella enterica* serovar Typhimurium in a paediatric patient. *J Med Microbiol.* 2015;64(12):1541-3.
- 179.** Day MR, Meunier D, Doumith M, de Pinna E, Woodford N, Hopkins KL. Carbapenemase-producing *Salmonella enterica* isolates in the UK. *J Antimicrob Chemother.* 2015;70(7):2165-7.
- 180.** Fischer J, Schmoger S, Jahn S, Helmuth R, Guerra B. NDM-1 carbapenemase-producing *Salmonella enterica* subsp. *enterica* serovar Corvallis isolated from a wild bird in Germany. *J Antimicrob Chemother.* 2013;68(12):2954-6.
- 181.** Fischer J, Rodriguez I, Schmoger S, Friese A, Roesler U, Helmuth R, *et al.* *Salmonella enterica* subsp. *enterica* producing VIM-1 carbapenemase isolated from livestock farms. *J Antimicrob Chemother.* 2013;68(2):478-80.
- 182.** Falgenhauer L, Ghosh H, Guerra B, Yao Y, Fritzenwanker M, Fischer J, *et al.* Comparative genome analysis of IncHI2 VIM-1 carbapenemase-encoding plasmids of *Escherichia coli* and *Salmonella enterica* isolated from a livestock farm in Germany. *Vet Microbiol.* 2015. [Epub ahead of print].
- 183.** Silva KC, Fontes LC, Moreno AM, Astolfi-Ferreira CS, Ferreira AJ, Lincopan N. Emergence of extended-spectrum- β -lactamase CTX-M-2-producing *Salmonella enterica*

serovars Schwarzengrund and Agona in poultry farms. *Antimicrob Agents Chemother.* 2013;57(7):3458-9.

184. Fitch FM, Carmo-Rodrigues MS, Oliveira VG, Gaspari MV, Dos Santos A, de Freitas JB, *et al.* β -Lactam Resistance Genes: Characterization, Epidemiology, and First Detection of *bla*_{CTX-M-1} and *bla*_{CTX-M-14} in *Salmonella* spp. Isolated from Poultry in Brazil-Brazil Ministry of Agriculture's Pathogen Reduction Program. *Microb Drug Resist.* 2015. [Epub ahead of print].

185. Rodriguez I, Barownick W, Helmuth R, Mendoza MC, Rodicio MR, Schroeter A, *et al.* Extended-spectrum β -lactamases and AmpC β -lactamases in ceftiofur-resistant *Salmonella enterica* isolates from food and livestock obtained in Germany during 2003-07. *J Antimicrob Chemother.* 2009;64(2):301-9.

186. Keelara S, Thakur S. Dissemination of plasmid-encoded AmpC β -lactamases in antimicrobial resistant *Salmonella* serotypes originating from humans, pigs and the swine environment. *Vet Microbiol.* 2014;173(1-2):76-83.

187. Hasman H, Mevius D, Veldman K, Olesen I, Aarestrup FM. β -Lactamases among extended-spectrum β -lactamase (ESBL)-resistant *Salmonella* from poultry, poultry products and human patients in The Netherlands. *J Antimicrob Chemother.* 2005;56(1):115-21.

188. Chiaretto G, Zavagnin P, Bettini F, Mancin M, Minorello C, Saccardin C, *et al.* Extended spectrum beta-lactamase SHV-12-producing *Salmonella* from poultry. *Vet Microbiol.* 2008;128(3-4):406-13.

189. Wittum TE, Mollenkopf DF, Erdman MM. Detection of *Salmonella enterica* isolates producing CTX-M Cephalosporinase in U.S. livestock populations. *Appl Environ Microbiol.* 2012;78(20):7487-91.

190. Hopkins KL, Liebana E, Villa L, Batchelor M, Threlfall EJ, Carattoli A. Replicon typing of plasmids carrying CTX-M or CMY beta-lactamases circulating among *Salmonella* and *Escherichia coli* isolates. *Antimicrob Agents Chemother.* 2006;50(9):3203-6.

191. Matsumoto Y, Kitazume H, Yamada M, Ishiguro Y, Muto T, Izumiya H, *et al.* CTX-M-14 type beta-lactamase producing *Salmonella enterica* serovar Enteritidis isolated from imported chicken meat. *Jpn J Infect Dis.* 2007;60(4):236-8.

192. Li L, Liao XP, Liu ZZ, Huang T, Li X, Sun J, *et al.* Co-spread of *oqxAB* and *bla*_{CTX-M-9G} in non-Typhi *Salmonella enterica* isolates mediated by ST2-IncHI2 plasmids. *Int J Antimicrob Agents.* 2014;44(3):263-8.

193. Tamang MD, Nam HM, Kim TS, Jang GC, Jung SC, Lim SK. Emergence of extended-spectrum beta-lactamase (CTX-M-15 and CTX-M-14)-producing nontyphoid

- Salmonella* with reduced susceptibility to ciprofloxacin among food animals and humans in Korea. J Clin Microbiol. 2011;49(7):2671-5.
- 194.** Kang MS, Kwon YK, Oh JY, Kim MJ, Call DR, An BK, *et al.* Evidence for recent acquisition and successful transmission of *bla*_(CTX-M-15) in *Salmonella enterica* in South Korea. Antimicrob Agents Chemother. 2013;57(5):2383-7.
- 195.** Clemente L, Manageiro V, Jones-Dias D, Correia I, Themudo P, Albuquerque T, *et al.* Antimicrobial susceptibility and oxymino-beta-lactam resistance mechanisms in *Salmonella enterica* and *Escherichia coli* isolates from different animal sources. Res Microbiol. 2015;166(7):574-83.
- 196.** Gonzalez F, Araque M. Association of Transferable Quinolone Resistance Determinant *qnrB19* with Extended-Spectrum β -Lactamases in *Salmonella* GIVE and *Salmonella* Heidelberg in Venezuela. Int J Microbiol. 2013;2013:628185.
- 197.** Pouget JG, Coutinho FJ, Reid-Smith RJ, Boerlin P. Characterization of *bla*_(SHV) genes on plasmids from *Escherichia coli* and *Salmonella enterica* isolates from Canadian food animals (2006-2007). Appl Environ Microbiol. 2013;79(12):3864-6.
- 198.** Glenn LM, Lindsey RL, Folster JP, Pecic G, Boerlin P, Gilmour MW, *et al.* Antimicrobial resistance genes in multidrug-resistant *Salmonella enterica* isolated from animals, retail meats, and humans in the United States and Canada. Microb Drug Resist. 2013;19(3):175-84.
- 199.** Jiang HX, Song L, Liu J, Zhang XH, Ren YN, Zhang WH, *et al.* Multiple transmissible genes encoding fluoroquinolone and third-generation cephalosporin resistance co-located in non-typhoidal *Salmonella* isolated from food-producing animals in China. Int J Antimicrob Agents. 2014;43(3):242-7.
- 200.** Lai J, Wang Y, Shen J, Li R, Han J, Foley SL, *et al.* Unique class 1 integron and multiple resistance genes co-located on IncHI2 plasmid is associated with the emerging multidrug resistance of *Salmonella* Indiana isolated from chicken in China. Foodborne Pathog Dis. 2013;10(7):581-8.
- 201.** Lin D, Chen K, Wai-Chi Chan E, Chen S. Increasing prevalence of ciprofloxacin-resistant food-borne *Salmonella* strains harboring multiple PMQR elements but not target gene mutations. Sci Rep. 2015;5:14754.
- 202.** Bai L, Lan R, Zhang X, Cui S, Xu J, Guo Y, *et al.* Prevalence of *Salmonella* Isolates from Chicken and Pig Slaughterhouses and Emergence of Ciprofloxacin and Cefotaxime Co-Resistant *S. enterica* Serovar Indiana in Henan, China. PLoS One. 2015;10(12):e0144532.
- 203.** Rayamajhi N, Jung BY, Cha SB, Shin MK, Kim A, Kang MS, *et al.* Antibiotic resistance patterns and detection of *bla*_{DHA-1} in *Salmonella* species isolates from chicken farms in South Korea. Appl Environ Microbiol. 2010;76(14):4760-4.

- 204.** Franco A, Leekitcharoenphon P, Feltrin F, Alba P, Cordaro G, Iurescia M, *et al.* Emergence of a Clonal Lineage of Multidrug-Resistant ESBL-Producing *Salmonella* Infantis Transmitted from Broilers and Broiler Meat to Humans in Italy between 2011 and 2014. PLoS One. 2015;10(12):e0144802.
- 205.** Noda T, Murakami K, Etoh Y, Okamoto F, Yatsuyanagi J, Sera N, *et al.* Increase in resistance to extended-spectrum cephalosporins in *Salmonella* isolated from retail chicken products in Japan. PLoS One. 2015;10(2):e0116927.
- 206.** Shahada F, Chuma T, Dahshan H, Akiba M, Sueyoshi M, Okamoto K. Detection and characterization of extended-spectrum β -lactamase (TEM-52)-producing *Salmonella* serotype Infantis from broilers in Japan. Foodborne Pathog Dis. 2010;7(5):515-21.
- 207.** Dahshan H, Chuma T, Shahada F, Akiba M, Fujimoto H, Akasaka K, *et al.* Characterization of antibiotic resistance and the emergence of AmpC-producing *Salmonella* Infantis from pigs. J Vet Med Sci. 2010;72(11):1437-42.
- 208.** Boyle F, Morris D, O'Connor J, Delappe N, Ward J, Cormican M. First report of extended-spectrum- β -lactamase-producing *Salmonella enterica* serovar Kentucky isolated from poultry in Ireland. Antimicrob Agents Chemother. 2010;54(1):551-3.
- 209.** Cloeckaert A, Praud K, Lefevre M, Doublet B, Pardos M, Granier SA, *et al.* IncI1 plasmid carrying extended-spectrum- β -lactamase gene *bla*_{CTX-M-1} in *Salmonella enterica* isolates from poultry and humans in France, 2003 to 2008. Antimicrob Agents Chemother. 2010;54(10):4484-6.
- 210.** Fernandes SA, Paterson DL, Ghilardi-Rodrigues AC, Adams-Haduch JM, Tavechio AT, Doi Y. CTX-M-2-producing *Salmonella* Typhimurium isolated from pediatric patients and poultry in Brazil. Microb Drug Resist. 2009;15(4):317-21.
- 211.** Hasman H, Mevius D, Veldman K, Olesen I, Aarestrup FM. β -lactamases among extended-spectrum β -lactamase (ESBL)-resistant *Salmonella* from poultry, poultry products and human patients in the Netherlands. J Antimicrob Chemother. 2005;56(1):115-21.
- 212.** O'Mahony R, Quinn T, Drudy D, Walsh C, Whyte P, Mattar S, *et al.* Antimicrobial resistance in nontyphoidal *Salmonella* from food sources in Colombia: evidence for an unusual plasmid-localized class 1 integron in serotypes Typhimurium and Anatum. Microb Drug Resist. 2006;12(4):269-77.
- 213.** Zaidi MB, Leon V, Canche C, Perez C, Zhao S, Hubert SK, *et al.* Rapid and widespread dissemination of multidrug-resistant *bla*_{CMY-2} *Salmonella* Typhimurium in Mexico. J Antimicrob Chemother. 2007;60(2):398-401.
- 214.** Politi L, Tassios PT, Lambiri M, Kansouzidou A, Pasiotou M, Vatopoulos AC, *et al.* Repeated occurrence of diverse extended-spectrum β -lactamases in minor serotypes of food-borne *Salmonella enterica* subsp. *enterica*. J Clin Microbiol. 2005;43(7):3453-6.

- 215.** Cardinale E, Colbachini P, Perrier-Gros-Claude JD, Gassama A, Aidara-Kane A. Dual emergence in food and humans of a novel multiresistant serotype of *Salmonella* in Senegal: *Salmonella enterica* subsp. *enterica* serotype 35:c:1,2. J Clin Microbiol. 2001;39(6):2373-4.
- 216.** Luo Y, Li J, Meng Y, Ma Y, Hu C, Jin S, *et al.* Joint effects of topoisomerase alterations and plasmid-mediated quinolone-resistant determinants in *Salmonella enterica* Typhimurium. Microb Drug Resist. 2011;17(1):1-5.
- 217.** Veldman K, Cavaco LM, Mevius D, Battisti A, Franco A, Botteldoorn N, *et al.* International collaborative study on the occurrence of plasmid-mediated quinolone resistance in *Salmonella enterica* and *Escherichia coli* isolated from animals, humans, food and the environment in 13 European countries. J Antimicrob Chemother. 2011;66(6):1278-86.
- 218.** Li L, Liao X, Yang Y, Sun J, Li L, Liu B, *et al.* Spread of *oqxAB* in *Salmonella enterica* serotype Typhimurium predominantly by IncHI2 plasmids. J Antimicrob Chemother. 2013;68(10):2263-8.
- 219.** Hooper DC, Jacoby GA. Mechanisms of drug resistance: quinolone resistance. Ann N Y Acad Sci. 2015;1354(1):12-31.
- 220.** Wasyl D, Hoszowski A, Zajac M. Prevalence and characterisation of quinolone resistance mechanisms in *Salmonella* spp. Vet Microbiol. 2014;171(3-4):307-14.
- 221.** Cavaco LM, Korsgaard H, Sorensen G, Aarestrup FM. Plasmid-mediated quinolone resistance due to *qnrB5* and *qnrS1* genes in *Salmonella enterica* serovars Newport, Hadar and Saintpaul isolated from turkey meat in Denmark. J Antimicrob Chemother. 2008;62(3):632-4.
- 222.** Jones-Dias D, Manageiro V, Francisco AP, Martins AP, Domingues G, Louro D, *et al.* Assessing the molecular basis of transferable quinolone resistance in *Escherichia coli* and *Salmonella* spp. from food-producing animals and food products. Vet Microbiol. 2013;167(3-4):523-31.
- 223.** Kehrenberg C, Friederichs S, de Jong A, Michael GB, Schwarz S. Identification of the plasmid-borne quinolone resistance gene *qnrS* in *Salmonella enterica* serovar Infantis. J Antimicrob Chemother. 2006;58(1):18-22.
- 224.** Karczmarczyk M, Martins M, McCusker M, Mattar S, Amaral L, Leonard N, *et al.* Characterization of antimicrobial resistance in *Salmonella enterica* food and animal isolates from Colombia: identification of a *qnrB19*-mediated quinolone resistance marker in two novel serovars. FEMS Microbiol Lett. 2010;313(1):10-9.
- 225.** Dolejska M, Villa L, Hasman H, Hansen L, Carattoli A. Characterization of IncN plasmids carrying *bla*_{CTX-M-1} and *qnr* genes in *Escherichia coli* and *Salmonella* from animals, the environment and humans. J Antimicrob Chemother. 2013;68(2):333-9.

- 226.** Hopkins KL, Wootton L, Day MR, Threlfall EJ. Plasmid-mediated quinolone resistance determinant *qnrS1* found in *Salmonella enterica* strains isolated in the UK. *J Antimicrob Chemother.* 2007;59(6):1071-5.
- 227.** Avsaroglu MD, Helmuth R, Junker E, Hertwig S, Schroeter A, Akcelik M, *et al.* Plasmid-mediated quinolone resistance conferred by *qnrS1* in *Salmonella enterica* serovar Virchow isolated from Turkish food of avian origin. *J Antimicrob Chemother.* 2007;60(5):1146-50.
- 228.** Kim JH, Cho JK, Kim KS. Prevalence and characterization of plasmid-mediated quinolone resistance genes in *Salmonella* isolated from poultry in Korea. *Avian Pathol.* 2013;42(3):221-9.
- 229.** Lucarelli C, Dionisi AM, Filetici E, Owczarek S, Luzzi I, Villa L. Nucleotide sequence of the chromosomal region conferring multidrug resistance (R-type ASSuT) in *Salmonella* Typhimurium and monophasic *Salmonella* Typhimurium strains. *J Antimicrob Chemother.* 2012;67(1):111-4.
- 230.** Wong MH, Yan M, Chan EW, Liu LZ, Kan B, Chen S. Expansion of *Salmonella* Typhimurium ST34 clone carrying multiple resistance determinants in China. *Antimicrob Agents Chemother.* 2013;57(9):4599-4601.
- 231.** Wong MH, Yan M, Chan EW, Biao K, Chen S. Emergence of clinical *Salmonella enterica* serovar Typhimurium isolates with concurrent resistance to ciprofloxacin, ceftriaxone, and azithromycin. *Antimicrob Agents Chemother.* 2014;58(7):3752-6.
- 232.** Frost LS, Leplae R, Summers AO, Toussaint A. Mobile genetic elements: the agents of open source evolution. *Nat Rev Microbiol.* 2005;3(9):722-32.
- 233.** Partridge SR. Analysis of antibiotic resistance regions in Gram-negative bacteria. *FEMS Microbiol Rev.* 2011;35(5):820-55.
- 234.** Silva C, Wiesner M, Calva E. The Importance of Mobile Genetic Elements in the Evolution of *Salmonella*: Pathogenesis, Antibiotic Resistance and Host Adaptation. In: Kumar Y, editor. *Salmonella - A Diversified Superbug.* 2012. Available at: <http://www.intechopen.com/books/salmonella-a-diversified-superbug/the-importance-of-mobile-genetic-elements-in-the-evolution-of-salmonella-pathogenesis-antibiotic-res>.
- 235.** Carattoli A. Resistance plasmid families in *Enterobacteriaceae*. *Antimicrob Agents Chemother.* 2009;53(6):2227-38.
- 236.** Carattoli A. Plasmids and the spread of resistance. *Int J Med Microbiol.* 2013;303(6-7):298-304.
- 237.** Carattoli A, Bertini A, Villa L, Falbo V, Hopkins KL, Threlfall EJ. Identification of plasmids by PCR-based replicon typing. *J Microbiol Methods.* 2005;63(3):219-28.

- 238.** Juhas M, van der Meer JR, Gaillard M, Harding RM, Hood DW, Crook DW. Genomic islands: tools of bacterial horizontal gene transfer and evolution. *FEMS Microbiol Rev.* 2009;33(2):376-93.
- 239.** Langille MG, Hsiao WW, Brinkman FS. Detecting genomic islands using bioinformatics approaches. *Nat Rev Microbiol.* 2010;8(5):373-82.
- 240.** Hall RM. *Salmonella* genomic islands and antibiotic resistance in *Salmonella enterica*. *Future Microbiol.* 2010;5(10):1525-38.
- 241.** Douard G, Praud K, Cloeckaert A, Doublet B. The *Salmonella* genomic island 1 is specifically mobilized in trans by the IncA/C multidrug resistance plasmid family. *PLoS One.* 2010;5(12):e15302.
- 242.** Doublet B, Boyd D, Mulvey MR, Cloeckaert A. The *Salmonella* genomic island 1 is an integrative mobilizable element. *Mol Microbiol.* 2005;55(6):1911-24.
- 243.** Mazel D. Integrons: agents of bacterial evolution. *Nat Rev Microbiol.* 2006;4(8):608-20.
- 244.** Antunes P, Machado J, Sousa JC, Peixe L. Dissemination amongst humans and food products of animal origin of a *Salmonella* Typhimurium clone expressing an integron-borne OXA-30 beta-lactamase. *J Antimicrob Chemother.* 2004;54(2):429-34.
- 245.** Scientific Committee on Emerging and Newly Identified Health Risks (SCENIHR). Assessment of the Antibiotic Resistance Effects of Biocides. 2009. Available at: http://ec.europa.eu/health/ph_risk/committees/04_scenihr/docs/scenihr_o_o21.pdf (last accessed 1 December 2015).
- 246.** Seiler C, Berendonk TU. Heavy metal driven co-selection of antibiotic resistance in soil and water bodies impacted by agriculture and aquaculture. *Front Microbiol.* 2012;3:399.
- 247.** Commission Regulation (EC) No 1831/2003 of the European Parliament and of the Council of 22 September 2003 on additives for use in animal nutrition [2003]. *OJ L268/29.*
- 248.** Scientific Committee on Emerging and Newly Identified Health Risks (SCENIHR). Research strategy to address the knowledge gaps on the antimicrobial resistance effects of biocides. 2010. Available at: http://ec.europa.eu/health/scientific_committees/emerging/docs/scenihr_o_o28.pdf (last accessed 1 December 2015).
- 249.** Baker-Austin C, Wright MS, Stepanauskas R, McArthur JV. Co-selection of antibiotic and metal resistance. *Trends Microbiol.* 2006;14(4):176-82.
- 250.** Gnanadhas DP, Marathe SA, Chakravorty D. Biocides-resistance, cross-resistance mechanisms and assessment. *Expert Opin Investig Drugs.* 2013;22(2):191-206.

- 251.** Directive 98/8/EC of the European Parliament and of the Council of 16 February 1998 concerning the placing of biocidal products on the market [1998]. OJ L123/1. 1998.
- 252.** Commission Regulation (EC) No 528/2012 of the European Parliament and of the Council of 22 May 2012 concerning the making available on the market and use of biocidal products [2012]. OJ L167/1.
- 253.** Buffet-Bataillon S, Tattevin P, Maillard JY, Bonnaure-Mallet M, Jolivet-Gougeon A. Efflux pump induction by quaternary ammonium compounds and fluoroquinolone resistance in bacteria. *Future Microbiol.* 2016;11:81-92.
- 254.** Condell O, Iversen C, Cooney S, Power KA, Walsh C, Burgess C, *et al.* Efficacy of biocides used in the modern food industry to control *Salmonella enterica*, and links between biocide tolerance and resistance to clinically relevant antimicrobial compounds. *Appl Environ Microbiol.* 2012;78(9):3087-97.
- 255.** Commission Regulation (EC) No 1003/2005 of 30 June 2005 implementing Regulation (EC) No 2160/2003 as regards a Community target for the reduction of the prevalence of certain *Salmonella* serotypes in breeding flocks of *Gallus gallus* and amending Regulation (EC) No 2160/2003 [2005]. OJ L170/12.
- 256.** Commission Regulation (EC) No 1086/2011 of 27 October 2011 amending Annex II to Regulation (EC) No 2160/2003 of the European Parliament and of the Council and Annex I to Commission Regulation (EC) No 2073/2005 as regards *Salmonella* in fresh poultry meat [2011]. OJ L281/7.
- 257.** Commission Regulation (EC) No 209/2013 of 11 March 2013 amending Regulation (EC) No 2073/2005 as regards microbiological criteria for sprouts and the sampling rules for poultry carcasses and fresh poultry meat [2013]. OJ L68/19.
- 258.** Commission Regulation (EC) No 217/2014 of 7 March 2014 amending Regulation (EC) No 2073/2005 as regards *Salmonella* in pig carcasses [2014]. OJ L69/93.
- 259.** Fanning S. Fanning S. Altered tolerance to biocides: links to antibiotic resistance?. Oral presentation at: European Symposium on Food Safety; 2011 May 19; The Netherlands. PowerPoint presentation available from: <https://www.dropbox.com/s/njirw7nfoxsc8jh/Refer%C3%A4ncia%20259%20-%20Fanning%20S.pdf?dl=0>.
- 260.** Wales A, Davies R. Co-Selection of Resistance to Antibiotics, Biocides and Heavy Metals, and Its Relevance to Foodborne Pathogens. *Antibiotics.* 2015;4(4):567.
- 261.** Yoshinaga M, Rosen BP. A CAs lyase for degradation of environmental organoarsenical herbicides and animal husbandry growth promoters. *Proc Natl Acad Sci U S A.* 2014;111(21):7701-6.
- 262.** Maillard JY, Hartemann P. Silver as an antimicrobial: facts and gaps in knowledge. *Crit Rev Microbiol.* 2013;39(4):373-83.

- 263.** Hobman JL, Crossman LC. Bacterial antimicrobial metal ion resistance. *J Med Microbiol.* 2015;64(Pt 5):471-97.
- 264.** Commission Regulation (EC) No 1334/2003 of 25 July 2003 amending the conditions for authorisation of a number of additives in feedingstuffs belonging to the group of trace elements [2003]. *OJ L187/11.*
- 265.** European Commission (EU). SCAN—Scientific Committee for Animal Nutrition; Opinion of the Scientific Committee for Animal Nutrition on the Use of Copper in Feedingstuffs. 2003. Available at: http://ec.europa.eu/food/fs/sc/scan/out115_en.pdf (last accessed 1 December 2015).
- 266.** Norwegian Scientific Committee for Food Safety (VKM); Zinc and copper in pig and poultry production - fate and effects in the food chain and the environment. 2014. Available at: <http://www.english.vkm.no/dav/3b1b6769dd.pdf> (last accessed 1 December 2015).
- 267.** Yazdankhah S, Rudi K, Bernhoft A. Zinc and copper in animal feed - development of resistance and co-resistance to antimicrobial agents in bacteria of animal origin. *Microb Ecol Health Dis.* 2014;25.
- 268.** European Food Safety Authority (EFSA) Panel on Additives and Products or Substances used in Animal Feed (FEEDAP); Scientific Opinion on the potential reduction of the currently authorised maximum zinc content in complete feed. *EFSA J.* 2014;12:3668. Available at: <http://www.efsa.europa.eu/en/efsajournal/pub/3668> (last accessed 1 December 2015).
- 269.** Jacela JY, DeRouchey JM, Tokach MD, Goodband RD, Nelssen JL, Renter DG, *et al.* Feed additives for swine: Fact sheets – high dietary levels of copper and zinc for young pigs, and phytase. *J Swine Health Prod.* 2010;18(2):87–91.
- 270.** European Food Safety Authority (EFSA) Panel on Additives and Products or Substances used in Animal Feed (FEEDAP); Scientific Opinion on the safety and efficacy of copper compounds (E4) as feed additives for all species: cupric chelate of amino acids hydrate, based on a dossier submitted by Zinpro Animal Nutrition Inc. *EFSA J.* 2013;11:3107. Available at: <http://www.efsa.europa.eu/en/efsajournal/pub/3107> (last accessed 1 December 2015).
- 271.** Gerber P, Opio C, Steinfeld H. Poultry production and the environment – a review. In: Thieme O, Pilling D, editors. *Proceedings of the International Conference Poultry in the Twenty-First Century: Avian Influenza and Beyond.* 2008. p. 379–405.
- 272.** Doyle MP, Busta F, Cords BR, Davidson PM, Hawker J, Hurd HS, *et al.* Antimicrobial Resistance: Implications for the Food System. *Compr Rev Food Sci F.* 2006;5(3):71-137.

- 273.** European Food Safety Authority (EFSA); Mercury as undesirable substance in animal feed Scientific opinion of the Panel on Contaminants in the Food Chain. EFSA J. 2008;654:1-76. Available at: <http://www.efsa.europa.eu/en/efsajournal/pub/654> (last accessed 1 December 2015).
- 274.** Bolan NS, Khan MA, Donaldson J, Adriano DC, Matthew C. Distribution and bioavailability of copper in farm effluent. Sci Total Environ. 2003;309(1-3):225-36.
- 275.** Xiong X, Yanxia L, Wei L, Chunye L, Wei H, Ming Y. Copper content in animal manures and potential risk of soil copper pollution with animal manure use in agriculture. Resour Conserv Recy. 2010;54(11):985-90.
- 276.** Zhang F, Li Y, Yang M, Li W. Content of Heavy Metals in Animal Feeds and Manures from Farms of Different Scales in Northeast China. Int J Environ Res Public Health. 2012;9(8):2658.
- 277.** Holzel CS, Muller C, Harms KS, Mikolajewski S, Schafer S, Schwaiger K, *et al.* Heavy metals in liquid pig manure in light of bacterial antimicrobial resistance. Environ Res. 2012;113:21-7.
- 278.** Ji X, Shen Q, Liu F, Ma J, Xu G, Wang Y, *et al.* Antibiotic resistance gene abundances associated with antibiotics and heavy metals in animal manures and agricultural soils adjacent to feedlots in Shanghai; China. J Hazard Mater. 2012;235-236:178-85.
- 279.** Zhu YG, Johnson TA, Su JQ, Qiao M, Guo GX, Stedtfeld RD, *et al.* Diverse and abundant antibiotic resistance genes in Chinese swine farms. Proc Natl Acad Sci U S A. 2013;110(9):3435-40.
- 280.** European Food Safety Authority (EFSA) Committed to ensuring that Europe's food is safe; Chemicals in food 2015 - Overview of Data Collection Reports. 2015. Available at: http://www.efsa.europa.eu/sites/default/files/corporate_publications/files/chemfood15.pdf (last accessed 1 December 2015).
- 281.** Zannoni D, Borsetti F, Harrison JJ, Turner RJ. The bacterial response to the chalcogen metalloids Se and Te. Adv Microb Physiol. 2008;53:1-72.
- 282.** European Commission Environment Directorate-General; Assessment of different options to address risks from the use phase of biocides. 2009. Available at: http://ec.europa.eu/environment/archives/ppps/pdf/final_reporto309.pdf (last accessed 1 December 2015).
- 283.** Pesticide Action Network (PAN) Germany; Biocide - treated consumer products: markets, policies and risks. 2013. Available at: http://www.pan-germany.org/download/biocides/biocide-treated_consumer_products.pdf (last accessed 1 December 2015).

- 284.** Gilbert P, McBain AJ. Potential impact of increased use of biocides in consumer products on prevalence of antibiotic resistance. *Clin Microbiol Rev.* 2003;16(2):189-208.
- 285.** Ortega Morente E, Fernandez-Fuentes MA, Grande Burgos MJ, Abriouel H, Perez Pulido R, Galvez A. Biocide tolerance in bacteria. *Int J Food Microbiol.* 2013;162(1):13-25.
- 286.** Hegstad K, Langsrud S, Lunestad BT, Scheie AA, Sunde M, Yazdankhah SP. Does the wide use of quaternary ammonium compounds enhance the selection and spread of antimicrobial resistance and thus threaten our health? *Microb Drug Resist.* 2010;16(2):91-104.
- 287.** Tezel U, Pavlostathis SG. Quaternary ammonium disinfectants: microbial adaptation, degradation and ecology. *Curr Opin Biotechnol.* 2015;33:296-304.
- 288.** Lemire JA, Harrison JJ, Turner RJ. Antimicrobial activity of metals: mechanisms, molecular targets and applications. *Nat Rev Microbiol.* 2013;11(6):371-84.
- 289.** Randall CP, Oyama LB, Bostock JM, Chopra I, O'Neill AJ. The silver cation (Ag⁺): antistaphylococcal activity, mode of action and resistance studies. *J Antimicrob Chemother.* 2013;68(1):131-8.
- 290.** Waldron KJ, Robinson NJ. How do bacterial cells ensure that metalloproteins get the correct metal? *Nat Rev Microbiol.* 2009;7(1):25-35.
- 291.** Dupont CL, Grass G, Rensing C. Copper toxicity and the origin of bacterial resistance-new insights and applications. *Metallomics.* 2011;3(11):1109-18.
- 292.** Ladomersky E, Petris MJ. Copper tolerance and virulence in bacteria. *Metallomics.* 2015;7(6):957-64.
- 293.** Chaturvedi KS, Henderson JP. Pathogenic adaptations to host-derived antibacterial copper. *Front Cell Infect Microbiol.* 2014;4:3.
- 294.** Osman D, Waldron KJ, Denton H, Taylor CM, Grant AJ, Mastroeni P, *et al.* Copper homeostasis in *Salmonella* is atypical and copper-CueP is a major periplasmic metal complex. *J Biol Chem.* 2010;285(33):25259-68.
- 295.** Hood MI, Skaar EP. Nutritional immunity: transition metals at the pathogen-host interface. *Nat Rev Microbiol.* 2012;10(8):525-37.
- 296.** Macomber L, Imlay JA. The iron-sulfur clusters of dehydratases are primary intracellular targets of copper toxicity. *Proc Natl Acad Sci U S A.* 2009;106(20):8344-9.
- 297.** Cerf O, Carpentier B, Sanders P. Tests for determining in-use concentrations of antibiotics and disinfectants are based on entirely different concepts: "resistance" has different meanings. *Int J Food Microbiol.* 2010;136(3):247-54.
- 298.** Russell AD. Bacterial adaptation and resistance to antiseptics, disinfectants and preservatives is not a new phenomenon. *J Hosp Infect.* 2004;57(2):97-104.
- 299.** Maillard JY, Bloomfield S, Coelho JR, Collier P, Cookson B, Fanning S, *et al.* Does microbicide use in consumer products promote antimicrobial resistance? A critical review

and recommendations for a cohesive approach to risk assessment. *Microb Drug Resist.* 2013;19(5):344-54.

300. Morrissey I, Oggioni MR, Knight D, Curiao T, Coque T, Kalkanci A, *et al.* Evaluation of epidemiological cut-off values indicates that biocide resistant subpopulations are uncommon in natural isolates of clinically-relevant microorganisms. *PLoS One.* 2014;9(1):e86669.

301. Mourão J, Rae J, Silveira E, Freitas AR, Coque TM, Peixe L, *et al.* Relevance of *tcrYAZB* operon acquisition for *Enterococcus* survival at high copper concentrations under anaerobic conditions. *J Antimicrob Chemother.* 2016;71(2):560-3.

302. Harrison JJ, Ceri H, Turner RJ. Multimetal resistance and tolerance in microbial biofilms. *Nat Rev Microbiol.* 2007;5(12):928-38.

303. Pal C, Bengtsson-Palme J, Rensing C, Kristiansson E, Larsson DG. BacMet: antibacterial biocide and metal resistance genes database. *Nucleic Acids Res.* 2014;42(Database issue):D737-43.

304. Hao X, Luthje FL, Qin Y, McDevitt SF, Lutay N, Hobman JL, *et al.* Survival in amoeba-a major selection pressure on the presence of bacterial copper and zinc resistance determinants? Identification of a "copper pathogenicity island". *Appl Microbiol Biotechnol.* 2015;99(14):5817-24.

305. Osman D, Patterson CJ, Bailey K, Fisher K, Robinson NJ, Rigby SE, *et al.* The copper supply pathway to a *Salmonella* Cu,Zn-superoxide dismutase (SodCII) involves P(1B)-type ATPase copper efflux and periplasmic CueP. *Mol Microbiol.* 2013;87(3):466-77.

306. Espariz M, Checa SK, Audero ME, Pontel LB, Soncini FC. Dissecting the *Salmonella* response to copper. *Microbiology.* 2007;153(Pt 9):2989-97.

307. Samanovic MI, Ding C, Thiele DJ, Darwin KH. Copper in microbial pathogenesis: meddling with the metal. *Cell Host Microbe.* 2012;11(2):106-15.

308. Achard ME, Tree JJ, Holden JA, Simpfordorfer KR, Wijburg OL, Strugnell RA, *et al.* The multi-copper-ion oxidase CueO of *Salmonella enterica* serovar Typhimurium is required for systemic virulence. *Infect Immun.* 2010;78(5):2312-9.

309. Pontel LB, Soncini FC. Alternative periplasmic copper-resistance mechanisms in Gram negative bacteria. *Mol Microbiol.* 2009;73(2):212-25.

310. Nies DH, Herzberg M. A fresh view of the cell biology of copper in enterobacteria. *Mol Microbiol.* 2013;87(3):447-54.

311. Checa SK, Espariz M, Audero ME, Botta PE, Spinelli SV, Soncini FC. Bacterial sensing of and resistance to gold salts. *Mol Microbiol.* 2007;63(5):1307-18.

- 312.** Pontel LB, Audero ME, Espariz M, Checa SK, Soncini FC. GolS controls the response to gold by the hierarchical induction of *Salmonella*-specific genes that include a CBA efflux-coding operon. *Mol Microbiol.* 2007;66(3):814-25.
- 313.** Pontel LB, Scampoli NL, Porwollik S, Checa SK, McClelland M, Soncini FC. Identification of a *Salmonella* ancillary copper detoxification mechanism by a comparative analysis of the genome-wide transcriptional response to copper and zinc excess. *Microbiology.* 2014;160(Pt 8):1659-69.
- 314.** Stevenson J, Barwinska-Sendra A, Tarrant E, Waldron KJ. Mechanism of action and applications of the antimicrobial properties of copper. In: Méndez-Vilas A, editor. *Microbial pathogens and strategies for combating them: science, technology and education.* 2013;468-79.
- 315.** Tetaz TJ, Luke RK. Plasmid-controlled resistance to copper in *Escherichia coli*. *J Bacteriol.* 1983;154(3):1263-8.
- 316.** Williams JR, Morgan AG, Rouch DA, Brown NL, Lee BT. Copper-resistant enteric bacteria from United Kingdom and Australian piggeries. *Appl Environ Microbiol.* 1993;59(8):2531-7.
- 317.** Zimmermann M, Udagedara SR, Sze CM, Ryan TM, Howlett GJ, Xiao Z, *et al.* PcoE-a metal sponge expressed to the periplasm of copper resistance *Escherichia coli*. Implication of its function role in copper resistance. *J Inorg Biochem.* 2012;115:186-97.
- 318.** Medardus JJ, Molla BZ, Nicol M, Morrow WM, Rajala-Schultz PJ, Kazwala R, *et al.* In-feed use of heavy metal micronutrients in U.S. swine production systems and its role in persistence of multidrug-resistant salmonellae. *Appl Environ Microbiol.* 2014;80(7):2317-25.
- 319.** Aarestrup FM, Hasman H. Susceptibility of different bacterial species isolated from food animals to copper sulphate, zinc chloride and antimicrobial substances used for disinfection. *Vet Microbiol.* 2004;100(1-2):83-9.
- 320.** Moreno Switt AI, den Bakker HC, Cummings CA, Rodriguez-Rivera LD, Govoni G, Raneiri ML, *et al.* Identification and characterization of novel *Salmonella* mobile elements involved in the dissemination of genes linked to virulence and transmission. *PLoS One.* 2012;7(7):e41247.
- 321.** Han J, Lynne AM, David DE, Tang H, Xu J, Nayak R, *et al.* DNA sequence analysis of plasmids from multidrug resistant *Salmonella enterica* serotype Heidelberg isolates. *PLoS One.* 2012;7(12):e51160.
- 322.** Peters JE, Fricker AD, Kapili BJ, Petassi MT. Heteromeric transposase elements: generators of genomic islands across diverse bacteria. *Mol Microbiol.* 2014;93(6):1084-92.
- 323.** Gupta A, Matsui K, Lo JF, Silver S. Molecular basis for resistance to silver cations in *Salmonella*. *Nat Med.* 1999;5(2):183-8.

- 324.** Randall CP, Gupta A, Jackson N, Busse D, O'Neill AJ. Silver resistance in Gram-negative bacteria: a dissection of endogenous and exogenous mechanisms. *J Antimicrob Chemother.* 2015;70(4):1037-46.
- 325.** Silver S. Bacterial silver resistance: molecular biology and uses and misuses of silver compounds. *FEMS Microbiol Rev.* 2003;27(2-3):341-53.
- 326.** Mijnenonckx K, Leys N, Mahillon J, Silver S, Van Houdt R. Antimicrobial silver: uses, toxicity and potential for resistance. *Biometals.* 2013;26(4):609-21.
- 327.** Woods EJ, Cochrane CA, Percival SL. Prevalence of silver resistance genes in bacteria isolated from human and horse wounds. *Vet Microbiol.* 2009;138(3-4):325-9.
- 328.** Sutterlin S, Tano E, Bergsten A, Tallberg AB, Melhus A. Effects of silver-based wound dressings on the bacterial flora in chronic leg ulcers and its susceptibility in vitro to silver. *Acta Derm Venereol.* 2012;92(1):34-9.
- 329.** Kremer AN, Hoffmann H. Subtractive hybridization yields a silver resistance determinant unique to nosocomial pathogens in the *Enterobacter cloacae* complex. *J Clin Microbiol.* 2012;50(10):3249-57.
- 330.** Sutterlin S, Edquist P, Sandegren L, Adler M, Tangden T, Drobni M, *et al.* Silver resistance genes are overrepresented among *Escherichia coli* isolates with CTX-M production. *Appl Environ Microbiol.* 2014;80(22):6863-9.
- 331.** Johnson TJ, Siek KE, Johnson SJ, Nolan LK. DNA sequence and comparative genomics of pAPEC-O2-R, an avian pathogenic *Escherichia coli* transmissible R plasmid. *Antimicrob Agents Chemother.* 2005;49(11):4681-8.
- 332.** Gilmour MW, Thomson NR, Sanders M, Parkhill J, Taylor DE. The complete nucleotide sequence of the resistance plasmid R478: defining the backbone components of incompatibility group H conjugative plasmids through comparative genomics. *Plasmid.* 2004;52(3):182-202.
- 333.** Chen YT, Chang HY, Lai YC, Pan CC, Tsai SF, Peng HL. Sequencing and analysis of the large virulence plasmid pLVPK of *Klebsiella pneumoniae* CG43. *Gene.* 2004;337:189-98.
- 334.** Wu KM, Li LH, Yan JJ, Tsao N, Liao TL, Tsai HC, *et al.* Genome sequencing and comparative analysis of *Klebsiella pneumoniae* NTUH-K2044, a strain causing liver abscess and meningitis. *J Bacteriol.* 2009;191(14):4492-501.
- 335.** Davis IJ, Richards H, Mullany P. Isolation of silver- and antibiotic-resistant *Enterobacter cloacae* from teeth. *Oral Microbiol Immunol.* 2005;20(3):191-4.
- 336.** Nascimento AM, Chartone-Souza E. Operon mer: bacterial resistance to mercury and potential for bioremediation of contaminated environments. *Genet Mol Res.* 2003;2(1):92-101.

- 337.** European Food Safety Authority (EFSA) Panel on Contaminants in the Food Chain (CONTAM); Scientific Opinion on the risk for public health related to the presence of mercury and methylmercury in food. EFSA J. 2012;10:2985. Available at: <http://www.efsa.europa.eu/en/efsajournal/pub/2985> (last accessed 1 December 2015).
- 338.** Mathema VB, Thakuri BC, Sillanpaa M. Bacterial mer operon-mediated detoxification of mercurial compounds: a short review. Arch Microbiol. 2011;193(12):837-44.
- 339.** Boyd ES, Barkay T. The mercury resistance operon: from an origin in a geothermal environment to an efficient detoxification machine. Front Microbiol. 2012;3:349.
- 340.** Liebert CA, Wireman J, Smith T, Summers AO. Phylogeny of mercury resistance (mer) operons of gram-negative bacteria isolated from the fecal flora of primates. Appl Environ Microbiol. 1997;63(3):1066-76.
- 341.** Levings RS, Partridge SR, Djordjevic SP, Hall RM. SGI1-K, a variant of the SGI1 genomic island carrying a mercury resistance region, in *Salmonella enterica* serovar Kentucky. Antimicrob Agents Chemother. 2007;51(1):317-23.
- 342.** Doublet B, Praud K, Weill FX, Cloeckert A. Association of IS26-composite transposons and complex In4-type integrons generates novel multidrug resistance loci in *Salmonella* genomic island 1. J Antimicrob Chemother. 2009;63(2):282-9.
- 343.** Crossman L. Large scale expansion of mobile elements in specific hotspot regions of the German outbreak *Escherichia coli* O104:H4. Available from Nature Precedings <<http://hdl.handle.net/10101/npre.2011.6466.1>> 2011.
- 344.** Grad YH, Godfrey P, Cerquiera GC, Mariani-Kurkdjian P, Gouali M, Bingen E, *et al.* Comparative genomics of recent Shiga toxin-producing *Escherichia coli* O104:H4: short-term evolution of an emerging pathogen. MBio. 2013;4(1):e00452-12.
- 345.** Welch TJ, Fricke WF, McDermott PF, White DG, Rosso ML, Rasko DA, *et al.* Multiple antimicrobial resistance in plague: an emerging public health risk. PLoS One. 2007;2(3):e309.
- 346.** Chen CY, Nace GW, Solow B, Fratamico P. Complete nucleotide sequences of 84.5- and 3.2-kb plasmids in the multi-antibiotic resistant *Salmonella enterica* serovar Typhimurium U302 strain G8430. Plasmid. 2007;57(1):29-43.
- 347.** Cain AK, Liu X, Djordjevic SP, Hall RM. Transposons related to Tn1696 in IncHI2 plasmids in multiply antibiotic resistant *Salmonella enterica* serovar Typhimurium from Australian animals. Microb Drug Resist. 2010;16(3):197-202.
- 348.** Garcia-Fernandez A, Carattoli A. Plasmid double locus sequence typing for IncHI2 plasmids, a subtyping scheme for the characterization of IncHI2 plasmids carrying extended-spectrum β -lactamase and quinolone resistance genes. J Antimicrob Chemother. 2010;65(6):1155-61.

- 349.** Lindsey RL, Frye JG, Fedorka-Cray PJ, Meinersmann RJ. Microarray-based analysis of IncA/C plasmid-associated genes from multidrug-resistant *Salmonella enterica*. *Appl Environ Microbiol.* 2011;77(19):6991-9.
- 350.** Wiesner M, Calva E, Fernandez-Mora M, Cevallos MA, Campos F, Zaidi MB, *et al.* *Salmonella* Typhimurium ST213 is associated with two types of IncA/C plasmids carrying multiple resistance determinants. *BMC Microbiol.* 2011;11(1):9.
- 351.** Reva O, Bezuidt O. Distribution of horizontally transferred heavy metal resistance operons in recent outbreak bacteria. *Mob Genet Elements.* 2012;2(2):96-100.
- 352.** Cain AK, Hall RM. Evolution of IncHI2 plasmids via acquisition of transposons carrying antibiotic resistance determinants. *J Antimicrob Chemother.* 2012;67(5):1121-7.
- 353.** Kariuki S, Okoro C, Kiiru J, Njoroge S, Omuse G, Langridge G, *et al.* Ceftriaxone-resistant *Salmonella enterica* serotype Typhimurium sequence type 313 from Kenyan patients is associated with the *bla*_{CTX-M-15} gene on a novel IncHI2 plasmid. *Antimicrob Agents Chemother.* 2015;59(6):3133-9.
- 354.** Nachman KE, Baron PA, Raber G, Francesconi KA, Navas-Acien A, Love DC. Roxarsone, inorganic arsenic, and other arsenic species in chicken: a U.S.-based market basket sample. *Environ Health Perspect.* 2013;121(7):818-24.
- 355.** Liu X, Zhang W, Hu Y, Hu E, Xie X, Wang L, *et al.* Arsenic pollution of agricultural soils by concentrated animal feeding operations (CAFOs). *Chemosphere.* 2015;119:273-81.
- 356.** Kaur S, Kamli MR, Ali A. Role of arsenic and its resistance in nature. *Can J Microbiol.* 2011;57(10):769-74.
- 357.** Chen J, Rosen BP. Biosensors for inorganic and organic arsenicals. *Biosensors (Basel).* 2014;4(4):494-512.
- 358.** Chen L, Chavda KD, Al Laham N, Melano RG, Jacobs MR, Bonomo RA, *et al.* Complete nucleotide sequence of a *bla*_{KPC}-harboring IncI2 plasmid and its dissemination in New Jersey and New York hospitals. *Antimicrob Agents Chemother.* 2013;57(10):5019-25.
- 359.** Shin J, Soo Ko K. Single origin of three plasmids bearing *bla*_{CTX-M-15} from different *Klebsiella pneumoniae* clones. *J Antimicrob Chemother.* 2014;69(4):969-72.
- 360.** Joerger RD, Hanning IB, Ricke SC. Presence of arsenic resistance in *Salmonella enterica* serovar Kentucky and other serovars isolated from poultry. *Avian Dis.* 2010;54(4):1178-82.
- 361.** Sampei G, Furuya N, Tachibana K, Saitou Y, Suzuki T, Mizobuchi K, *et al.* Complete genome sequence of the incompatibility group I1 plasmid R64. *Plasmid.* 2010;64(2):92-103.

- 362.** Chasteen TG, Fuentes DE, Tantalean JC, Vasquez CC. Tellurite: history, oxidative stress, and molecular mechanisms of resistance. *FEMS Microbiol Rev.* 2009;33(4):820-32.
- 363.** Turner RJ. Bacterial Tellurite Resistance. In: Kretsinger RH, Uversky VN, Permyakov EA, editors. *Encyclopedia of Metalloproteins*. Volume 1: Springer-Verlag New York; 2013. p. 219-22.
- 364.** Taylor DE. Bacterial tellurite resistance. *Trends Microbiol.* 1999;7(3):111-5.
- 365.** Orth D, Grif K, Dierich MP, Wurzner R. Variability in tellurite resistance and the *ter* gene cluster among Shiga toxin-producing *Escherichia coli* isolated from humans, animals and food. *Res Microbiol.* 2007;158(2):105-11.
- 366.** Bielaszewska M, Middendorf B, Tarr PI, Zhang W, Prager R, Aldick T, *et al.* Chromosomal instability in enterohaemorrhagic *Escherichia coli* O157:H7: impact on adherence, tellurite resistance and colony phenotype. *Mol Microbiol.* 2011;79(4):1024-44.
- 367.** Passet V, Brisse S. Association of tellurite resistance with hypervirulent clonal groups of *Klebsiella pneumoniae*. *J Clin Microbiol.* 2015;53(4):1380-2.
- 368.** Scientific Committee on Emerging and Newly Identified Health Risks (SCENIHR). Opinion on Nanosilver: safety, health and environmental effects and role in antimicrobial resistance. 2014. Available at: http://ec.europa.eu/health/scientific_committees/emerging/docs/scenih_r_o_039.pdf (last accessed 1 December 2015).
- 369.** Gaze WH, Krone SM, Larsson DG, Li XZ, Robinson JA, Simonet P, *et al.* Influence of humans on evolution and mobilization of environmental antibiotic resistome. *Emerg Infect Dis.* 2013;19(7).
- 370.** Berg J, Thorsen MK, Holm PE, Jensen J, Nybroe O, Brandt KK. Cu exposure under field conditions coselects for antibiotic resistance as determined by a novel cultivation-independent bacterial community tolerance assay. *Environ Sci Technol.* 2010;44(22):8724-8.
- 371.** Knapp CW, McCluskey SM, Singh BK, Campbell CD, Hudson G, Graham DW. Antibiotic resistance gene abundances correlate with metal and geochemical conditions in archived Scottish soils. *PLoS One.* 2011;6(11):e27300.
- 372.** Karatzas KA, Webber MA, Jorgensen F, Woodward MJ, Piddock LJ, Humphrey TJ. Prolonged treatment of *Salmonella enterica* serovar Typhimurium with commercial disinfectants selects for multiple antibiotic resistance, increased efflux and reduced invasiveness. *J Antimicrob Chemother.* 2007;60(5):947-55.
- 373.** Randall LP, Cooles SW, Coldham NG, Penuela EG, Mott AC, Woodward MJ, *et al.* Commonly used farm disinfectants can select for mutant *Salmonella enterica* serovar

Typhimurium with decreased susceptibility to biocides and antibiotics without compromising virulence. *J Antimicrob Chemother.* 2007;60(6):1273-80.

374. Karatzas KA, Randall LP, Webber M, Piddock LJ, Humphrey TJ, Woodward MJ, *et al.* Phenotypic and proteomic characterization of multiply antibiotic-resistant variants of *Salmonella enterica* serovar Typhimurium selected following exposure to disinfectants. *Appl Environ Microbiol.* 2008;74(5):1508-16.

375. Whitehead RN, Overton TW, Kemp CL, Webber MA. Exposure of *Salmonella enterica* serovar Typhimurium to high level biocide challenge can select multidrug resistant mutants in a single step. *PLoS One.* 2011;6(7):e22833.

376. Guo W, Cui S, Xu X, Wang H. Resistant mechanism study of benzalkonium chloride selected *Salmonella* Typhimurium mutants. *Microb Drug Resist.* 2014;20(1):11-6.

377. Webber MA, Whitehead RN, Mount M, Loman NJ, Pallen MJ, Piddock LJ. Parallel evolutionary pathways to antibiotic resistance selected by biocide exposure. *J Antimicrob Chemother.* 2015;70(8):2241-8.

378. Hansen LH, Jensen LB, Sorensen HI, Sorensen SJ. Substrate specificity of the OqxAB multidrug resistance pump in *Escherichia coli* and selected enteric bacteria. *J Antimicrob Chemother.* 2007;60(1):145-7.

379. Pal C, Bengtsson-Palme J, Kristiansson E, Larsson DG. Co-occurrence of resistance genes to antibiotics, biocides and metals reveals novel insights into their co-selection potential. *BMC Genomics.* 2015;16(1):964.

380. Antunes P, Machado J, Peixe L. Dissemination of *sul3*-containing elements linked to class 1 integrons with an unusual 3' conserved sequence region among *Salmonella* isolates. *Antimicrob Agents Chemother.* 2007;51(4):1545-8.

381. Khemtong S, Chuanchuen R. Class 1 integrons and *Salmonella* genomic island 1 among *Salmonella enterica* isolated from poultry and swine. *Microb Drug Resist.* 2008;14(1):65-70.


382. Domingues S, da Silva GJ, Nielsen KM. Global dissemination patterns of common gene cassette arrays in class 1 integrons. *Microbiology.* 2015;161(7):1313-37.

383. Gaze WH, Zhang L, Abdouslam NA, Hawkey PM, Calvo-Bado L, Royle J, *et al.* Impacts of anthropogenic activity on the ecology of class 1 integrons and integron-associated genes in the environment. *ISME J.* 2011;5(8):1253-61.

384. Wilson NL, Hall RM. Unusual class 1 integron configuration found in *Salmonella* genomic island 2 from *Salmonella enterica* serovar Emek. *Antimicrob Agents Chemother.* 2010;54(1):513-6.

385. Gullberg E, Albrecht LM, Karlsson C, Sandegren L, Andersson DI. Selection of a multidrug resistance plasmid by sublethal levels of antibiotics and heavy metals. *MBio.* 2014;5(5):e01918-14.

- 386.** Cavaco LM, Hasman H, Stegger M, Andersen PS, Skov R, Fluit AC, *et al.* Cloning and occurrence of *czrC*, a gene conferring cadmium and zinc resistance in methicillin-resistant *Staphylococcus aureus* CC398 isolates. *Antimicrob Agents Chemother.* 2010;54(9):3605-8.
- 387.** Cavaco LM, Hasman H, Aarestrup FM. Zinc resistance of *Staphylococcus aureus* of animal origin is strongly associated with methicillin resistance. *Vet Microbiol.* 2011;150(3-4):344-8.
- 388.** Aarestrup FM, Cavaco L, Hasman H. Decreased susceptibility to zinc chloride is associated with methicillin resistant *Staphylococcus aureus* CC398 in Danish swine. *Vet Microbiol.* 2010;142(3-4):455-7.



Chapter | 2

Objectives and outline of the study

"Imagination is more important than knowledge."

Albert Einstein

2.1. Statement of objectives

Non-typhoidal *Salmonella enterica* constitutes a worldwide major foodborne zoonotic pathogen. Its main reservoir is the intestinal tract of food-producing animals, which readily leads to contamination of farm surrounding environments and the food chain.

In recent years, expansion of particularly successful *Salmonella* serotypes and clones have been reported globally. They are transmitted to humans mostly by a wide range of foodstuffs of animal origin, they are involved in endemic/epidemic situations and they frequently show resistance to multiple antibiotics limiting therapeutic options when necessary. Interventions related to the limitation of antibiotics use (e.g. growth promoters ban in EU) were unsuccessful to control antibiotic resistance expansion, revealing the need for new approaches. In fact, recent priority programs included the study of the factors promoting antibiotic resistance in animal-environment-human interface (http://ec.europa.eu/health/scientific_committees/emerging/docs/scenihhr_o_028.pdf; <http://www.onehealthinitiative.com/about.php>), and stressed the need to overcome several knowledge gaps. Among them are included the effects of non-antibiotic compounds with antimicrobial activity, such as metals/biocides (incorporated in disinfectants, antiseptics, preservatives and as feed additives) in the selection and evolution of antibiotic resistant bacteria, namely in food-animal production environments.

The currently available information concerning the emergence and persistence of successful MDR *Salmonella* serotypes/clones in different environments and hosts is mostly related to the acquisition and maintenance of genetic elements harbouring antibiotic resistance genes. Indeed, most of these studies have been focused in the selection pressure created by the massive use of antibiotics, other than the human applications widely present in different food-animal production settings. More recently, diverse acquired genes coding for metals/biocides tolerance mechanisms have been described in scarce studies of *Salmonella* and other Gram-negative bacteria, but they only covered a few isolates/serotypes, sources and did not used a phenotypic/genotypic multilayered approach. As so, a comprehensive analysis of metals/biocides tolerance, including their acquired gene dispersion and genetic contexts within *Salmonella* from diverse ecological niches or belonging to successful or emergent MDR clones, remains poorly explored.

Our **hypothesis** was that tolerance to metals/biocides contributes for the survival and emergence/persistence of particular multidrug-resistant (MDR) *Salmonella* serotypes and clones in diverse environments and hosts.

The general **goal** of this work was to assess the occurrence of acquired metals/biocides tolerance genes as well as to characterize their associated genetic platforms and tolerance phenotypes in different *Salmonella* serotypes and emergent clones.

To accomplish this purpose, 406 isolates from a collection of ~2500 Portuguese non-typhoidal *Salmonella enterica*, representative of different serotypes and clones (years 2000 to 2014) were included. They were selected based on the source (human clinical cases, food products, food-animal production setting and the aquatic environment), geographical regions (North, Centre, South, Islands) and antibiotic susceptibility phenotypes and genotypes, Pulsed-Field Gel Electrophoresis (PFGE) types and/or Sequence Types (ST; obtained by Multilocus Sequence Typing-MLST).

This study focused on the following **specific aims**:

1. *To analyse the occurrence and dispersion of genes associated with tolerance to copper, silver, mercury, arsenic and tellurite.* We intended to determine whether *Salmonella* isolates from different ecological communities and hosts, are important reservoirs of metals/biocides tolerance genes.
2. *To understand the epidemiological and genetic backgrounds of Salmonella isolates carrying metals/biocides tolerance genes.* We intended to verify if the acquisition of metals/biocides tolerance was more associated with frequent or emergent MDR *Salmonella* serotypes and/or clones involved in human infections and if this feature could be an additional advantage for their maintenance in different ecological niches and hosts.
3. *To assess the susceptibility of diverse Salmonella serotypes/clones with different genetic backgrounds to copper and silver, two commonly used metals in food-animal setting.* We expected to unveil the advantage conferred by the acquisition of copper/silver tolerance genes to overcome metal toxic concentrations occurring in different food-animal settings, which might facilitate particular serotypes/clones survival and persistence in these environments. The phenotypic assays under

anaerobic atmospheres besides the aerobic classical ones intended to mimic oxygen conditions in diverse copper contaminated environments and hosts (e.g. manure, animal gut, waste lagoons) and to evaluate if anaerobic environments might have an important role in the selection of metal tolerant *Salmonella*.

4. *To characterize the genetic platforms involved in the acquisition of metal/biocide tolerance and to assess their transfer ability along with antibiotic resistance.* We intended to assess if metals/biocides tolerance genes were co-located in the same genetic platforms as antibiotic resistance genes, able to be co-selected by different compounds present in the food-animal setting. A detailed analysis of *Salmonella* available genomes was also performed in order to evaluate if predominant genetic platforms associated with several metals/biocides tolerance genes are spread in different *Salmonella* serotypes/clones recovered from different sources.

2.2. Outline of the thesis

The findings that would answer the *specific aims* of the thesis yielded different publications (n=6; 5 publications in peer review journals and 1 manuscript under final revision). Such results have been organized throughout the papers, according to the rationale proposed as follows:


I. Insights on the population structure and antimicrobial resistance of clinically-relevant *Salmonella* serotypes from clinical and non-clinical settings. In addition to isolates of food and human origin previously characterized in studies performed by our group, these works further characterized *Salmonella* serotypes with human clinical relevance recovered from pigs farms (e.g. *S. Rissen*, *S. Typhimurium*) and also belonging to the worldwide emergent serotype, *S. 4,[5],12:i:-*. The introduction of these pig-related serotypes was crucial since metals, such as copper and silver, are currently widely used in pig-production as animal feed for growth promotion and as disinfectants/antiseptics, respectively.

- a. Antunes P, **Mourão J**, Pestana N, Peixe L. 2011. Leakage of emerging clinically relevant multidrug-resistant *Salmonella* clones from pig farms. *Journal of Antimicrobial Chemotherapy*. 66(9):2028-2032.
- b. **Mourão J**, Machado J, Novais C, Antunes P, Peixe L. 2014. Characterization of the emerging clinically-relevant multidrug-resistant *Salmonella enterica* serotype

4,[5],12:i:- (monophasic variant of *S. Typhimurium*) clones. *European Journal of Clinical Microbiology & Infectious Diseases*. 33(12):2249-57.

II. Analysis of the dispersion of metals/biocides tolerance genes and tolerance phenotype as well as their associated genetic platforms in diverse non-typhoidal *Salmonella* serotypes and clones.

- c. **Mourão J**, Novais C, Machado J, Peixe L, Antunes P. 2015. Metal tolerance in emerging clinically relevant multidrug-resistant *Salmonella enterica* serotype 4,[5],12:i:- clones circulating in Europe. *International Journal of Antimicrobial Agents*. 45(6):610-6.
- d. **Mourão J**, Marçal S, Ramos P, Machado J, Peixe L, Novais C, Antunes P. 2015. Tolerance to multiple metal stressors in emerging non-typhoidal MDR *Salmonella* serotypes: a relevant role for copper in anaerobic conditions. *Submitted to the Journal of Antimicrobial Chemotherapy (under the second editorial revision)*.
- e. Antunes P, **Mourão J**, Alves T, Campos J, Novais C, Novais A, Peixe L. 2013. *Salmonella enterica* serotype Bovismorbificans, a new host for CTX-M-9. *International Journal of Antimicrobial Agents*. 41(1):91-93.
- f. Campos J, **Mourão J**, Marçal S, Machado J, Novais C, Peixe L, Antunes P. 2015. Clinical *Salmonella* Typhimurium ST34 with metal tolerance genes and an IncHI2 plasmid carrying *oqxAB-aac(6')-Ib-cr* from Europe. *Journal of Antimicrobial Chemotherapy*. DOI: <http://dx.doi.org/10.1093/jac/dkv409>.



Chapter | 3

Results and discussion

"You can never solve a problem on the same level on which it was created."

Albert Einstein

3.1. Insights on the population structure and antimicrobial resistance of clinically-relevant *Salmonella* serotypes from clinical and non-clinical settings

- 3.1.1. Leakage of emerging clinically relevant multidrug-resistant *Salmonella* clones from pig farms.
- 3.1.2. Characterization of the emerging clinically-relevant multidrug-resistant *Salmonella enterica* serotype 4,[5],12:i:- (monophasic variant of *S. Typhimurium*) clones.

**Leakage of emerging clinically relevant multidrug-resistant
Salmonella clones from pig farms**

Patrícia Antunes^{1, 2}, Joana Mourão², Nazaré Pestana² and Luísa Peixe^{1*}

¹ Faculdade de Ciências da Nutrição e Alimentação, Universidade do Porto, Rua Dr. Roberto Frias, 4200 Porto, Portugal; ² REQUIMTE, Laboratório de Microbiologia, Faculdade de Farmácia, Universidade do Porto, Rua Aníbal Cunha, no. 164, 4050-047 Porto, Portugal.

Journal of Antimicrobial Chemotherapy 2011 September; 66(9):2028-2032

Oxford University Press has authorized the reproduction of the final published PDF version of this paper in this thesis through the License Agreement Number 3786170804909 on January 12, 2016.

Leakage of emerging clinically relevant multidrug-resistant *Salmonella* clones from pig farms

Patrícia Antunes^{1,2}, Joana Mourão², Nazaré Pestana² and Luísa Peixe^{2*}

¹Faculdade de Ciências da Nutrição e Alimentação, Universidade do Porto, Rua Dr. Roberto Frias, 4200 Porto, Portugal; ²REQUIMTE, Laboratório de Microbiologia, Faculdade de Farmácia, Universidade do Porto, Rua Aníbal Cunha, no. 164, 4050-047 Porto, Portugal

*Corresponding author. Laboratório de Microbiologia, Faculdade de Farmácia, Universidade do Porto, Rua Aníbal Cunha, no. 164, 4050-047 Porto, Portugal. Tel: +351-22-2078972; Fax: +351-22-2003977; E-mail: lpeixe@ff.up.pt

Received 19 April 2011; returned 2 May 2011; revised 5 May 2011; accepted 9 May 2011

Objectives: To assess the presence of multidrug-resistant (MDR) *Salmonella* with human clinical relevance in pig farms from different regions of Portugal and to analyse their mobile genetic elements associated with antibiotic resistance.

Methods: Seventy-nine samples were collected from six piggeries and analysed for the presence of *Salmonella*. All isolates were examined for susceptibility to antimicrobial agents and representative isolates for resistance genes and class 1 integrons (PCR/restriction fragment length polymorphism). Clonality was determined by PFGE and multilocus sequence typing (MLST). Plasmid analysis included determination of size, content and characterization of the incompatibility group (rep-PCR and I-CeuI/S1-hybridization).

Results: Sixty *Salmonella* isolates were recovered from five samples (two manure, two waste lagoons and one animal feed) in half of the piggeries studied. All isolates were resistant to at least one antibiotic (tetracycline) and 97% to at least four antibiotics from different families. In 10 isolates representing different serogroup and resistance phenotype combinations a diversity of resistance genes and integrons was detected. These isolates belonged to the internationally widespread *Salmonella* Rissen (ST469) and *Salmonella* Typhimurium DT104 (ST19) clones, as well as to the emerging *Salmonella* Typhimurium monophasic variant with examples of Spanish (carrying a *sul3*-atypical integron within IncA/C plasmids, here assigned to ST19) and European (ASSuT phenotype, assigned to ST34) clones.

Conclusions: This is one of the few studies reporting emerging MDR *Salmonella* clones and the first one detecting *Salmonella* Typhimurium monophasic variant in the pig production setting. The survival of these strains in manure and waste lagoons is of concern, since these environments might allow spread of MDR bacteria beyond pig farms' boundaries.

Keywords: antimicrobial agents, piggeries, PFGE clones, MLST

Introduction

In recent years, changing trends in salmonellosis and associated serotypes have been observed, with a marked increase in certain multidrug-resistant (MDR) clones of *Salmonella* Typhimurium and its monophasic variant in different countries.^{1,2} Among these, *Salmonella* Typhimurium DT104, OXA-30-producing *Salmonella* Typhimurium clone and *Salmonella* Typhimurium monophasic variant (4,[5],12:i:-) are the most frequently reported in Europe.^{2,3} The latter, which have recently emerged worldwide, range from pan-susceptible (USA, Brazil) to MDR (Europe) strains and seem to be largely distributed in animal hosts and their derived products (e.g. pork, poultry products, cattle).² Although there is evidence that the animal setting seems to be a reservoir of MDR strains, data concerning the spread of emerging *Salmonella*

MDR clones with features of clinical interest within and off pig farms are still missing. Here, we assessed the presence of MDR *Salmonella* with human clinical relevance on pig farms from different regions of Portugal and analysed their mobile genetic elements associated with antibiotic resistance.

Materials and methods

Sample processing and *Salmonella* identification

Seventy-nine samples were collected within six geographically separated Portuguese piggeries (five with intensive and one with extensive production) during 2006 and 2007. They included samples from pigs ($n=21$; faeces, nostril/surface swabs), feed/medicines ($n=22$; feed, water, medicine, antiseptics), residues ($n=17$; swine waste lagoons,

residual waters, manure, septic tank) and piggery facilities ($n=19$; water, walls/floors dust, soil). The presence of *Salmonella* was screened by the conventional method following ISO 6579, which includes two stages of enrichment and plating out in two selective solid media (four plates per sample).⁴ Suspected colonies (up to five from each of the four plates) were identified by slide agglutination (*Salmonella* O poly antisera and serogroup-specific antisera for serogroups B, C1 and D; BD, USA), biochemical tests (API 32 GN; bioMérieux, Marcy l'Étoile, France) and a PCR assay (targeting the *invA* gene and a DT104/U302 phage type-specific DNA sequence).⁵ The serotypes of representative isolates were determined at the National Centre of *Salmonella*. The *Salmonella* Typhimurium monophasic variant (4,[5],12:i:-) isolates were confirmed using PCR as previously described.²

Antimicrobial susceptibility testing

All *Salmonella* isolates were tested for susceptibility to 10 antimicrobial agents (μg) [amoxicillin (10), gentamicin (10), kanamycin (30), streptomycin (10), ciprofloxacin (5), nalidixic acid (30), chloramphenicol (30), tetracycline (30), sulfamethoxazole (300) and trimethoprim (5)] by the disc diffusion method following CLSI standards.⁶ *Escherichia coli* ATCC 25922 was used as the control strain. Amoxicillin-resistant isolates were further tested for susceptibility to several extended-spectrum β -lactams (ceftazidime, ceftriaxone, cefotaxime, cefepime, ceftiofur, aztreonam and imipenem) and the double disc synergy test (DDST) for ESBL detection was also conducted.⁶ From each sample, the selection of representative isolates for further studies was based on serogroup, presence of the *invA* gene and/or phage type DT104/U302 phage type-specific DNA sequence and antibiotic resistance phenotype.

Characterization of antimicrobial resistance genes, integrons and plasmids

Genes coding for resistance to sulfamethoxazole (*sul1*, *sul2* and *sul3*), tetracycline [*tet(A)*, *tet(B)* and *tet(G)*], chloramphenicol (*floR*, *cmlA* and *catA*), amoxicillin (*bla_{TEM}*, *bla_{PSE-1}* and *bla_{OXA-30}*), gentamicin [*aac(3)-IV*], streptomycin (*aadA* and *strA-strB*) and trimethoprim (*dfrA1* and *dfrA12*) were searched for by PCR using primers and conditions previously described.^{7,8} The detection and characterization of class 1 integrons was performed by PCR and restriction fragment length polymorphism (RFLP) analysis with *TaqI* as previously reported.^{7,8} Positive and negative controls were included in all PCRs. Plasmid content and the genetic localization of integrons and *sul* genes were investigated by *S1* nuclease (Takara Bio Inc., Shiga, Japan) and *I-CeuI* (New England Biolabs, Ipswich, MA, USA) digestion of total genomic DNA followed by PFGE. Identification of plasmid incompatibility groups was determined by a rep-PCR typing method³ including three additional PCR assays for the IncU, IncR and ColE groups.⁹ Southern blot hybridization was performed by standard methods using *int1*, *sul1*, *sul2*, *sul3* and *rep* intragenic probes, following the manufacturer's instructions (Gene Images Alkphos Direct Labelling System Kit; Amersham GB/GE Healthcare Life Sciences UK Limited).

PFGE and MLST analysis

Clonal relatedness among isolates was assessed by PFGE following *XbaI* digestion of genomic DNA according to the standard 1 day protocol of the CDC.⁷ *Salmonella enterica* serotype Braenderup H9812 (CDC) was used as a molecular size marker. Multilocus sequence typing (MLST) analysis was performed using specific primers to amplify a set of seven housekeeping genes (*aroC*, *dhxN*, *hemD*, *hisD*, *purE*, *sucA* and *thrA*) and sequence type (ST) was assigned according to the MLST database (<http://mlst.ucc.ie/mlst>).

Results and discussion

Sixty *Salmonella* isolates were recovered from five positive samples (two manure, two waste lagoons and one animal feed) collected on three pig farms with intensive production (one located in the north and two located in the south of Portugal). Forty-seven isolates were serogroup B, of which 22 were positive for the DT104/U302 phage type-specific region, and 13 were serogroup C1. All isolates were resistant to at least one antibiotic (tetracycline) and 58 out of 60 were resistant to at least four antibiotics from different families (ranging from four to eight antibiotics). Resistances to tetracycline ($n=60$ isolates), streptomycin ($n=58$), sulfamethoxazole ($n=57$) and amoxicillin ($n=56$) were the most frequently detected and might reflect the high usage of these antibiotics in food-producing animals.⁷ Lesser rates of resistance were observed for the other antimicrobial agents tested; chloramphenicol ($n=22$), trimethoprim ($n=19$), nalidixic acid ($n=10$), gentamicin ($n=8$), ciprofloxacin ($n=0$) and kanamycin ($n=0$). Susceptibility to extended-spectrum β -lactams and absence of extended-spectrum β -lactamases (ESBLs) and AmpC were observed. Because enrichment steps allowed selection of more than one isolate of the same strain from the same sample, 10 isolates representing different combinations of serogroup and antibiotic resistance phenotype were selected for further studies (Table 1). The selected isolates belonged to three *Salmonella* serotypes (four Rissen, two Typhimurium and four from its monophasic variant 4,[5],12:i:-) and four PFGE clones spread among different farms and samples (Figure 1 and Table 1). Among them a diversity of antibiotic resistance genes were detected. These strains harboured common MDR genetic elements (e.g. integrons) and shared clonal relationships with previously nationally widespread/emerging clones in human and food products,^{7,8} showing their potential transmission along the food chain and their ability to cause human infections (Figure 1).

More than one resistant genotype or clone was detected within the same piggery, suggesting enrichment of the local metagenome with a broad range of drug-resistant strains and genetic elements, such as integrons. Isolates belonging to the emerging *Salmonella* Rissen clone ($n=4$), detected in the manure of two geographically separated piggeries, were assigned to ST469, only previously reported in the few European isolates of the same serotype allocated in the MLST database (<http://mlst.ucc.ie/mlst>). The human acquisition of *Salmonella* Rissen was previously associated with tourism or international trade outside the EU,¹⁰ but the data of this and a few other studies suggest a more local source of infection with this emerging serotype.⁷ Different resistance patterns and the presence of a specific class 1 integron (*dfrA12-orfF-aadA2*), located in IncR plasmids (35 kb), were observed, which may provide a selective advantage for this emergent serotype in the animal niche. Two isolates from the widely disseminated clone of *Salmonella* Typhimurium, DT104, showing the chromosomal located R-type ACSSuT (ampicillin, chloramphenicol, streptomycin, sulfamethoxazole, tetracycline), were detected in environmental samples from one piggery (manure and waste lagoon), suggesting that, beside the classical dissemination routes (animals and foods of animal origin), environmental reservoirs might also play a role in the spread of such strains. Of particular interest was the

Table 1. Characterization of the *Salmonella* clones from Portuguese piggeries

Serotype (phage type)/ PFGE type ^a	No. of isolates	MLST – ST (no. of isolates) ^b	Date of isolation	Piggery	Sample	Resistance phenotype ^c /resistance genes profile (no. of isolates; sample)	Class 1 integron genes (bp)	Class 1 integron and/or <i>sul2</i> location Chr or PL (kb, Inc) ^d
Rissen/N	4	ST469 (n=2)	2006	B, C	manure B1; manure C7	AMX, STR, SUL, TET, TMP <i>bla</i> _{TEM} , <i>aadA2</i> , <i>sul1</i> , <i>tet</i> (A), <i>dfrA12</i> (n=2; B1 and C7) AMX, STR, TET, TMP <i>bla</i> _{TEM} , <i>tet</i> (A) (n=1; B1) TET <i>tet</i> (A) (n=1; C7)	<i>intI1</i> , <i>sul1</i> <i>dfrA12</i> , <i>orfF</i> , <i>aadA2</i> (2000) <i>intI1</i>	PL (35, R) PL (30, R) —
Typhimurium (DT104/ U302)/A	2	ST19 (n=1)	2007	E	waste lagoon E32; manure E34	AMX, CHL, STR, SUL, TET <i>bla</i> _{PSE-1} , <i>floR</i> , <i>aadA2</i> , <i>sul1</i> , <i>tet</i> (G) (n=2; E32 and E34)	<i>intI1</i> , <i>sul1</i> <i>aadA2</i> (1000) <i>bla</i> _{PSE-1} (1200)	Chr
Typhimurium (DT104/ U302) 4,[5],12:i:-/O	2	ST19 (n=1)	2006	C	manure C7	AMX, CHL, GEN, NAL, STR, SUL, TET, TMP <i>bla</i> _{TEM} , <i>cmiA1</i> , <i>aac</i> (3)-IV, <i>aadA2</i> , <i>sul1</i> - <i>sul2</i> - <i>sul3</i> , <i>tet</i> (A), <i>dfrA12</i> (n=1; C7) CHL, NAL, STR, TET, SUL <i>cmiA1</i> , <i>aadA</i> , <i>tet</i> (A), <i>sul1</i> - <i>sul2</i> - <i>sul3</i> (n=1; C7)	<i>intI1</i> , <i>sul1</i> <i>dfrA12</i> , <i>orfF</i> , <i>aadA2</i> (2000) type III- <i>sul3</i> ^e	PL (170, A/C) PL (130, A/C)
Typhimurium 4,[5],12:i:-/Y	2	ST34 (n=2)	2007	E	feed E2; manure E34	AMX, STR, SUL, TET <i>bla</i> _{TEM} , <i>strA</i> - <i>strB</i> , <i>sul2</i> , <i>tet</i> (B) (n=2; E2 and E34)	—	Chr (<i>sul2</i>)

^aClones are designated by capital letters, as previously published.^{7,8}^bNumber of isolates submitted to MLST database.^cAMX, amoxicillin; CHL, chloramphenicol; GEN, gentamicin; NAL, nalidixic acid; STR, streptomycin; SUL, sulfamethoxazole; TET, tetracycline; TMP, trimethoprim.^dChromosomal (Chr) and/or plasmid (PL) location of integrons and the *sul2* gene was assessed by hybridization of 1-CeuI/S1-digested genomic DNA using *int1*, *sul1*, *sul2*, *sul3* and *rep* probes.^eStructure of the type III *sul3* integron: 5'CS-*estX*-*psp*-*aadA2*-*cmiA1*-*aadA1*-*qacH*-15440-*sul3*.⁸




Figure 1. PFGE patterns of *Salmonella* isolates from piggeries and others previously characterized.^{7,8} Lanes: M, *Salmonella* Braenderup H9812 (CDC); 1, clone O, human isolate (*Salmonella* Typhimurium 4,[5],12:i:-, 459/02); 2, clone O, pork isolate (*Salmonella* Typhimurium 4,[5],12:i:-, A30); 3, C7.5 isolate (*Salmonella* Typhimurium 4,[5],12:i:-, piggy C; clone O); 4, C7.17 isolate (*Salmonella* Typhimurium 4,[5],12:i:-, piggy C; clone O); 5, clone O, pork isolate (*Salmonella* Typhimurium 4,[5],12:i:-, B296); 6, C7.7 isolate (*Salmonella* Rissen, piggy C; clone N); 7, B1.1 isolate (*Salmonella* Rissen, piggy B; clone N); 8, clone N, pork isolate (*Salmonella* Rissen, A27); 9, C7.16 isolate (*Salmonella* Rissen, piggy C; clone N); 10, B1.3 isolate (*Salmonella* Rissen, piggy B; clone N); 11, E32.2 isolate (*Salmonella* Typhimurium DT104, piggy E; clone A); 12, E2.5 isolate (*Salmonella* Typhimurium 4,[5],12:i:-, piggy E; clone Y); 13, E34.1 isolate (*Salmonella* Typhimurium 4,[5],12:i:-, piggy E; clone Y); and 14, E34.11 isolate (*Salmonella* Typhimurium DT104, piggy E; clone A).

detection of two MDR clones of the emerging *Salmonella* Typhimurium monophasic variant, related to the so-called European and Spanish clones.^{11,12} The pattern ASSuT was observed in the isolates from one piggy associated with the chromosomal resistance genes [*bla*_{TEM-1}, *strA-strB*, *sul2* and *tet(B)*] typical of the European monophasic strains,¹² here demonstrated to belong to ST34 (a single-locus variant of ST19). The other monophasic variant MDR clone, belonging to the worldwide-distributed ST19 and DT104/U302 phage type, presented an IncA/C plasmid (130–170 kb) with the recently characterized atypical *sul3*-carrying integron type III.⁸ Interestingly, isolates of this monophasic variant clone carrying similar MDR features have been recovered for more than 10 years in Spain¹¹ and from humans and food products in Portugal, and were initially identified as *Salmonella* Typhimurium.⁸ Successful adaptation of this pathogen and the intense commercial trade in live pigs and pig meat products between these countries might explain the maintenance and dissemination of the monophasic MDR so-called Spanish clone in the Iberian Peninsula.

This study indicates that clinically relevant MDR clones are prevalent and disseminated in piggeries. The success of these bacterial strains might be related to antimicrobial resistance features, but also to other properties, such as virulence factors, as recently described for the IncA/C plasmids in the Spanish clone.¹¹ Also, the detection of these widely frequent MDR genetic lineages of *Salmonella* in manure and swine waste

lagoons raises concerns due to potential water, air and soil contamination, as well as of wild animals, indicating potentially high risks of transmission of the pathogen in the environment, animals and humans.^{13–15} The frequent use of manure as a fertilizer and the long-term survival potential of this zoonotic pathogen seem to contribute to the increasing reports of salmonellosis outbreaks caused by consumption of fresh products (e.g. ready-to-eat salads).^{13,14} This alerts us to the ability of the emergent clones, such as subtypes of *Salmonella* Typhimurium monophasic variant to be transmitted to humans by sources other than pigs and pork, until now considered sporadic.

In conclusion, this is one of the few studies reporting emerging MDR *Salmonella* clones carrying features with clinical interest and the first one identifying *Salmonella* Typhimurium monophasic variant in pig farms. The presence of isolates resistant to several antibiotic families and belonging to clonal lineages increasingly associated with human infections, particularly the emergence of *Salmonella* Typhimurium monophasic variants in piggeries, is a public health threat.

Acknowledgements

We are deeply grateful to Carla Novais for critical review of this paper prior to submission. We are also grateful to Alessandra Carattoli for kindly providing the positive controls for the IncU, IncR and ColE plasmids, to Centro Nacional de *Salmonella* (Lisboa, Portugal) for serotyping the strains and CDC for the PFGE protocols and the control strain *Salmonella* Braenderup H9812.

Funding

This work was partially supported by Fundação para a Ciência e a Tecnologia (FCT), which belongs to the Ministry of Science, Technology and Innovation from Portugal (POCI/AMB/61814/2004).

Transparency declarations

None to declare.

References

- 1 Parry CM, Threlfall EJ. Antimicrobial resistance in typhoidal and nontyphoidal salmonellae. *Curr Opin Infect Dis* 2008; **21**: 531–8.
- 2 European Food Safety Authority. EFSA Panel on Biological Hazards (BIOHAZ). Scientific opinion on monitoring and assessment of the public health risk of “*Salmonella* Typhimurium-like” strains. *EFSA J* 2010; **8**: 1826.
- 3 Antunes P, Coque T, Peixe L. Emergence of an IncI γ plasmid encoding CMY-2 β -lactamase associated with the international ST19 OXA-30-producing β -lactamase *Salmonella* Typhimurium multidrug-resistant clone. *J Antimicrob Chemother* 2010; **65**: 2097–100.
- 4 International Organization for Standardization. ISO 6579, *Microbiology of Food and Animal Feeding Stuffs - Horizontal Method for the Detection of Salmonella spp.* International Organization for Standardization, 2002.
- 5 Pritchett LC, Konkel ME, Gay JM *et al.* Identification of DT104 and U302 phage types among *Salmonella enterica* serotype Typhimurium isolates by PCR. *J Clin Microbiol* 2000; **38**: 3484–8.

Antunes et al.

- 6 Clinical and Laboratory Standards Institute. *Performance Standards for Antimicrobial Susceptibility Testing: Seventeenth Informational Supplement M100-S17*. CLSI, Wayne, PA, USA, 2007.
- 7 Antunes P, Machado J, Peixe L. Characterization of antimicrobial resistance and class 1 and 2 integrons in *Salmonella enterica* isolates from different sources in Portugal. *J Antimicrob Chemother* 2006; **58**: 297–304.
- 8 Antunes P, Machado J, Peixe L. Dissemination of *sul3*-containing elements linked to class 1 integrons with an unusual 3'CS region among *Salmonella* isolates. *Antimicrob Agents Chemother* 2007; **51**: 1545–8.
- 9 García-Fernández A, Fortini D, Veldman K et al. Characterization of plasmids harbouring *qnrS1*, *qnrB2* and *qnrB19* genes in *Salmonella*. *J Antimicrob Chemother* 2009; **63**: 274–81.
- 10 Hendriksen RS, Bangtrakulnonth A, Pulsrikarn C et al. Antimicrobial resistance and molecular epidemiology of *Salmonella* Rissen from animals, food products, and patients in Thailand and Denmark. *Foodborne Pathog Dis* 2008; **5**: 605–19.
- 11 García P, Guerra B, Bances M et al. IncA/C plasmids mediate antimicrobial resistance linked to virulence genes in the Spanish clone of the emerging *Salmonella enterica* serotype 4,[5],12:i:-. *J Antimicrob Chemother* 2011; **66**: 543–9.
- 12 Lucarelli C, Dionisi AM, Torpdahl M et al. Evidence for a second genomic island conferring multidrug resistance in a clonal group of strains of *Salmonella enterica* serovar Typhimurium and its monophasic variant circulating in Italy, Denmark, and the United Kingdom. *J Clin Microbiol* 2010; **48**: 2103–9.
- 13 Baloda SB, Christensen L, Trajcevska S. Persistence of a *Salmonella enterica* serovar Typhimurium DT12 clone in a piggery and in agricultural soil amended with *Salmonella*-contaminated slurry. *Appl Environ Microbiol* 2001; **67**: 2859–62.
- 14 United States Environmental Protection Agency. *Detecting and Mitigating the Environmental Impact of Fecal Pathogens Originating From Confined Animal Feeding Operations: Review*. EPA National Risk Management Research Laboratory, 2005.
- 15 Allen SE, Boerlin P, Janecko N et al. Antimicrobial resistance in generic *Escherichia coli* isolates from wild small mammals living in swine farm, residential, landfill, and natural environments in Southern Ontario, Canada. *Appl Environ Microbiol* 2011; **77**: 882–8.

Characterization of the emerging clinically-relevant multidrug-resistant *Salmonella enterica* serotype 4,[5],12:i:- (monophasic variant of *S. Typhimurium*) clones

Joana Mourão¹, Jorge Machado², Carla Novais¹, Patrícia Antunes^{1,3} and Luísa Peixe^{1*}

¹ REQUIMTE, Laboratório de Microbiologia, Faculdade de Farmácia, Universidade do Porto, Porto, Portugal; ² Laboratório Nacional de Referência de Infecções Gastrointestinais, Departamento de Doenças Infecciosas, Instituto Nacional de Saúde Dr. Ricardo Jorge, Lisboa, Portugal; ³ Faculdade de Ciências da Nutrição e Alimentação, Universidade do Porto, Porto, Portugal.

**European Journal of Clinical Microbiology & Infectious Diseases 2014
December; 33(12):2249-2257**

Springer has authorized the reproduction of the final published PDF version of this paper in this thesis through the License Agreement Number 3787890799997 on January 15, 2016.

ARTICLE

Characterization of the emerging clinically-relevant multidrug-resistant *Salmonella enterica* serotype 4,[5],12:i:- (monophasic variant of *S. Typhimurium*) clonesJ. Mourão · J. Machado · C. Novais · P. Antunes ·
L. PeixeReceived: 9 April 2014 / Accepted: 1 June 2014 / Published online: 15 July 2014
© Springer-Verlag Berlin Heidelberg 2014

Abstract To better understand the recent success/emergence of *Salmonella enterica* serotype 4,[5],12:i:- we characterized the population diversity, *fljAB* deletion patterns, antibiotic resistance features and associated genetic elements of a comprehensive collection obtained in the last decade from Portugal (2002–2010). One hundred thirty-one isolates from human clinical specimens, food, environment and piggeries, verified by PCR as *S. 4,[5],12:i:-*, were studied for clonality (Pulsed-Field Gel Electrophoresis and Multilocus Sequence Typing), antibiotic resistance by phenotypic (disk diffusion and/or agar dilution) and genotypic (PCR/Restriction Fragment Length Polymorphism and sequencing, genomic location) methods and *fljAB*-deletions (PCR). Plasmid analysis included determination of size, content and characterization of the incompatibility group (PCR-Based Replicon Typing and I-CeuI/S1-hybridization). Results showed three multidrug-resistant (MDR) clones circulating and causing infections, associated with particular phenotypic and genotypic features. Most of the isolates belonged to the widespread European (ASSuT phenotype, RR1-RR2 resistance regions, ST34) and Spanish (carrying a *sul3*-type III integron within

IncA/C plasmids, ST19) clones circulating in Europe. A third clone, here designated Southern European clone (carrying a *sul3*-type I integron within IncR plasmids, ST19), presents a *fljAB* region different from the previous clones and similar to the US strains, despite differences in the MDR mobile genetic platforms. The success of *S. 4,[5],12:i:-* might be related to the selective advantage offered by MDR profiles associated with stable genetic elements, also carrying virulence features, along with well adapted clones to the animal food production and causing human infections.

Introduction

Non-typhoidal *Salmonella* is one of the leading causes of zoonotic food borne diseases, presenting public health relevance, including such industrialized regions as the European Union (EU) [1, 2]. Currently, changing trends in salmonellosis and associated serotypes have been observed, with predominant multidrug resistant (MDR) *Salmonella* clonal lineages scattered worldwide in a wide range of food animals and involved in endemic/epidemic scenarios, such as described for *Salmonella enterica* serotype 4,[5],12:i:- [1, 3]. This serotype is a monophasic variant of the recognized pathogenic *Salmonella enterica* serotype Typhimurium and has become a new epidemic strain causing human infections transmitted by the food chain, suggesting good adaptability to different environmental conditions and colonization/infection of a wide range of animals and humans [1, 3–5].

In Europe, including Portugal, *S. 4,[5],12:i:-* has been the third most frequent serotype among human salmonellosis [1] and responsible for several recent large outbreaks [6, 7]. Food animals, particularly pigs, were pointed out as its reservoir and their derived products as the main infection vehicles [3, 8, 9]. Those *S. 4,[5],12:i:-* isolates present MDR profiles associated with particular widespread clonal lineages, designated

Electronic supplementary material The online version of this article (doi:10.1007/s10096-014-2180-1) contains supplementary material, which is available to authorized users.

J. Mourão · C. Novais · P. Antunes · L. Peixe
REQUIMTE, Laboratório de Microbiologia, Faculdade de Farmácia,
Universidade do Porto, Porto, Portugal

J. Machado
Laboratório Nacional de Referência de Infecções Gastrointestinais,
Departamento de Doenças Infecciosas, Instituto Nacional de Saúde
Dr. Ricardo Jorge, Lisboa, Portugal

P. Antunes (✉)
Faculdade de Ciências da Nutrição e Alimentação, Universidade do
Porto, Rua Dr. Roberto Frias, 4200-465 Porto, Portugal
e-mail: patriciaantunes@fcna.up.pt

European and Spanish clones [3, 4]. However, few studies comprehensively analyse the genetic elements (genes, integrons, plasmids) spreading antibiotic resistance [10–12] and the *fljAB* genomic deletion patterns in *S. 4,[5],12:i:-* [13]. This serotype is currently targeted by EU actions to detect and control *Salmonella* serotypes of concern [14], requiring further surveillance and mitigation strategies, including its epidemiological success characterization. To better understand the recent emergence of *S. 4,[5],12:i:-* we characterized the population diversity, antibiotic resistance features and associated genetic elements, plus *fljAB* deletion patterns, of a comprehensive collection obtained in the last decade from Portugal.

Materials and methods

Bacterial isolates

A total of 131 isolates of *S. 4,[5],12:i:-* were studied, with isolates recovered from human clinical specimens ($n=114$; faeces, blood, peritoneal fluid and urine; from 28 hospitals and two community laboratories), food of animal origin ($n=9$; pork, beef and poultry), environment ($n=4$; drinking and bathing water) and piggeries ($n=4$; manure and feed) between 2002 and 2010 from different geographical areas of Portugal. Those isolates were serologically identified as *S. 4,[5],12:i:-* by the National Reference Centre (INSA, Lisbon, Portugal) and confirmed by PCR targeting the intergenic region *fliB-fliA* and *fljB* gene coding for phase-2 flagellar antigen [3, 15]. For evolutionary analyses, all isolates were screened for genomic deletions patterns surrounding *fljAB* operon (*fljA*, *fljB*, *hin*, *iroB*, STM2740 and STM2757 genes and STM1053-1997 region) by PCR [13]. Also, isolates were screened by multiplex PCR targeting *invA* gene and DT104/U302 phage-type-specific DNA [16].

Molecular subtyping

Clonal relatedness among isolates was established by pulsed-field gel electrophoresis (PFGE) of *XbaI*-digested genomic DNA [8]. Band patterns were analysed with InfoQuest™ FP software v4.5 (BioRad Laboratories) and cluster analysis with unweighted pair group method with arithmetic averages based on Dice similarity coefficient with 1.0 % optimization and tolerance. Isolates with Dice band-based similarity coefficient ≥ 85 % were considered to belong to the same PFGE-type (designated by capital letters). Multilocus sequence typing (MLST) was also performed in representative isolates ($n=16$; different PFGE-types/clones) (<http://mlst.ucc.ie/mlst/dbs/Senterica>).

Study of antibiotic susceptibility and resistance genes

Susceptibility to amoxicillin, gentamicin, kanamycin, streptomycin, ciprofloxacin, nalidixic acid, chloramphenicol, tetracycline, sulfamethoxazole and trimethoprim was determined by disc diffusion or agar dilution methods (ciprofloxacin) following CLSI guidelines [17]. Amoxicillin resistant isolates were further tested for several extended spectrum β -lactams (ceftazidime, cefotaxime, cefepime and ceftiofur) and by double disk synergy test for ESBL [17]. MDR was considered when isolates were resistant to three or more antibiotics of different families. The search for genes encoding resistance to amoxicillin (*bla*_{TEM}), chloramphenicol (*catA*, *cmlA1* and *floR*), ciprofloxacin [*qnrA*, *qnrB*, *qnrC*, *qnrD*, *qnrS*, *qepA*, *oqxA*, *oqxB* and *aac(6)-Ib-cr*], gentamicin [*aac(3)-IV*], kanamycin (*aphA*) streptomycin (*strA-strB*, *aadA*), sulfamethoxazole (*sul1*, *sul2*, *sul3*), tetracycline [*tet(A)*, *tet(B)*, *tet(G)*] and trimethoprim (*dfrA1*, *dfrA12*) was performed by PCR [8]. Detection and characterization of class 1 integrons was performed by PCR and restriction fragment length polymorphism (*TaqI*) [18, 19]. Representative isolates ($n=22$; different PFGE-types) were also PCR tested for presence of left, right and internal regions of genomic resistance region RR1 (RR1LJ, RR1RJ, *tnpB* and *repC* regions/genes) and RR2 (RR2LJ, RR2RJ, *IS10* and *tetC(B)* Δ regions/genes) [11]. A positive amplicon from each PCR was sequenced and compared with known GenBank sequences to validate the method.

Characterization of the genetic elements carrying antibiotic resistance genes

Conjugation assays were done for isolates representative of different PFGE-types using *Escherichia coli* K802N (rifampicin and nalidixic acid resistant) as recipient strain [19]. Transconjugants were selected on Mueller-Hinton 2 agar containing nalidixic acid (64 mg/L) plus tetracycline (8 mg/L) or sulfamethoxazole (256 mg/L). Classification of plasmids was based on PCR-based replicon typing [20, 21] and location of antibiotic resistance genes by hybridization of S1-PFGE and I-CeuI-PFGE with specific probes [8, 19].

Results

Typing of *Salmonella 4,[5],12:i:-* isolates

The 131 isolates corresponded to 35 PFGE profiles, which were grouped into ten PFGE-types and three clones (showing 67.8 to 75.2 % identity) (Online Resource 1). Two showed phenotypic and genotypic features similar to previously described European and Spanish clones (see further sections). A third one presented unique features and was here designated

Southern European clone. Within the European clone were included 50 % of monophasic strains, non DT104/U302 phage-type. They were classified in three prevalent PFGE-types, with type-A gathering strains from humans, environment and piggeries (Table 1). The Spanish clone (DT104/U302 phage-type) included 35 % of the isolates, dominated by PFGE type O ($n=44$), including also human, environment, food and piggeries isolates. Finally, the less prevalent Southern European clone includes 15 % of the isolates of human and food origins, all non DT104/U302 phage-type (Table 1). The MLST typing assigned Spanish and Southern European clones in ST19 and the European clone in its single-locus variant ST34. Also, genomic deletions patterns surrounding the *fljAB* region were constant in all isolates from MDR Spanish and Southern European clones, but diverse in European clone isolates, in spite of 70 % ($n=46/66$) presenting a predominant profile (Table 1).

Antibiotic resistance study

All but two *S.* 4,[5],12:i:- isolates ($n=129/131$; 98 %) were resistant to at least one of the antibiotics tested. They expressed resistance to sulfamethoxazole ($n=121/92$ %), tetracycline ($n=119/91$ %), streptomycin ($n=119/91$ %), amoxicillin ($n=89/68$ %), chloramphenicol ($n=60/46$ %), trimethoprim ($n=46/35$ %), gentamicin ($n=35/27$ %) and kanamycin ($n=4/3$ %). Decreased susceptibility to ciprofloxacin ($n=6/5$ %), with ($n=4$) or without ($n=2$) resistance to nalidixic acid, was only observed in isolates from PFGE-types B (European clone) and O (Spanish clone), but none carried the plasmid-mediated quinolone resistance genes screened. Susceptibility to extended-spectrum β -lactams and absence of ESBL/AmpC were also observed.

Diverse antibiotic resistance profiles (ranging from one to eight antibiotics) were identified among isolates belonging to the three clones, with 92 % ($n=121/131$) resistant to three or more antibiotics. The most frequent profiles were ASSuT (amoxicillin, streptomycin, sulfamethoxazole, tetracycline) typical of the European clone ($n=51/66$), ACGSSuTTm (amoxicillin, chloramphenicol, gentamicin, streptomycin, sulfamethoxazole, tetracycline, trimethoprim) typical of the Spanish clone ($n=26/46$) and CSSuTTm (chloramphenicol, streptomycin, sulfamethoxazole, tetracycline, trimethoprim) on the Southern European clone ($n=15/19$) (Table 1).

Genetic elements carrying antibiotic resistance genes

Isolates of the European clone presented the typical antibiotic resistance genes [*bla*_{TEM-1}, *strA-strB*, *sul2* and *tet*(B)] chromosomally located and non-transferable. Absence of class 1 integrons in all but two isolates was also detected (Table 1). None of the tested isolates from this clone ($n=22$; all ASSuT) was simultaneously positive for presence of left, right and

internal genes/regions of chromosomal genomic regions RR1 (RR1LJ⁺ - 32 %, RR1RJ⁺ - 18 % regions and *tnpB*⁺ - 55 %, *repC*⁺ - 73 % genes) and RR2 (RR2LJ⁺ - 14 %, RR2RJ⁺ - 95 % regions and *IS10*⁺ - 73 %, *tetC*(B) Δ - 77 % genes). Although it was not possible to establish an association between those profiles and specific PFGE-types, six out of nine isolates from PFGE-type A exhibited particular patterns (RR1LJ + *tnpB* + *repC* + *tetC*(B) Δ + RR2RJ, $n=4$ and *tnpB* + *repC* + RR1RJ + RR2LJ + *IS10* + RR2RJ, $n=2$).

In contrast, the Spanish and Southern European clones, despite being included in the same ST19 lineage, carried different atypical class 1 integrons [*sul3*-type III (5'CS-*estX*-*psp*-*aadA2*-*cmlA1*-*aadA1*-*qacH*-*IS440*-*sul3*) on Spanish clone; *sul3*-type I (5'CS-*dfrA12*-*orfF*-*aadA2*-*cmlA1*-*aadA1*-*qacH*-*IS440*-*sul3*) on Southern European clone], located in the non-transferable plasmids, IncA/C (110–220 Kb) or IncR (80–160 Kb), respectively. Furthermore, most isolates belonging to the Spanish clone also carried one typical class 1 integron (*dfrA12*-*orfF*-*aadA2*-*qacE* Δ 1-*sul1*), *tet*(A), *aac*(3)-IV and *bla*_{TEM} within the same plasmid (Table 1). None of the Southern European clone isolates exhibited resistance to amoxicillin (*bla*_{TEM}) and the tetracycline resistance was associated with *tet*(B) located in IncR plasmids.

Discussion

A decreasing trend in human salmonellosis cases is observed in European countries, mainly due to successful control programs in poultry/eggs production [1]. Nevertheless, occurrence of MDR clonal lineages causing human infections and resident in food animal settings becomes increasingly frequent, with the emergent *S.* 4,[5],12:i:- suggested as a new epidemic serotype in Europe [3, 4, 8]. In Portugal, the number of reported cases associated with this monophasic variant of *S.* Typhimurium had a nine-fold increase during this study period and nowadays ranks third in frequency (1,1 % in 2002 versus 22.6 % in 2010) [22], similar to other European countries [1, 3].

This study reports high rates (92 %) of MDR in *S.* 4,[5],12:i:- isolates from different origins and geographical regions of Portugal. Resistance to sulfamethoxazole, tetracycline, streptomycin and amoxicillin were the most frequent and might reflect their high use in food-production animals in Europe [23, 24]. Most of *S.* 4,[5],12:i:- isolates studied were grouped in the European clone/ST34, corroborating its recent epidemic emergence in Europe [4, 9, 25, 26] and, more recently, also in America [5, 27], which in all cases presented similar phenotypic and genotypic ASSuT markers. Nevertheless, a considerable diversity in PFGE-types is currently disseminated in Europe, particularly STYMXB.0131, STYMXB.0079 and STYMXB.0010 (PulseNet Europe) [4, 9,

Table 1 Genotypic and phenotypic characterization of *Salmonella enterica* serotype 4,[5],12:i:- clones from Portugal

Clone designation Phage type ^a MLST-ST (eBG) (no. isolates) ^b Predominant <i>fljAB</i> deletion pattern ^c	PFGE-type (no. isolates/ no. PFGE subtypes) ^d	Source (no. isolates)	Year/ region ^e	Resistance phenotype (no. isolates) ^f / Resistance genes profile	Class 1 integron (bp) ⁱ (no. isolates)	Class 1 integron and/or <i>sul2</i> location Chr or PL (Kb, Inc group) ^j (no. isolates)
European Clone	A (n=29; 5)	Human (n=26; 16 hospitals; 2 community); Environment (n=1; drinking water); Piggeries (n=2; manure, animal feed)	2003-2010 /N, C, S	A, S, Su, T, Tm (n=1)/ <i>bla</i> _{TEM} , <i>aadA2</i> , <i>su1</i> - <i>sul2</i> , <i>tet</i> (B), <i>dfiA1</i> (1500) (n=1) A, S, Su, T (n=19)/ <i>bla</i> _{TEM} , <i>strA</i> - <i>strB</i> , <i>sul2</i> , <i>tet</i> (B) A, S, Su (n=4)/ <i>bla</i> _{TEM} , <i>strA</i> - <i>strB</i> , <i>sul2</i> S, Su, T (n=1)/ <i>strA</i> - <i>strB</i> , <i>sul2</i> , <i>tet</i> (B) A, T (n=2)/ <i>bla</i> _{TEM} , <i>tet</i> (B) T (n=1)/ <i>tet</i> (B) (-) (n=1) ^g A, S, Su, T (C, G, Tm) (n=16)/ <i>bla</i> _{TEM} , <i>strA</i> - <i>strB</i> , <i>sul2</i> , <i>tet</i> (B) (<i>floR</i> , <i>dfiA12</i> , <i>aadA2</i> , <i>su1</i>) ^h T (n=1)/ <i>tet</i> (B)	<i>dfiA1</i> - <i>orfF</i> - <i>aadA2</i> (1500) (n=1) (-) (n=28)	PL (55, 11) (n=1) and Chr (<i>sul2</i>) (n=4)
non DT104/U302						
ST34 (1) (n=6)						
STM2740 ⁺ , STM2757 ^{+/+} , STM1053-199T ⁻ , <i>fljA</i> ⁻ , <i>fljB</i> ⁻ , <i>hin</i> ⁻ , <i>iroB</i> ⁺						
	B (n=17; 5)	Human (n=17; 10 hospitals)	2008-2010/N, C, S, I	A, S, Su, T (n=16)/ <i>bla</i> _{TEM} , <i>strA</i> - <i>strB</i> , <i>sul2</i> , <i>tet</i> (B) (<i>floR</i> , <i>dfiA12</i> , <i>aadA2</i> , <i>su1</i>) ^h T (n=1)/ <i>tet</i> (B) (-) (n=16)	<i>dfiA12</i> - <i>orfF</i> - <i>aadA2</i> ^h (2000) (n=1) (-) (-) (n=16)	PL (230, HI2) ^h (n=1) and Chr (<i>sul2</i>) (n=8)
	C (n=19; 5)	Human (n=19; 15 hospitals)	2007-2010/N, C, S, I	A, S, Su, T (K) (n=14)/ <i>bla</i> _{TEM} , <i>strA</i> - <i>strB</i> , <i>sul2</i> , <i>tet</i> (B) A, S, Su (n=1)/ <i>bla</i> _{TEM} , <i>strA</i> - <i>strB</i> , <i>sul2</i> S, Su, T (n=3)/ <i>strA</i> - <i>strB</i> , <i>sul2</i> , <i>tet</i> (B) T (n=1)/ <i>tet</i> (B) A, S, Su, T (n=1)/ <i>bla</i> _{TEM} , <i>strA</i> - <i>strB</i> , <i>sul2</i> , <i>tet</i> (B)	(-) (n=19) (-) (n=1)	Chr (<i>sul2</i>) (n=5) Chr (<i>sul2</i>) (n=1)
Spanish clone	O (n=44; 8)	Human (n=30; 20 hospitals); Food (n=8; poultry, pork, beef); Environment (n=3; bathing water); Piggery (n=2; manure); Unknown (n=1)	2002-2010/N, C, S	A, C, G, S, Su, T, Tm (K, Nx) (n= 24)/ <i>bla</i> _{TEM} , <i>cmiA1</i> , <i>aac</i> (3)-IV, <i>aadA2</i> - <i>aadA1</i> , <i>su1</i> - <i>sul2</i> - <i>sul3</i> , <i>tet</i> (A), <i>dfiA12</i> (<i>floR</i>)	<i>dfiA12</i> - <i>orfF</i> - <i>aadA2</i> (2000) (n=27)	PL (110–220, A/C) (n=8)

Table 1 (continued)

Clone designation Phage type ^a MLST-ST (eBG) (no. isolates) ^b Predominant <i>fljAB</i> deletion pattern ^c	PFGE-type (no. isolates/ no. PFGE subtypes) ^d	Source (no. isolates)	Year/ region ^e	Resistance phenotype (no. isolates) ^f / Resistance genes profile	Class 1 integron (bp) ⁱ (no. isolates)	Class 1 integron and/or <i>sul2</i> location Chr or PL (Kb, Inc group) ^g (no. isolates)
DT104/U302				A, C, G, S, Su, Tm (n=2)/ <i>bla</i> _{TEM} , <i>cmlA1</i> , <i>aac</i> (3)-IV, <i>aadA2</i> - <i>aadA1</i> , <i>sul1-sul2-sul3</i> , <i>dfiA12</i> (<i>floR</i>)	Type III- <i>sul3</i> (7304) (n=43) (-) (n=1)	PL (170; A/C, N) (n=1)
ST19 (I) (n=5)				A, C, G, S, Su, T (n=2)/ <i>bla</i> _{TEM} , <i>cmlA1</i> , <i>aac</i> (3)-IV, <i>aadA2</i> - <i>aadA1</i> , <i>sul1-sul2-sul3</i> , <i>tet</i> (A)		
STM2740 ⁺ , STM2757 ⁺ , STM1053-1997 ⁺ , <i>fljA</i> ⁺ , <i>fljB</i> ⁻ , <i>hin</i> ⁻ , <i>iroB</i> ⁻				A, C, S, Su, T, Tm (n=1)/ <i>bla</i> _{TEM} , <i>cmlA1</i> , <i>aadA2-aadA1</i> , <i>sul1-</i> <i>sul2-sul3</i> , <i>tet</i> (A), <i>dfiA12</i> C, G, Su, T (S) (n=4)/ <i>cmlA1</i> , <i>aac</i> (3)-IV, <i>sul1-sul2-sul3</i> , <i>tet</i> (A), <i>aadA2-aadA1</i> C, S, Su, T (Nx) (n=7)/ <i>cmlA1</i> , <i>aadA2-aadA1</i> , <i>sul1-sul2-sul3</i> , <i>tet</i> (A) C, G, S, Su (n=1)/ <i>cmlA1</i> , <i>aac</i> (3)- IV, <i>aadA2-aadA1</i> , <i>sul1-sul2-sul3</i> C, S, Su (n=1)/ <i>cmlA1</i> , <i>aadA2-</i> <i>aadA1</i> , <i>sul1-sul2-sul3</i> S, Su, T (n=1)/ <i>aadA2-aadA1</i> , <i>sul1-sul2-sul3</i> , <i>tet</i> (A) Su (n=1)/ <i>sul1-sul2</i>		
	X (n=2; 1)	Human (n=2; 1 hospital)	2004/S	A, C, G, S, Su, T, Tm (n=2)/ <i>bla</i> _{TEM} , <i>cmlA1</i> , <i>aac</i> (3)-IV, <i>aadA2-aadA1</i> , <i>sul1-sul2-sul3</i> , <i>tet</i> (A), <i>dfiA12</i>	<i>dfiA12-orfF-aadA2</i> (2000) (n=2) Type III- <i>sul3</i> (7304) (n=2)	PL (130, A/C) (n=1)
Southern European Clone	Q (n=9; 4)	Human (n=8; 8 hospitals); Food (n=1; pork)	2004, 2006–2010/N, C, S	C, S, Su, T, Tm (n=7)/ <i>cmlA1</i> , <i>aadA2-aadA1</i> , <i>sul3</i> , <i>tet</i> (B), <i>dfiA12</i> (<i>strA-strB</i>) T (n=2)/ <i>tet</i> (B)	Type I- <i>sul3</i> (7085) (n=7) (-) (n=2)	PL (80–160, R) (n=3)

Table 1 (continued)

Clone designation Phage type ^a MLST-ST (eBG) (no. isolates) ^b Predominant <i>fljAB</i> deletion pattern ^c	PFGE-type (no. isolates/ no. PFGE subtypes) ^d	Source (no. isolates)	Year/ region ^e	Resistance phenotype (no. isolates) ^f / Resistance genes profile	Class 1 integron (bp) ^j (no. isolates)	Class 1 integron and/or <i>sul2</i> location Chr or PL (Kb, Inc group) ^j (no. isolates)
non DT104/U302	G (n=7; 3)	Human (n=7; 6 hospitals)	2003–2004, 2009– 2010/N	C, S, Su, T, Tm (n=7)/ <i>cmiA1</i> , <i>aadA2-aadA1</i> , <i>sul3</i> , <i>tet(B)</i> , <i>dfiA12 (strA-strB)</i>	Type I- <i>sul3</i> (7085) (n=7)	PL (110–140, R) (n=5)
ST19 (I) (n=5)	F (n=2; 2)	Human (n=2; 2 hospitals)	2008–2009/N	C, S, Su, T, Tm (n=1)/ <i>cmiA1</i> , <i>aadA2-aadA1</i> , <i>sul3</i> , <i>tet(B)</i> , <i>dfiA12 (strA-strB)</i>	Type I- <i>sul3</i> (7085) (n=1)	PL (nd, R) (n=1)
STM2740 ⁻ , STM2757 ⁻ , STM1053-1997 ⁺ , <i>fljA</i> ⁻ , <i>fljB</i> ⁻ , <i>hin</i> ⁺ , <i>iroB</i> ⁺	E (n=1; 1)	Human (n=1; 1 hospital)	2007/S	T (n=1)/ <i>tet(B)</i> (-) (n=1) ^g (-) (n=1)	(-) (n=1) (-) (n=1)	(-)

^a PCR assay for the identification of *Salmonella enterica* serotype Typhimurium DT104/U302 [16]^b ST, sequence type; eBG, eBurst Groups identified by Multilocus Sequence Typing. Number of isolates submitted to *Salmonella* MLST Database^c Pattern of genomic deletions surrounding the *fljAB* operon, including *fljA*, *fljB*, *hin*, *iroB*, *STM2740* and *STM2757* genes and *STM1053*-1997 region by PCR according to Soyer et al. [13]; +, presence; -, absence^d Clones are designated by capital letters^e N, North; C, Centre; S, South and I, Islands (Azores)^f Variable presence of resistance phenotype and genotype is indicated in parenthesis. Antimicrobial abbreviations: A, amoxicillin; C, chloramphenicol; Cp, ciprofloxacin; G, gentamicin; K, kanamycin; Nx, nalidixic acid; S, streptomycin; Su, sulfamethoxazole; T, tetracycline; Tm, trimethoprim^g (-), susceptible to all the antibiotics tested^h The transferability of underlined genes was only verified in one isolateⁱ (-), absence of class 1 integron. Atypical integron characterized: Type I-*sul3* (5'CS-*dfiA12-orfF-aadA2-cmiA1-aadA1-qacH-IS440-sul3*); Type III-*sul3* (5'CS-*extX-psp-aadA2-cmiA1-qacH-IS440-sul3*) [19]^j Chromosomal (Chr) and/or plasmid (PL) location of integron and/or the *sul2* genes was assessed by hybridization of I-*CeuI*/S1-digested genomic DNA using *int1*, *sul1*, *sul2*, *sul3* and rep probes; nd, not determined

25, 26]. Here, the dominant PFGE-type A, corresponding to STYMXB.0079 profile particularly widespread in Italy [11, 25], was found in human clinical samples and piggeries (manure and animal feed), suggesting that pig-production may be its main reservoir in Portugal, as demonstrated for other European *S.* 4,[5],12:i:- PFGE-types (e.g., STYMXB.0131 in Germany; STYMXB.0010 in Greece) [9, 26]. Common pig breeding lines and commercial trade in living animals, feed and meat products could also explain the spread of these MDR clones in Europe [28]. The ASSuT pattern was detected in all PFGE-types and was associated with chromosomal resistance genes [*bla*_{TEM-1}, *strA-strB*, *sul2* and *tet(B)*], recently localized in two highly conserved resistance regions in STYMXB.0079-ASSuT strains [11]. However, the detection of isolates with other resistant patterns (ASSu, SSuT, AT, T) along with diverse results obtained for RR1 and RR2 left, right and internal regions in ASSuT isolates, suggests rearrangements or deletions within those chromosomal resistance regions of the European clone. Furthermore, we found several patterns of genomic deletions surrounding the *fljAB* region, including those previously described for European clone isolates [11] or in atypical US strains (32 % of our isolates) [13], suggesting that multiple independent emergence events occurred in ST34 clonal lineage, most likely from *S.* Typhimurium.

The isolates belonging to the Spanish clone presented phenotypic and genotypic markers previously described [10, 29]. This clonal lineage, originally identified in Spain where it still predominates [10, 29], was subsequently reported in food products, environment or piggeries and is causing human infections in neighboring Portugal since 2002 [8, 19]. Intense commercial trade between Portugal and Spain might explain the maintenance and dissemination of the Spanish clone in the Iberian Peninsula. Most isolates studied presented the core ACGSSuTTm R-type, related with class 1 integrons, located on large non-transferable IncA/C or IncA/C-N plasmids, as described in Spain [10]. Acquisition of those epidemic plasmids, carrying multiple antibiotic resistance determinants and/or virulence genes [10, 30], could contribute to the Spanish clone maintenance and spread. Moreover, all MDR isolates carried the same genomic deletion pattern surrounding the *fljAB* region as described for few Spanish isolates studied, suggesting that this clone represents a distinct genetic lineage that evolved through an independent event from *S.* Typhimurium [13].

Of note, is the description of a third MDR *S.* 4,[5],12:i:- group within the worldwide distributed ST19, circulating, at least, since 2003 in Portugal and initially identified as *S.* Typhimurium [19]. Their discrimination from European and Spanish clones comprises several features, including PFGE profiles, MDR phenotypes and plasmid-integron encoded resistance determinants. Moreover, when tested by PCR according to Soyer et al. [13] the same particular genomic deletion

pattern surrounding the *fljAB* operon was found in all isolates, presenting *hin* and *iroB* genes and the STM1053-1997 region (all absent from Spanish clone) but not *fljA*, *fljB*, STM2740 and STM2757 genes. This pattern is also shared by ST19 USA strains [13], although IncR plasmids and MDR profile characterizing our isolates were absent (reference strain CVM23701, GenBank accession no. NZ_ABAO00000000.1). These data suggest a common evolutionary origin for US strains and Southern European clone, with the acquisition of MDR IncR plasmids by the latter, possibly driven by antibiotic selective pressure and availability of IncR in the European metagenome [8, 21, 31]. Moreover, all but three Southern European clone isolates presented the MDR pattern CSSuTTm associated with a genetic platform including an atypical *sul3* class 1 integron, previously described in Portuguese isolates [19], and *tet(B)*, both located on large non-transferable IncR plasmids. Identification of those features in few sporadic *S.* 4,[5],12:i:- Italian (human) and Spanish (swine) isolates [12] suggests that this clone might be spread in other countries. IncR plasmids have been described in different *Enterobacteriaceae* and associated with high-risk clones and/or clinically relevant genes (e.g. *bla*_{VIM}, *bla*_{CTX-M}, and *qnr* genes) [21, 30, 31], including in the animal setting (e.g. *S.* Rissen in Portuguese piggeries) [8]. Simultaneous presence of resistance genes to different antibiotics and/or virulence [12] could contribute to successful persistence of IncR plasmids by diverse co-selection events and provide selective advantages for this emerging *S.* 4,[5],12:i:- clone. Its potential reservoir and transmission pathways (e.g., food and animal sources, travel outside European Union), however, remain unknown and require further studies.

In summary, in this nine-year period study, we identified three *S.* 4,[5],12:i:- clones of particular PFGE-types, specific MDR profiles and genetic elements circulating in Portugal, with two of those related to the highly disseminated European and Spanish clones. We described a third successful MDR clonal lineage associated with human infections, originated by a *fljAB*-deletion event distinct from the European and Spanish clones and similar to the US strains, carrying several clinically important antibiotic resistance genes located in newly described IncR plasmids, which could contribute to higher prevalence of *Salmonella* serotype 4,[5],12:i:-. Moreover, a combination of molecular features typical of each clone (e.g., PFGE-type, *sul3*-type I integron and IncR-Southern European clone; PFGE-type, *sul3*-type III integron and IncA/C-Spanish clone; PFGE-type, ASSuT-RR1/RR2-European clone) may constitute biomarkers for tracing MDR *Salmonella* 4,[5],12:i:- clones with potential for clinical diagnosis and epidemiological surveillance. Nevertheless, further studies are needed to assess the contribution of other driving forces (e.g., wide use of non-antibiotic compounds in biocides and/or animal feed supplements) and/or adaptive traits (e.g., biofilm production) for the success of *S.* 4,[5],12:i:- in order to contain its ongoing dissemination.


Acknowledgments The authors wish to thank Claudia Lucarelli for kindly providing the *Xba*I PFGE profiles designated according to PulseNet for comparison and to Laura Dias, Liliana Ferreira and Tatiana Alves for technical assistance. This study was funded by research grants from the Fundação para a Ciência e a Tecnologia (PEst-C/EQB/LA0006/2013) and Investigação Jovem da Universidade do Porto/Santander Totta (PP_IJUP2010-17). Joana Mourão was supported by a Ph.D. fellowship from Fundação para a Ciência e a Tecnologia (grant number SFRH/BD/77518/2011; POPH-QREN).

Conflict of interest The authors declare that they have no conflict of interest.

References

- European Food Safety Authority (2014) The European union summary report on trends and sources of zoonoses, zoonotic agents and food-borne outbreaks in 2012. EFSA J 12(2):35–47
- Majowicz SE, Musto J, Scallan E, Angulo FJ, Kirk M, O'Brien SJ, Jones TF, Fazil A, Hoekstra RM (2010) The global burden of nontyphoidal *Salmonella* gastroenteritis. Clin Infect Dis 50(6):882–889
- European Food Safety Authority (2010) EFSA panel on biological hazards (BIOHAZ): Scientific opinion on monitoring and assessment of the public health risk of “*Salmonella* Typhimurium-like” strains. EFSA J 8(10):1826
- Hopkins KL, Kirchner M, Guerra B, Granier SA, Lucarelli C, Porrero MC, Jakubczak A, Threlfall EJ, Mevius DJ (2010) Multiresistant *Salmonella enterica* serovar 4,[5],12:i:- in Europe: a new pandemic strain? Euro Surveill 15(22):19580
- Switt AI, Soyer Y, Warnick LD, Wiedmann M (2009) Emergence, distribution, and molecular and phenotypic characteristics of *Salmonella enterica* serotype 4,5,12:i:-. Foodborne Pathog Dis 6(4):407–415
- Gossner CM, van Cauteren D, Le Hello S, Weill FX, Terrien E, Tessier S, Janin C, Brisabois A, Dusch V, Vaillant V, Jourdan-da Silva N (2012) Nationwide outbreak of *Salmonella enterica* serotype 4,[5],12:i:- infection associated with consumption of dried pork sausage, France, November to December 2011. Euro Surveill 17(5)
- Raguenaud ME, Le Hello S, Salah S, Weill FX, Brisabois A, Delmas G, Germonneau P (2012) Epidemiological and microbiological investigation of a large outbreak of monophasic *Salmonella* Typhimurium 4,5,12:i:- in schools associated with imported beef in Poitiers, France, October 2010. Euro Surveill 17(40):20289
- Antunes P, Mourão J, Pestana N, Peixe L (2011) Leakage of emerging clinically relevant multidrug-resistant *Salmonella* clones from pig farms. J Antimicrob Chemother 66(9):2028–2032
- Hauser E, Tietze E, Helmuth R, Junker E, Blank K, Prager R, Rabsch W, Appel B, Fruth A, Malorny B (2010) Pork contaminated with *Salmonella enterica* serovar 4,[5],12:i:-, an emerging health risk for humans. Appl Environ Microbiol 76(14):4601–4610
- Garcia P, Guerra B, Bances M, Mendoza MC, Rodicio MR (2011) IncA/C plasmids mediate antimicrobial resistance linked to virulence genes in the Spanish clone of the emerging *Salmonella enterica* serotype 4,[5],12:i:-. J Antimicrob Chemother 66(3):543–549
- Lucarelli C, Dionisi AM, Filetici E, Owczarek S, Luzzi I, Villa L (2012) Nucleotide sequence of the chromosomal region conferring multidrug resistance (R-type ASSuT) in *Salmonella* Typhimurium and monophasic *Salmonella* Typhimurium strains. J Antimicrob Chemother 67(1):111–114
- Garcia P, Hopkins KL, Garcia V, Beutlich J, Mendoza MC, Threlfall J, Mevius D, Helmuth R, Rodicio MR, Guerra B, Med-Vet-Net WPPG (2014) Diversity of plasmids encoding virulence and resistance functions in *Salmonella enterica* subsp. *enterica* Serovar Typhimurium monophasic variant 4,[5],12:i:- Strains circulating in Europe. PLoS One 9(2):e89635
- Soyer Y, Moreno Switt A, Davis MA, Maurer J, McDonough PL, Schoonmaker-Bopp DJ, Dumas NB, Root T, Warnick LD, Grohn YT, Wiedmann M (2009) *Salmonella enterica* serotype 4,5,12:i:-, an emerging *Salmonella* serotype that represents multiple distinct clones. J Clin Microbiol 47(11):3546–3556
- Council Regulation (EC) No. 517/2011 of 25 May 2011 implementing Regulation (EC) No 2160/2003 of the European Parliament and of the Council as regards a Union target for the reduction of the prevalence of certain *Salmonella* serotypes in laying hens of *Gallus gallus* and amending Regulation (EC) No 2160/2003 and Commission Regulation (EU) No 200/2010. Official Journal of the European Communities L138/45
- Tennant SM, Diallo S, Levy H, Livio S, Sow SO, Tapia M, Fields PI, Mikoleit M, Tamboura B, Kotloff KL, Nataro JP, Galen JE, Levine MM (2010) Identification by PCR of non-typhoidal *Salmonella enterica* serovars associated with invasive infections among febrile patients in Mali. PLoS Negl Trop Dis 4(3):e621
- Pritchett LC, Konkel ME, Gay JM, Besser TE (2000) Identification of DT104 and U302 phage types among *Salmonella enterica* serotype typhimurium isolates by PCR. J Clin Microbiol 38(9):3484–3488
- Clinical and Laboratory Standards Institute (CLSI) (2013) Performance standards for antimicrobial susceptibility testing: twenty-first informational supplement M100-S23. CLSI, Wayne, IN, USA
- Antunes P, Machado J, Peixe L (2006) Characterization of antimicrobial resistance and class 1 and 2 integrons in *Salmonella enterica* isolates from different sources in Portugal. J Antimicrob Chemother 58(2):297–304
- Antunes P, Machado J, Peixe L (2007) Dissemination of *sul3*-containing elements linked to class 1 integrons with an unusual 3' conserved sequence region among *Salmonella* isolates. Antimicrob Agents Chemother 51(4):1545–1548
- Carattoli A, Bertini A, Villa L, Falbo V, Hopkins KL, Threlfall EJ (2005) Identification of plasmids by PCR-based replicon typing. J Microbiol Methods 63(3):219–228
- Garcia-Fernandez A, Fortini D, Veldman K, Mevius D, Carattoli A (2009) Characterization of plasmids harbouring *qnrS1*, *qnrB2* and *qnrB19* genes in *Salmonella*. J Antimicrob Chemother 63(2):274–281
- Silveira L, Marques A, Machado J (2013) Infecções por *Salmonella enterica* no período entre 2000–2012. Boletim Epidemiológico do Instituto Nacional de Saúde Dr Ricardo Jorge 2013. <http://repositorio.insa.pt/>. Accessed 08 April 2014
- European Medicines Agency, European Surveillance of Veterinary Antimicrobial Consumption (2013) Sales of veterinary antimicrobial agents in 25 EU/EEA countries in 2011. EMA/236501/2013
- DANMAP (2012) Use of Antimicrobial Agents and Occurrence of Antimicrobial Resistance in Bacteria from Food Animals, Foods and Humans in Denmark. <http://www.danmap.org/>. Accessed 08 April 2014
- Lucarelli C, Dionisi AM, Torpdahl M, Villa L, Graziani C, Hopkins K, Threlfall J, Caprioli A, Luzzi I (2010) Evidence for a second genomic island conferring multidrug resistance in a clonal group of strains of *Salmonella enterica* serovar Typhimurium and its monophasic variant circulating in Italy, Denmark, and the United Kingdom. J Clin Microbiol 48(6):2103–2109
- Mandilara G, Lambiri M, Polemis M, Passiotou M, Matopoulos A (2013) Phenotypic and molecular characterisation of multiresistant

- monophasic *Salmonella* Typhimurium (1,4,[5],12:i:-) in Greece, 2006 to 2011. Euro Surveill 18(22)
27. Mulvey MR, Finley R, Allen V, Ang L, Bekal S, El Bailey S, Haldane D, Hoang L, Horsman G, Louie M, Robberts L, Wylie J, McCracken M, Langner S, Ahmed R, Tabor H, Gilmour M (2013) Emergence of multidrug-resistant *Salmonella enterica* serotype 4,[5],12:i:- involving human cases in Canada: results from the Canadian integrated program on antimicrobial resistance surveillance (CIPARS), 2003–10. J Antimicrob Chemother 68(9):1982–1986
28. European Food Safety Authority (2011) Scientific report of EFSA: analysis of the baseline survey of *Salmonella* in holdings with breeding pigs, in the EU, 2008; Part B: analysis of factors potentially associated with *Salmonella* pen positivity. EFSA J 9(7):2329
29. Garcia P, Malorny B, Hauser E, Mendoza MC, Rodicio MR (2013) Genetic types, gene repertoire, and evolution of isolates of the *Salmonella enterica* serovar 4,5,12:i:- Spanish clone assigned to different phage types. J Clin Microbiol 51(3):973–978
30. Carattoli A (2009) Resistance plasmid families in Enterobacteriaceae. Antimicrob Agents Chemother 53(6):2227–2238
31. Novais A, Rodrigues C, Branquinho R, Antunes P, Grosso F, Boaventura L, Ribeiro G, Peixe L (2012) Spread of an OmpK36-modified ST15 *Klebsiella pneumoniae* variant during an outbreak involving multiple carbapenem-resistant Enterobacteriaceae species and clones. Eur J Clin Microbiol Infect Dis 31(11):3057–3063



Online Resource 1 XbaI profiles of *Salmonella enterica* 4,[5],12:i:- isolates were analysed by InfoQuest FP version 5.4 software (BioRad Laboratories), and the percentage of similarity was calculated by applying the unweighted-pair group method using average linkages (UPGMA) algorithm based on the Dice coefficient (1.0% band tolerance; 1.0% optimization). A rectangle highlights the most prominent PFGE profiles in Europe, A and B corresponding to STYMXB.0079 and STYMXB.0131 respectively (designation according to PulseNet Europe)

3.2. Analysis of the dispersion of metals/biocides tolerance genes and tolerance phenotype as well as their associated genetic platforms in diverse non-typhoidal *Salmonella* serotypes and clones

- 3.2.1. Metal tolerance in emerging clinically relevant multidrug-resistant *Salmonella enterica* serotype 4,[5],12:i:- clones circulating in Europe.
- 3.2.2. Tolerance to multiple metal stressors in emerging non-typhoidal MDR *Salmonella* serotypes: a relevant role for copper in anaerobic conditions.
- 3.2.3. *Salmonella enterica* serotype Bovismorbificans, a new host for CTX-M-9.
- 3.2.4. Clinical *Salmonella* Typhimurium ST34 with metal tolerance genes and an IncHI2 plasmid carrying *oqxAB-aac(6')-Ib-cr* from Europe.

Metal tolerance in emerging clinically relevant multidrug-resistant *Salmonella enterica* serotype 4,[5],12:i:- clones circulating in Europe

Joana Mourão¹, Carla Novais¹, Jorge Machado², Luísa Peixe¹ and Patrícia Antunes^{1,3*}

¹ UCIBIO/REQUIMTE, Departamento de Ciências Biológicas, Laboratório de Microbiologia, Faculdade de Farmácia, Universidade do Porto, Porto, Portugal; ² Laboratório Nacional de Referência de Infecções Gastrointestinais, Departamento de Doenças Infecciosas, Instituto Nacional de Saúde Dr. Ricardo Jorge, Lisboa, Portugal; ³ Faculdade de Ciências da Nutrição e Alimentação, Universidade do Porto, Porto, Portugal.

International Journal of Antimicrobial Agents 2015 June; 45(6):610-616

Elsevier has authorized the reproduction of the final published PDF version of this paper in this thesis through the License Agreement Number 3791960546294 on January 18, 2016.



Contents lists available at ScienceDirect

International Journal of Antimicrobial Agents

journal homepage: <http://www.elsevier.com/locate/ijantimicag>

Metal tolerance in emerging clinically relevant multidrug-resistant *Salmonella enterica* serotype 4,[5],12:i:– clones circulating in Europe

Joana Mourão^a, Carla Novais^a, Jorge Machado^b, Luísa Peixe^a, Patrícia Antunes^{a,c,*}^a UCIBIO/REQUIMTE, Departamento de Ciências Biológicas, Laboratório de Microbiologia, Faculdade de Farmácia, Universidade do Porto, Porto, Portugal^b Laboratório Nacional de Referência de Infecções Gastrointestinais, Departamento de Doenças Infecciosas, Instituto Nacional de Saúde Dr Ricardo Jorge, Lisbon, Portugal^c Faculdade de Ciências da Nutrição e Alimentação, Universidade do Porto, Porto, Portugal

ARTICLE INFO

Article history:

Received 3 October 2014

Accepted 16 January 2015

Keywords:

Biocides

Copper

Silver

Multidrug resistance

Salmonella enterica Typhimurium

monophasic variant

ABSTRACT

The occurrence of acquired metal tolerance genes in emerging MDR *Salmonella enterica* serotype 4,[5],12:i:– clones was assessed and their associated platforms and tolerance phenotype were characterised. *Salmonella* 4,[5],12:i:– from different sources belonging to European, Spanish and Southern European clones were studied. Screening for copper (*pcoA-pcoD/tcrB*), silver/copper (*silA-silE*), mercury (*merA*), arsenic (*arsB*) and tellurite (*terF*) tolerance genes was performed by PCR/sequencing. $\text{CuSO}_4/\text{AgNO}_3$ MICs were determined in aerobic/anaerobic atmospheres by agar dilution. Conjugation assays, genomic location and plasmid analysis were performed by standard procedures. Most isolates from European (98%) and Spanish (74%) clones carried *silA-silE*, contrasting with the Southern European clone (26%). *merA*/62% (European and Spanish clones) and *pcoA-pcoD*/50% (European clone) were also detected. *merA*±*pco*±*sil* were chromosomally located in the European clone, whereas in Spanish and Southern European clones *sil*±*merA* were within plasmids, both with antibiotic resistance genes. The *pcoA-pcoD/silA-silE*⁺ isolates showed higher $\text{MIC}_{\text{CuSO}_4}$ in anaerobiosis than those without these genes (MIC_{50} = 24–28 vs. 2 mM). Different $\text{MIC}_{\text{AgNO}_3}$ of *silA-silE*⁺ (MIC_{50} = 0.25 mM) and *silA-silE*[–] (MIC_{50} = 0.16 mM) isolates were observed in both atmospheres, with an MIC increment after prior exposure to silver (>3 vs. 0.08–0.125 mM) in aerobiosis. A high frequency of copper and silver tolerance, particularly among the two major *Salmonella* 4,[5],12:i:– MDR clones (European/Spanish) circulating in Europe and causing human infections, might facilitate adaptation/expansion of these strains in metal-contaminated environments, particularly copper in anaerobiosis. Furthermore, metal toxic concentrations in food-animal environments can contribute to persistence of genetic platforms carrying metal/antibiotic resistance genes in this foodborne zoonotic pathogen.

© 2015 Elsevier B.V. and the International Society of Chemotherapy. All rights reserved.

1. Introduction

Salmonella enterica subsp. *enterica* serotype 4,[5],12:i:–, a monophasic variant of pathogenic *S. enterica* subsp. *enterica* serotype Typhimurium, has become a new epidemic multidrug-resistant (MDR) serotype in Europe associated with human infections [1,2]. Currently, two major clones are recognised and were previously designated as ‘European’ and ‘Spanish’. They are spread in diverse geographical regions of Europe particularly since

2002 [2,3] or are mainly located in the Iberian Peninsula, respectively [4,5]. A third less frequent clone, the Southern European clone, was recently described in Portugal [5] and was also found among a few sporadic Italian and Spanish isolates [4]. All of these clones, carrying a combination of antibiotic resistance molecular features, account for *Salmonella* 4,[5],12:i:– spread through the food chain, with food animals (e.g. pigs) indicated as their reservoir [2–5]. Selective pressure due to massive use of antibiotics in food-producing animals has been implicated in their selection and maintenance, but the possibility of co-selection of these successful MDR *Salmonella* clones by other compounds widely used in animal settings (e.g. biocides, metals) remains less explored.

Non-antibiotic compounds with antimicrobial activity are currently used in animal farming management (biosecurity, hygiene) owing to antibiotic restriction directives and/or to limit foodborne zoonotic pathogenic bacteria such as

* Corresponding author. Present address: Faculdade de Ciências da Nutrição e Alimentação, Universidade do Porto, Rua Dr Roberto Frias, 4200–465 Porto, Portugal. Tel.: +351 22 507 4320; fax: +351 22 507 4329.

E-mail addresses: patriciaantunes@fcna.up.pt, patantunes@gmail.com (P. Antunes).

Table 1
Epidemiological and antibiotic and metal tolerance genetic features of *Salmonella enterica* subsp. *enterica* serotype 4,[5],12:i:– isolates from Portugal.

Clone designation Phage type (no. of isolates)/ST-eBG; ^a Predominant antibiotic resistance phenotype (no. of isolates) and genotype	PFGE type (no. of isolates; no. of PFGE subtypes) ^b	Source (no. of isolates)	Year(s)/region(s) ^c	Occurrence of metal tolerance genes [no. of isolates (%)]			Metal tolerance genes; location (kb, Inc) (no. of isolates)	Co-location with antibiotic resistance genes (no. of isolates)
				<i>pcoA-pcoD</i>	<i>silA-silE</i>	<i>merA</i>		
European clone Non DT104/U302 (n=66/ST34-1) AMX, STR, SUL, TET (n=51) <i>bla</i> _{TEM-1} , <i>strA-strB</i> , <i>sul2</i> , <i>tet(B)</i>	A (n=29; 5)	Human (n=26; 16 hospitals, 2 community labs)	2003–2010/N, C, S	29 (100%)	29 (100%)	11 (38%)	<i>pcoD</i> + <i>silA</i> + <i>merA</i> ; <i>Chr</i> ^d (n=4) <i>pcoD</i> + <i>silA</i> ; <i>Chr</i> (n=3)	<i>bla</i> _{TEM-1} and/or <i>strA-strB</i> , <i>sul2</i> , <i>tet(B)</i> (n=7)
		Environment (n=1; drinking water) Piggeries (n=2; manure, animal feed)						
	B (n=17; 5)	Human (n=17; 10 hospitals)	2008–2010/N, C, S, I	17 (100%)	17 (100%)	14 (82%)	<i>pcoD</i> + <i>silA</i> + <i>merA</i> ; <i>Chr</i> ^d (n=9) <i>pcoD</i> + <i>silA</i> ; <i>Chr</i> (n=1) <i>merA</i> and/or <i>pcoD</i> + <i>silA</i> ; <i>Chr</i> ^d (n=4) <i>pcoD</i> + <i>silA</i> ; <i>Chr</i> (n=1)	<i>bla</i> _{TEM-1} , <i>strA-strB</i> , <i>sul2</i> , <i>tet(B)</i> (n=9) <i>tet(B)</i> (n=1)
		Human (n=19; 15 hospitals)						
Spanish clone DT104/U302 (n=46/ST19-1) AMX, CHL, GEN, STR, SUL, TET, TMP (n=26) <i>bla</i> _{TEM-1} , <i>cmiA1</i> , <i>aac(3)-IV</i> , <i>addA2-addA1</i> , <i>sul1-sul2-sul3</i> , <i>tet(A)</i> , <i>dfiA12</i>	D (n=1; 1)	Human (n=1; 1 hospital)	2009/N	1 (100%)	1 (100%)	1 (100%)	<i>pcoD</i> + <i>silA</i> + <i>merA</i> ; <i>Chr</i> ^d (n=1)	<i>bla</i> _{TEM-1} , <i>strA-strB</i> , <i>sul2</i> , <i>tet(B)</i> (n=1)
		Human (n=31; 20 hospitals)						
	O (n=44; 8)	Food (n=8; poultry, pork, beef) Environment (n=3; bathing water) Piggery (n=2; manure)	2002–2010/N, C, S	n=65 (98%) 0 (0%)	n=65 (98%) 32 (73%)	n=39 (59%) 40 (91%)	<i>silA</i> + <i>merA</i> ; <i>PL</i> (170; A/C, N) (n=1)	<i>bla</i> _{TEM-1} , <i>cmiA1</i> , <i>aac(3)-IV</i> , <i>addA2-addA1</i> , <i>sul1-sul2-sul3</i> , <i>tet(A)</i> , <i>dfiA12</i> ^e (n=1)
		Human (n=2; 1 hospital)						
X (n=2; 1)	n=46	Human (n=2; 1 hospital)	2004/S	0 (0%)	2 (100%)	2 (100%)	<i>silA</i> + <i>merA</i> ; <i>PL</i> (110; A/C) (n=1) <i>silA</i> + <i>merA</i> ; <i>PL</i> (220; A/C) (n=1)	<i>bla</i> _{TEM-1} , <i>cmiA1</i> , <i>aac(3)-IV</i> , <i>addA2-addA1</i> , <i>sul1-sul3</i> , <i>tet(A)</i> , <i>dfiA12</i> ^e (n=1)

Table 1 (Continued)

Clone designation Phage type (no. of isolates) ^a Predominant antibiotic resistance phenotype (no. of isolates) and genotype	PFGE type (no. of isolates; no. of PFGE subtypes) ^b	Source (no. of isolates)	Year(s)/region(s) ^c	Occurrence of metal tolerance genes [no. of isolates (%)]			Metal tolerance genes; location (kb, Inc) (no. of isolates)	Co-location with antibiotic resistance genes (no. of isolates)
				<i>pcoA-pcoD</i>	<i>silA-silE</i>	<i>merA</i>		
Southern European clone Non DT104/U302 (n = 19/ST19-1) CHL, STR, SUL, TET, TMP (n = 15)	Q (n = 9; 4)	Human (n = 8; 8 hospitals) Food (n = 1; pork)	2004, 2006–2010/N, C, S 2003–2004, 2009–2010/N	0 (0%)	0 (0%)	0 (0%)	N/A	N/A
<i>cmlA1</i> , <i>aadA2-aadA1</i> , <i>sil3</i> , <i>ter(B)</i> , <i>dfiA12</i>	G (n = 7; 3)	Human (n = 7; 6 hospitals)		0 (0%)	5 (71%)	0 (0%)	<i>silA</i> ; <i>PL</i> (110–140; R) (n = 5)	<i>cmlA1</i> , <i>aadA2-aadA1</i> , <i>sil3</i> , <i>ter(B)</i> , <i>dfiA12</i> ^c (n = 5)
<i>ter(B)</i> , <i>dfiA12</i>	E (n = 1; 1) F (n = 2; 2) n = 19	Human (n = 1; 1 hospital) Human (n = 2; 2 hospitals)	2007/S 2008–2009/N	0 (0%) 0 (0%) n = 0 (0%)	0 (0%) 0 (0%) n = 5 (26%)	0 (0%) 0 (0%) n = 0 (0%)	N/A N/A	N/A N/A

ST, sequence type; eBG, eBURST groups identified by multilocus sequence typing (<http://mlst.ucc.ie/mlst/dbs/Serientica>); PFGE, pulsed-field gel electrophoresis; AMX, amoxicillin; STR, streptomycin; SUL, sulfamethoxazole; TET, tetracycline; CHL, chloramphenicol; GEN, gentamicin; TMP, trimethoprim; Chr, chromosome; PL, plasmid; N/A, not applicable.

^a PCR assay for identification of *Salmonella enterica* serotype Typhimurium DT104/U302 as previously described [5].

^b Clones were designated by capital letters and also include PFGE subtypes as previously described [5].

^c N, North; C, Centre; S, South; I, Islands (Azores).

^d Co-location in the same *I-CeuI* chromosomal fragment as antibiotic resistance genes was observed for all isolates with the *merA* gene (n = 18), including 5 isolates also carrying *pcoD* + *silA* (PFGE types B and D).

^e Some genes belong to the atypical *sil3*-type III integron (5'CS-*estX*-*psp*-*aadA2-cmlA1-aadA1-qacH-IS440-sul3*) and/or the conventional class I integron (*dfiA12-orfI-aadA2*) in the Spanish clone or the *sil3*-type I integron (5'CS-*dfiA12-orfI-aadA2-cmlA1-aadA1-qacH-IS440-sul3*) in the Southern European clone [5].

Salmonella (including *Salmonella* 4,[5],12:i:–) in the food supply (<http://eur-lex.europa.eu/legal-content/EN/TXT/PDF/?uri=CELEX:32011R0517&from=EN>) [6,7]. Several metals are suitable for use as disinfectants, antiseptics or preservatives (e.g. silver, copper) and as feed additives for growth promotion of farm animals (e.g. copper, zinc) (<http://eur-lex.europa.eu/legal-content/EN/TXT/PDF/?uri=CELEX:32003R1831&from=EN>) or can be found as contaminants in animal feed (e.g. mercury) (<http://www.efsa.europa.eu/en/efsajournal/doc/654.pdf>) [6,7]. Metals can remain in the production environment and accumulate in toxic concentrations, representing a long-term selective pressure potentially driving co-selection of antibiotic-resistant bacteria, including *Salmonella* [6–8]. Diverse acquired genes encoding for metal tolerance mechanisms have been described in scarce studies of *Salmonella* and other Gram-negative bacteria (e.g. Enterobacteriaceae, *Aeromonas*, *Campylobacter*, *Pseudomonas*) commonly sharing the same ecosystems [9–15]. Among them are efflux systems to overcome toxic concentrations of copper (e.g. *pcoABCDRE* and *trcYAZB*), silver (e.g. *silCFBAPRSE*) or arsenic (e.g. *arsRBC*); mercuric detoxification systems (e.g. *mer* operons); and the tellurite resistance system (e.g. *terZABCDE*). Nevertheless, a comprehensive analysis of metal tolerance, including acquired metal gene dispersion and their genetic context within *Salmonella* from diverse ecological niches or belonging to successful MDR clones, remains poorly explored.

This study assessed the occurrence of acquired metal tolerance genes in emerging MDR *Salmonella* 4,[5],12:i:– clones and characterised their associated genetic platforms and tolerance phenotype.

2. Materials and methods

2.1. Strains and epidemiological background

This study included a comprehensive collection of 131 *Salmonella* 4,[5],12:i:– isolates obtained between 2002 and 2010 from human clinical cases (n = 114 from 28 hospitals and 2 community laboratories) covering all geographical regions of Portugal (North, Centre, South and Azores islands) as well as from food of animal origin (n = 9; pork, beef and poultry), the environment (n = 4; drinking and bathing water) and piggeries (n = 4; manure and animal feed) (Table 1). Clonality by pulsed-field gel electrophoresis (PFGE) and multilocus sequence typing (MLST), MDR profiles, *fljAB* deletion patterns and genetic elements linked to antibiotic resistance were previously characterised for the three discriminated clones (European, Spanish and Southern European) [5]. The occurrence of diverse acquired metal tolerance genes was assessed in all of the isolates from this collection. Representative isolates from different clonal lineages, PFGE types, sources, isolation date and antibiotic resistance phenotypes and genotypes, harbouring or lacking acquired copper (Cu)/silver (Ag) tolerance genes, were selected for phenotypic assays [minimum inhibitory concentration (MIC) determination] and for characterisation of genetic platforms.

2.2. Study of metal tolerance and acquired genes

Different acquired genes previously described among Gram-negative and Gram-positive bacteria and associated with tolerance to toxic concentrations of metals, widely used or present in animal settings, were searched by PCR using the conditions and primers described in Supplementary Table S1. They include genes from different metal export systems, namely two genes of the *pcoABCDRE* cluster [*pcoA* (multicopper oxidase) and *pcoD* (copper inner membrane pump)] [9,16], one gene of the *trcYAZB* cluster [*trcB* (copper export ATPase)] [17], two genes of the *silCFBAPRSE*

cluster [*silA* (silver inner-membrane proton/cation antiporter) and *silE* (silver/copper periplasmic metal binding protein)] [18] and *arsB* (arsenite transmembrane pump) [11]. Other mechanisms related to metal tolerance, such as *merA* (mercuric reductase) [10] and *terF* (tellurite resistance protein), were also included [11]. A positive amplicon from each PCR was sequenced and compared with known GenBank sequences to validate the screening method (Supplementary Table S1).

Supplementary Table S1 related to this article can be found, in the online version, at <http://dx.doi.org/10.1016/j.ijantimicag.2015.01.013>.

MICs were determined by the agar dilution method using Mueller–Hinton II agar plates (bioMérieux, Marcy-l'Étoile, France) supplemented with CuSO₄ (0.25, 0.5, 1, 2, 4, 8, 12, 16, 20, 24, 28, 32 and 36 mM, adjusted to pH 7.2) or AgNO₃ (0.025, 0.06, 0.08, 0.125, 0.16, 0.25, 0.32, 0.5, 1.0 and 1.5 mM, adjusted to pH 7.4) as described previously [19]. Briefly, 1 µL of a 10⁷ CFU/mL suspension of each isolate was applied to the supplemented plates, which were incubated at 37 °C in aerobic and anaerobic (GENbox jar with GENbox anaer. and an anaerobic indicator; bioMérieux) atmospheres and read at 16–20 h. The first CuSO₄/AgNO₃ concentration without visible bacterial growth was considered as the MIC. Since *sil* genes have been described as inducible by silver in an aerobic atmosphere [20], MICs to AgNO₃ were also determined following overnight *sil*⁺ and *sil*[−] cultures (14–16 h) in BBL™ Mueller–Hinton II broth (BD, Franklin Lakes, NJ) supplemented with 25 µM of this metal. *Enterococcus faecium* BM4105RF [17] or *Staphylococcus aureus* ATCC 29213 [21] (both negative for all genes tested) and *Escherichia coli* ED8739 (plasmid pRJ1004 with *pco* genes) [9] or *E. coli* J53 (plasmid pMG101 with *sil* genes) [21] were used as controls in CuSO₄ or AgNO₃ tolerance assays, respectively.

2.3. Characterisation of the genetic elements carrying metal and antibiotic resistance genes

Conjugative transfer of antibiotic resistance and metal tolerance genes was analysed for 20 representative isolates of different clonal lineages, PFGE types, resistance phenotypes/genotypes and sources. Conjugative transfer was performed by the filter-mating method on Mueller–Hinton II agar plates at 1:1 and 1:2 donor:recipient ratios using *E. coli* K802N or *E. coli* K-12 strain BM21R as recipient strains, respectively (both rifampicin- and nalidixic acid-resistant) at 30 °C, 37 °C and 42 °C. Transconjugants were recovered after incubation at 37 °C from Mueller–Hinton II agar or MacConkey agar plates supplemented with nalidixic acid (64 mg/L) plus tetracycline (8 mg/L) or sulfamethoxazole (256 mg/L). Co-transfer of antibiotic resistance and/or metal tolerance was confirmed at phenotypic and genotypic levels as described above. Identification of plasmid incompatibility groups was determined by the PCR-based replicon typing (PBRT) method, including three additional PCR assays for the IncU, IncR and ColE groups as previously described [5]. The genetic location of metal and antibiotic determinants was determined by S1 nuclease (Takara Bio Inc., Shiga, Japan) and I-CeuI (New England Biolabs, Ipswich, MA) digested genomic DNA followed by PFGE and Southern blot hybridisation. This was performed by standard methods using antibiotic [*aac*(3)-IV, *bla*_{TEM-1}, *strA*, *strB*, *sul1*, *sul2*, *sul3*, *tet*(A), *tet*(B), *dfrA12*] and metal [*pcoD*, *silA*, *merA*, *arsB*, *terF*] resistance genes, *int11*, rep intragenic probes (IncA/C, R) and 16S rRNA following the manufacturer's instructions (Gene Images AlkPhos Direct™ Labelling System; Amersham GB/GE Healthcare Life Sciences UK Ltd., Little Chalfont, UK). *Xba*I-digested *S. enterica* serotype Braenderup H9812 (CDC) was used as the reference strain.

3. Results

3.1. Occurrence of acquired genes encoding metal tolerance

The genes *silA*–*silE*, *merA* and *pcoA*–*pcoD* were detected in 79% (104/131), 62% (81/131) and 50% (65/131) of the isolates, respectively. The *silA*–*silE* genes were dispersed among the majority of isolates belonging to the European (98%; all PFGE types) and Spanish (74%; all PFGE types) clones and in a few belonging to the Southern European clone (26%; only PFGE type G), regardless of their source (human, environment, piggeries and food) or resistance profile. In contrast, the *pcoA*–*pcoD* genes were only detected in *silA*–*silE*⁺ isolates from the European clone (98%; *n* = 65) (Table 1). Combinations of *sil* + *merA* (34/42) or *pco* + *sil* + *merA* (38/39) were frequently found among Spanish and European clones, respectively. None of the other searched genes was detected.

3.2. Genetic elements carrying metal and antibiotic resistance genes

Isolates of the European clone presented *pco* + *sil* and/or *merA* in the chromosome. In addition, these metal tolerance genes were often co-located in the same chromosomal I-CeuI PFGE fragment of ca. 750 kb and/or ca. 850 kb as antibiotic resistance genes typical of this clonal lineage [*bla*_{TEM-1}, *strA*–*strB*, *sul2* and *tet*(B)], none of them being transferable by conjugation assays (Table 1). In contrast, Spanish clone isolates carried *sil* + *merA* within ca. 110–220 kb non-transferable IncA/C plasmids, along with a *sul3*-type III class 1 integron (*estX*–*psp*–*aadA2*–*cmlA1*–*aadA1*–IS440–*qacH*–*sul3*), *bla*_{TEM-1}, *aac*(3)-IV, *tet*(A), *dfrA12*, *sul1*–*sul2*, typical of this clonal lineage (Table 1). In the Southern European clone, *sil* was found co-located in ca. 110–140 kb non-transferable IncR plasmids with *tet*(B) and an atypical *sul3*-type I class 1 integron (*dfrA12*–*orfF*–*aadA2*–*cmlA1*–*aadA1*–IS440–*qacH*–*sul3*) genes, only in isolates from PFGE type G (Table 1).

3.3. Susceptibility to CuSO₄ and AgNO₃ in aerobic and anaerobic atmospheres

The MIC_{CuSO₄} distribution was similar for isolates of the three clones, sources and independently of location (chromosome or plasmid) of metal tolerance determinants. Higher MIC_{CuSO₄} values were obtained in aerobic conditions (MIC₅₀ = 32–36 mM) irrespective of the presence or absence of *pcoA*–*pcoD* and/or *silA*–*silE*. However, in an anaerobic atmosphere, the *pcoA*–*pcoD*/*silA*–*silE*⁺ isolates showed higher MIC_{CuSO₄} values than those without these genes (MIC₅₀ = 24–28 mM vs. 2 mM) (Table 2). Regarding AgNO₃, the MIC₅₀ was identical both in aerobic and anaerobic conditions, but differences between isolates with *silA*–*silE* (MIC₅₀ = 0.25 mM) and without *silA*–*silE* (MIC₅₀ = 0.16 mM) in the two atmospheres were observed. In addition, in isolates that were subjected to prior exposure to subinhibitory concentrations of AgNO₃, MICs for those carrying *silA*–*silE* increased to >3 mM, contrasting with *silA*–*silE*[−] isolates (0.08–0.125 mM) (Table 2).

4. Discussion

This is the first study documenting a high occurrence of genes encoding tolerance to copper and silver, particularly dispersed among the two major *Salmonella* 4,5,12:i:–MDR clonal lineages (European and Spanish clones) circulating in Europe and causing human infections. These features might have a potential role in the adaptation and expansion of these strains in metal-contaminated environments.

An association between silver tolerance and presence of the *sil* efflux system among isolates belonging to *Salmonella* epidemic

Table 2
Minimum inhibitory concentrations (MICs) of CuSO₄ and AgNO₃ for *Salmonella enterica* subsp. *enterica* serotype 4,[5],12:i:– isolates.

Metal	Environment	<i>silA-silE</i>	<i>pcoA-pcoD</i>	No. of isolates ^a	MIC (mM) ^b										MIC ₅₀ (mM)	MIC ₉₀ (mM)					
					0.08	0.125	0.16	0.25	0.32	0.5	1	2	>3	20	24	28	32	36			
CuSO ₄	Aerobic	+	+	30												12	14	4	32	36	
		+	–	19													1	8	10	36	
	Anaerobic	–	–	15													7	7	1	32	32
		+	+	30											7	12	10	1	1	24	28
		+	–	19											1	7	9	1	1	28	32
AgNO ₃	Aerobic	–	–	15							1	5	9						2	2	
		+	+	48															0.25	0.32	
	Anaerobic	–	–	16															0.16	0.16	
		+	+	48															0.25	0.5	
		–	–	16															0.16	0.25	
Aerobic/previous contact with 0.025 mM AgNO ₃	+	+	16										16					>3	>3		
	–	–	8															>3	0.125	0.125	

^a The number of isolates included representatives of the three clones: European clone (*n* = 31); Spanish clone (*n* = 19); and Southern European clone (*n* = 14).^b MIC values of CuSO₄ (4, 8, 12 and 16 mM) and AgNO₃ (0.75 and 1.5 mM) that were not assigned to the isolates tested were omitted. *Escherichia coli* ED8739 (plasmid pRJ1004 with *pco* genes; MIC_{CuSO4} = 32–36 mM, aerobiosis) [9] and *Enterococcus faecium* BM4105RF (negative for all genes tested; MIC_{CuSO4} = 8–12 mM, aerobiosis) [17] were used as control strains in Cu assays. Those strains were also used in anaerobic assays: *E. coli* ED8739 (plasmid pRJ1004 with *pco* genes; MIC_{CuSO4} = 16–20 mM) and *E. faecium* BM4105RF (MIC_{CuSO4} = 2–4 mM). *Escherichia coli* J53 (plasmid pMG101 with *sil* genes; MIC_{AgNO3} > 3 mM, aerobiosis) [21] and *Staphylococcus aureus* ATCC 29213 (negative for all genes tested; MIC_{AgNO3} = 0.16–0.25 mM, aerobiosis) [21] were used as control strains in Ag assays. Those strains were also used in anaerobic assays: *E. coli* J53 (plasmid pMG101 with *sil* genes; MIC_{AgNO3} > 3 mM) and *S. aureus* ATCC 29213 (MIC_{AgNO3} = 0.32–0.5 mM).

clones was established in this study. Although *sil* genes are scattered in different bacterial genera [12–14,18], including diverse *Salmonella* serotypes (e.g. GenBank accession nos. AF067954, CP005995, CP006055 and CP007505), an association between the genotype and the corresponding phenotypic behaviour has never been established in a comprehensive *Salmonella* collection. In addition, scarce studies searching for *sil* tolerance genes are confined to non-*Salmonella* Enterobacteriaceae isolates recovered from nosocomial and animal clinical settings [12–14] or environmental bacteria [22], impairing an evaluation of *sil* tolerance gene dispersion and their potential impact on the success of emergent *Salmonella* in particular environments, namely where silver could be widely used (e.g. disinfectants used in water or surfaces in the animal production setting; human and animal antiseptics) [22,23].

Higher tolerance to CuSO₄ in anaerobic conditions for *Salmonella* 4,[5],12:i:– isolates carrying *sil* or *sil*+*pco* genes was detected, suggesting that they could overcome the highly toxic cuprous form (Cu¹⁺) present in anoxic environments [24] (e.g. animal gastrointestinal tract and manure, waste lagoons, or forage/feed from food-producing settings). It was proposed that the *silE* gene may have a role in copper tolerance under anaerobic conditions [18], contrasting with other acquired aerobic copper tolerance gene clusters such as the plasmid-borne *pcoABCDRE* efflux system (e.g. *pcoA* requires oxygen to convert Cu¹⁺ into the less toxic Cu²⁺) in *E. coli* [9,16]. However, similar to another recent study [15], the high MIC_{CuSO4} detected in aerobic conditions, both in *Salmonella* carrying *pco* or not, indicate that these genes might not lead to an increase in copper tolerance in *Salmonella* 4,[5],12:i:– isolates in such environmental conditions. The common use of copper as a feed additive in animal growth promotion is a current concern since it might contribute to the selection and maintenance of copper-tolerant bacteria (<http://www.efsa.europa.eu/en/efsajournal/doc/2969.pdf>) [25]. In fact, the emergence of *Salmonella* 4,[5],12:i:– clones causing human infections has been associated particularly with pig production [1–3,26], which are within the animal settings where higher copper concentrations in feed are allowed (<http://eur-lex.europa.eu/LexUriServ.do?uri=OJ:C:2004:050:0001:0144:EN:PDF>) by the European Union.

Co-localisation of the metal tolerance genes within chromosomal regions or plasmids previously associated with MDR profiles in *Salmonella* 4,[5],12:i:– clones [4,5,27], irrespective of their source, was shown by the molecular studies performed. In fact, it is widely demonstrated in industrialised countries that *Salmonella* MDR genetic platforms are the same both in food-production animals and humans, suggesting a direct link between the two bacterial populations. In the case of the globally spread European *Salmonella* 4,[5],12:i:– clone, *mer* genes were recently found downstream of the RR1–RR2 resistance region [27], similar to *Salmonella* genomic island 1 (SGI-1) variants [28], but genetic data pointing out the potential to adapt simultaneously to other metal environmental stresses (e.g. copper and silver) is described here first. The chromosomal co-localisation of *pco*+*sil* and *mer*, frequently associated with antibiotic resistance genes, appears to be typical of the European clone. These data, along with the recently reported chromosomal integration of Tn7-like transposons carrying *pco*+*sil* in an *S. enterica* serotype Senftenberg [29], suggests that the accumulation of diverse metal tolerance genetic determinants with other features might relevantly contribute to the evolution or emergence of global widespread emergent clones. Although several studies related the emergence and/or spread of Spanish and Southern European clones to the presence of IncA/C or IncR plasmids harbouring additional MDR features and virulence factors [4,5,26], this study also suggests a role for metal tolerance genes. In fact, *sil*+*merA* and antibiotic resistance genes were linked together on the IncA/C plasmid typical of the second major clone (Spanish clone) circulating

in Europe, contrasting with the less frequent clone (Southern European clone) where the presence of *sil* genes occurred in few isolates associated with IncR MDR plasmids. Metal tolerance genes are dispersed in diverse plasmid backbones (e.g. *merA* in IncA/C or *sil* in IncHI) [11,20,29,30] within different bacterial hosts potentially facing similar chemical challenges, which might favour the sharing and persistence of these genetic elements in particular environments [30]. The ability to acquire diverse genetic platforms with genes coding for a phenotype plethora (e.g. antibiotic resistance, metal tolerance, virulence) might facilitate adaptation of zoonotic clinically relevant strains or epidemic clones such as *Salmonella* 4,[5],12:i:–.

5. Conclusions

In summary, we report a high occurrence of genes encoding tolerance to toxic concentrations of CuSO₄ and AgNO₃ dispersed among the two major *Salmonella* 4,[5],12:i:– MDR clonal lineages (European and Spanish clones) circulating in Europe and causing human infections, which might facilitate the adaptation/expansion of these strains in metal-contaminated environments, particularly copper in anoxic conditions. Furthermore, metal toxic concentrations in the food-animal farm environment can contribute to the persistence of genetic platforms carrying metal and antibiotic resistance genes in this clinically relevant foodborne zoonotic pathogen. More studies are needed to evaluate the impact of the continuous contact with subinhibitory concentrations of several metal environmental contaminants from feed, disinfectants and/or other anthropogenic sources, and their impact on the dynamics of the emergence of different *Salmonella* serotypes and clones.

Acknowledgments

The authors wish to thank to Dr^a Lina Cavaco (Division of Microbiology and Risk Assessment, DTU Food, Technical University of Denmark, National Food Institute, Denmark) for the discussion of agar gel dilution protocols for metal substances. The authors would also like to thank Dr Jill Williams (Department of Genetics of the University of Melbourne, Melbourne, Victoria, Australia) and Drs Christopher P. Randall and Alex J. O'Neill (Antimicrobial Research Centre, School of Molecular and Cellular Biology, University of Leeds, Leeds, UK) who kindly provided *Escherichia coli* ED8739 (pRJ1004) and *E. coli* J53 (pMG101), respectively.

Funding

This work received financial support from the European Union (FEDER funds through the Operational Programme for Competitiveness Factors – COMPETE) and national funds [Foundation for Science and Technology (FCT)] through project Pest-C/EQB/LA0006/2013, and from Investigação Jovem da Universidade do Porto/Santander Totta. J. Mourão was supported by a PhD fellowship from the Foundation for Science and Technology [grant no. SFRH/BD/77518/2011; POPH-QREN]. The authors are greatly indebted to all of the financing sources.

Competing interests

None declared.

Ethical approval

Not required.

References

- [1] European Food Safety Authority (EFSA) Panel on Biological Hazards (BIOHAZ). Scientific opinion on monitoring and assessment of the public health risk of 'Salmonella Typhimurium-like strains'. EFSA J 2010;8:1826.
- [2] Hopkins KL, Kirchner M, Guerra B, Granier SA, Lucarelli C, Porrero MC, et al. Multiresistant *Salmonella enterica* serovar 4,[5],12:i:– in Europe: a new pandemic strain? Euro Surveill 2010;15:19580.
- [3] Hauser E, Tietze E, Helmuth R, Junker E, Blank K, Prager R, et al. Pork contaminated with *Salmonella enterica* serovar 4,[5],12:i:–, an emerging health risk for humans. Appl Environ Microbiol 2010;76:4601–10.
- [4] García P, Hopkins KL, García V, Beutlich J, Mendoza MC, Threlfall J, et al. Diversity of plasmids encoding virulence and resistance functions in *Salmonella enterica* subsp. *enterica* serovar Typhimurium monophasic variant 4,[5],12:i:– strains circulating in Europe. PLoS One 2014;9:e89635.
- [5] Mourão J, Machado J, Novais C, Peixe L, Antunes P. Characterization of the emerging clinically-relevant multidrug-resistant *Salmonella enterica* serotype 4,[5],12:i:– (monophasic variant of *S. Typhimurium*) clones. Eur J Clin Microbiol Infect Dis 2014;33:2249–57.
- [6] Scientific Committee on Emerging and Newly Identified Health Risks (SCENIHR). Assessment of the antibiotic resistance effects of biocides; January 2009. http://ec.europa.eu/health/ph_risk/risk_en.htm [accessed 13.07.14].
- [7] Seiler C, Berendonk TU. Heavy metal driven co-selection of antibiotic resistance in soil and water bodies impacted by agriculture and aquaculture. Front Microbiol 2012;3:399.
- [8] Baker-Austin C, Wright MS, Stepanauskas R, McArthur JV. Co-selection of antibiotic and metal resistance. Trends Microbiol 2006;14:176–82.
- [9] Williams JR, Morgan AG, Rouch DA, Brown NL, Lee BT. Copper-resistant enteric bacteria from United Kingdom and Australian piggeries. Appl Environ Microbiol 1993;59:2531–7.
- [10] Liebert CA, Wireman J, Smith T, Summers AO. Phylogeny of mercury resistance (*mer*) operons of Gram-negative bacteria isolated from the fecal flora of primates. Appl Environ Microbiol 1997;63:1066–76.
- [11] García Fernández A, Cloeckaert A, Bertini A, Praud K, Doublet B, Weill FX, et al. Comparative analysis of IncHI2 plasmids carrying *bla*_{CTX-M-2} or *bla*_{CTX-M-9} from *Escherichia coli* and *Salmonella enterica* strains isolated from poultry and humans. Antimicrob Agents Chemother 2007;51:4177–80.
- [12] Woods EJ, Cochrane CA, Percival SL. Prevalence of silver resistance genes in bacteria isolated from human and horse wounds. Vet Microbiol 2009;138:325–9.
- [13] Sütterlin S, Tano E, Bergsten A, Tallberg AB, Melhus A. Effects of silver-based wound dressings on the bacterial flora in chronic leg ulcers and its susceptibility in vitro to silver. Acta Derm Venereol 2012;92:34–9.
- [14] Kremer AN, Hoffmann H. Subtractive hybridization yields a silver resistance determinant unique to nosocomial pathogens in the *Enterobacter cloacae* complex. J Clin Microbiol 2012;50:3249–57.
- [15] Medardus JJ, Molla BZ, Nicol M, Morrow WM, Rajala-Schultz PJ, Kazwala R, et al. In-feed use of heavy metal micronutrients in U.S. swine production systems and its role in persistence of multidrug-resistant salmonellae. Appl Environ Microbiol 2014;80:2317–25.
- [16] Zimmermann M, Udagedara SR, Sze CM, Ryan TM, Howlett GJ, Xiao Z, et al. PcoE—a metal sponge expressed to the periplasm of copper resistance *Escherichia coli*. Implication of its function role in copper resistance. J Inorg Biochem 2012;115:186–97.
- [17] Hasman H, Kempf I, Chidaïne B, Cariolet R, Erbsoll AK, Houe H, et al. Copper resistance in *Enterococcus faecium*, mediated by the *tcrB* gene, is selected by supplementation of pig feed with copper sulfate. Appl Environ Microbiol 2006;72:5784–9.
- [18] Silver S. Bacterial silver resistance: molecular biology and uses and misuses of silver compounds. FEMS Microbiol Rev 2003;27:341–53.
- [19] Cavaco LM, Hasman H, Aarestrup FM. Zinc resistance of *Staphylococcus aureus* of animal origin is strongly associated with methicillin resistance. Vet Microbiol 2011;150:344–8.
- [20] Gupta A, Phung LT, Taylor DE, Silver S. Diversity of silver resistance genes in IncH incompatibility group plasmids. Microbiology 2001;147:3393–402.
- [21] Randall CP, Oyama LB, Bostock JM, Chopra I, O'Neill AJ. The silver cation (Ag⁺): antistaphylococcal activity, mode of action and resistance studies. J Antimicrob Chemother 2013;68:131–8.
- [22] Mijndonckx K, Leys N, Mahillon J, Silver S, Van Houdt R. Antimicrobial silver: uses, toxicity and potential for resistance. Biometals 2013;26:609–21.
- [23] Maillard JY, Hartemann P. Silver as an antimicrobial: facts and gaps in knowledge. Crit Rev Microbiol 2013;39:373–83.
- [24] Porcheron G, Garénaux A, Proulx J, Sabri M, Dozois CM. Iron, copper, zinc, and manganese transport and regulation in pathogenic Enterobacteria: correlations between strains, site of infection and the relative importance of the different metal transport systems for virulence. Front Cell Infect Microbiol 2013;3:90.
- [25] Hölzel CS, Müller C, Harms KS, Mikolajewski S, Schäfer S, Schwaiger K, et al. Heavy metals in liquid pig manure in light of bacterial antimicrobial resistance. Environ Res 2012;113:21–7.
- [26] Antunes P, Mourão J, Pestana N, Peixe L. Leakage of emerging clinically relevant multidrug-resistant *Salmonella* clones from pig farms. J Antimicrob Chemother 2011;66:2028–32.

- [27] Lucarelli C, Dionisi AM, Filetici E, Owczarek S, Luzzi J, Villa L. Nucleotide sequence of the chromosomal region conferring multidrug resistance (R-type ASSuT) in *Salmonella* Typhimurium and monophasic *Salmonella* Typhimurium strains. *J Antimicrob Chemother* 2012;67:111–4.
- [28] Doublet B, Praud K, Weill FX, Cloeckaert A. Association of IS26-composite transposons and complex *Int*-type integrons generates novel multidrug resistance loci in *Salmonella* genomic island 1. *J Antimicrob Chemother* 2009;63:282–9.
- [29] Moreno Switt AI, den Bakker HC, Cummings CA, Rodriguez-Rivera LD, Govoni G, Raneiri ML, et al. Identification and characterization of novel *Salmonella* mobile elements involved in the dissemination of genes linked to virulence and transmission. *PLoS One* 2012;7:e41247.
- [30] Lindsey RL, Frye JG, Fedorka-Cray PJ, Meinersmann RJ. Microarray-based analysis of *IncA/C* plasmid-associated genes from multidrug-resistant *Salmonella enterica*. *Appl Environ Microbiol* 2011;77:6991–9.

Supplementary Table S1
Primers used in the PCR to search for acquired metal tolerance genes

Target gene	Gene coding for	Primer	Primer sequence (5'-3')	Amplicon size (bp)	Annealing temperature (°C)	Nucleotide position (GenBank accession no.)	Reference
<i>pcoA</i>	Multicopper oxidase	pcoA_978_F pcoA_1481_R	CTCGGGGATGTCAGTGGCTACACCT ATCCGGAAGGTCAGACACCGTCCATAGAC	504	60	978–1002 1481–1454 (X83541.1)	This study
<i>pcoD</i>	Copper inner membrane pump	pcoD_F pcoD_R	CTGGCCACACTTGCCCTGGGG CACGCTACGGCGCCCAAGAT	500	55	3801–3820 4300–4281 (X83541.1)	This study
<i>tcrB</i>	Copper export ATPase	tcrB_F tcrB_R	CATCACGGTAGCTTTAAGGAGATTTC ATAGAGGACTCCGCCACCATTG	663	66	4954–4980 5616–5595 (AY048044.2)	[17]
<i>silA</i>	Silver inner-membrane proton/cation antiporter	silA_Fw silA_Rv	GCAAGACCGGTAAAGCAGAG CCTGCCAGTACAGGAACCAT	936	59	8936–8955 9871–9852 (AF067954.1)	This study
<i>silE</i>	Silver/copper periplasmic binding protein	sile_1105_F sile_1368_R	GTTCGTCATGGTYTCATGAGC GTACTYCCCCGGACATCACTAATT	264	62	1105–1125 1368–1345 (AF067954.1)	This study
<i>merA</i>	Mercuric reductase	merA_1F merA_5R	ACCATCGGGCGCACCTGCGT ACCATCGTCAGGTAGGGGAAC	1238	65	2140–2159 3377–3357 (K03089.1)	[10]
<i>arsB</i>	Arsenite transmembrane pump	arsB_Fw arsB_Rv	AGTGAAAGACAGACGAAAGCG GGCAGATAGTGTGGAATGCG	1136	60	159735–159754 160870–160851 (BX664015.1)	[11]
<i>terF</i>	Tellurite resistance protein	terF_Fw terF_Rv	ATGCAGGCTCAAGGAATCGC TTTCATCGATCCACGGTCTG	894	60	80270–80289 81163–81145 (BX664015.1)	[11]

Tolerance to multiple metal stressors in emerging non-typhoidal MDR *Salmonella* serotypes: a relevant role for copper in anaerobic conditions

Joana Mourão¹, Sara Marçal², Paula Ramos², Joana Campos¹, Jorge Machado³, Luísa Peixe¹, Carla Novais¹ and Patrícia Antunes^{1,2*}

¹ UCIBIO/REQUIMTE, Departamento de Ciências Biológicas, Laboratório de Microbiologia, Faculdade de Farmácia, Universidade do Porto, Porto, Portugal; ² Faculdade de Ciências da Nutrição e Alimentação, Universidade do Porto, Porto, Portugal; ³ Laboratório Nacional de Referência de Infecções Gastrointestinais, Departamento de Doenças Infecciosas, Instituto Nacional de Saúde Dr. Ricardo Jorge, Lisboa, Portugal.

Running title: Tolerance to metal stressors in MDR *Salmonella*

Keywords: Anaerobic, biocides, copper, metals, multidrug resistant, non-typhoidal *Salmonella*, silver

***Corresponding author:**

Patrícia Antunes, Ph.D.

Faculdade de Ciências da Nutrição e Alimentação, Universidade do Porto.

Rua Dr. Roberto Frias, 4200-465 Porto, Portugal

E-mail address: patriciaantunes@fcna.up.pt

Tel.: +351 22 507 4320

Fax: +351 22 507 4329

Submitted to the Journal of Antimicrobial Chemotherapy

Under the Second Editorial Revision

SYNOPSIS

Objectives: The driving factors for the expansion of particular MDR *Salmonella* serotypes/clones are not completely understood. Here we assessed if emergent MDR *Salmonella* serotypes/clones were more enriched in metal tolerance genes (e.g. to Cu/Ag) than other less frequent ones, as an additional survival adaptive feature to environments contaminated with metals.

Methods: Metal (Cu-*pco*/Ag, Cu-*sil*/Hg-*mer*/As-*ars*/Te-*ter*) tolerance genes screening (PCR/sequencing), MICs to CuSO₄/AgNO₃ (aerobiosis/anaerobiosis), genetic elements characterisation (S1/I-Ceul PFGE) and conjugation assays were performed in a well-characterised *Salmonella* collection (n=275 isolates; 2000-2014; 49 serotypes/clones).

Results: The *sil±pco* genes were detected in 37% of isolates from diverse serotypes, mainly in emergent Rissen/ST469 and Typhimurium/ST34 European clone (100%) which are mostly associated with pig-setting where copper is highly used. These genes were frequently co-located with *merA±terF* and/or antibiotic resistance genes in plasmids (100-270kb; IncHI2/IncHI1/IncN/IncFIIA; mostly transferable by conjugation) or in chromosome. Most *sil±pco*⁺ isolates (77%) were MDR contrasting with *sil±pco*⁻ ones (48%). The *sil±pco*⁺ isolates presented significantly higher MIC_{CuSO₄} under anaerobiosis (MIC₅₀/MIC₉₀=28/32mM) and MIC_{AgNO₃} after previous Ag contact (MIC₅₀/MIC₉₀>3mM) than *sil*⁻ ones (MIC₅₀/MIC₉₀=2/8mM to CuSO₄; MIC₅₀/MIC₉₀=0.125/0.16 to AgNO₃). The use of these modified methodological approaches allowed the establishment of tolerance cut-offs to CuSO₄/AgNO₃, here firstly proposed.

Conclusions: This study demonstrates that Cu/Ag tolerance genes acquisition might account for the emergence of particular clinically relevant MDR *Salmonella* serotypes/clones by facilitating their adaptation to diverse metal contaminated settings, especially to pig production. An assessment of control measures for the use and/or accumulation of metals in diverse environments is needed to prevent a wider expansion of such strains or the emergence of new ones.

INTRODUCTION

Non-typhoidal *Salmonella enterica* is a worldwide major foodborne pathogen with a main reservoir in the intestinal tract of food-producing animals, from which it is readily transmitted to farm surrounding environments and the food chain.^{1,2} In recent years, expansion of particularly successful *Salmonella* serotypes and clonal lineages have been reported globally. They are scattered to humans mostly by a wide range of foodstuffs of animal origin, they are involved in endemic/epidemic situations and they frequently show resistance to multiple antibiotics limiting therapeutic options when necessary.¹⁻⁵

The exact mechanisms of successful *Salmonella* serotypes persistence and spread among animals and in the environment still remain largely unknown. However, recent studies pointed out tolerance to antimicrobial non-antibiotic compounds widely used in food-producing animals as a potential adaptive advantage in environments and hosts (animal/human) contaminated with these type of compounds.^{4,6} Among them are heavy metals, currently associated with diverse anthropogenic activities as they are incorporated in biocidal products, such as disinfectants, antiseptics/medicines or preservatives (e.g. silver-Ag; copper-Cu), in feed additives for animal growth promotion (e.g. copper, zinc) or in soil fertilisers (e.g. arsenic).⁷⁻¹¹ Consequently, metals as environmental pollutants can accumulate for example in manure, waste lagoons or feed in the animal production setting,^{9,10,12} and then be spread by sewage sludge and liquid manure applied in amended soils worldwide or by water bodies.^{9,13-15} This extensive environmental metal contamination is a topic of increasing concern recognized by diverse authorities^{8,10,16,17} as they can contribute for the selection of metal tolerant and/or antibiotic resistant bacteria by mechanisms of cross or co-resistance.^{9,11,18} Metals can also potentially participate as a catalyst of gene transfer among diverse microorganisms sharing the same ecologic niche.^{9,17} A recent study from our group showed that *S. 4,[5],12:i:-*, a worldwide emergent MDR serotype, carries diverse metal tolerance genes to handle with metal toxicity, such as those encoding for Cu/Ag efflux (*sil/pco* genes) and mercury-Hg enzymatic reduction (*mer* genes).⁶ We also showed that the *sil* gene cluster was frequent among the two major *S. 4,[5],12:i:-* clones (European/Spanish) and that its presence was particular relevant for Cu tolerance under anaerobic conditions,⁶ as suggested by others.^{19,20} Nevertheless, a comprehensive analysis of acquired metal tolerance by additional *Salmonella* serotypes remains scarce, including for other emergent ones.

Here we hypothesise that emergent antibiotic resistant *Salmonella* serotypes/clones are more enriched in metal tolerance genes (e.g. to Cu and Ag) than other less frequent ones, and that this is an additional survival adaptive feature

contributing to their emergence in metal contaminated environments. To test this hypothesis, we used a large Portuguese *Salmonella* collection representative of different serotypes/clones, in order to search and characterise diverse metal tolerance genes as well as their associated genetic platforms and tolerance phenotype.

MATERIALS AND METHODS

Bacterial isolates and epidemiological background

From a collection of 2500 Portuguese non-typhoidal *Salmonella enterica*, 275 isolates belonging to 49 serotypes were selected based on source, antibiotic susceptibility phenotypes and genotypes (MDR was considered when resistance to ≥ 3 antibiotics of different families was observed), PFGE types and/or Sequence types (ST) obtained by MLST (Table 1 and Table S1, available as Supplementary data at JAC Online).^{1,21-25} They were collected between 2000-2014 in several geographical regions of Portugal (North, Centre, South, Islands) from human clinical cases [biological products from hospitals (n=108; articular liquid, blood, faeces, peritoneal fluid, pus or urine) and community laboratories (n=11; faeces)], food products (n=89; beef, cow, pork, poultry, quail, cooked meal), food-animal production setting (n=50-pigs/piggeries environment; n=4-aquacultures environment) and the aquatic environment (n=13; drinking, bathing or river water). These isolates belonged to the two most frequent serotypes worldwide [*Salmonella* Enteritidis (n=32/ST11) and *Salmonella* Typhimurium (n=96/ST19 and ST34)], but also to others that are emergent and have clinical relevance [e.g. *Salmonella* Rissen (n=50/ST469), *Salmonella* Derby (n=16), *Salmonella* Hadar (n=4), *Salmonella* Heidelberg (n=2), *Salmonella* Infantis (n=3), *Salmonella* Newport (n=1), *Salmonella* Saintpaul (n=3), *Salmonella* Stanley (n=2), *Salmonella* Virchow (n=4)]. Thirty-eight less frequently serotypes detected in surveillance studies were also included (Table 1 and Table S1, available as Supplementary data at JAC Online).

Search of acquired metal tolerance genes and copper/silver susceptibility

The occurrence of diverse acquired metal tolerance genes was assessed in all isolates (n=275) and was performed by PCR using conditions and primers previously described for the S. 4,[5],12:i:- collection.⁶ The searched genes encoded different metal tolerance mechanisms (e.g. export, reduction) related to copper, silver, mercury, arsenite and tellurite: two genes of *pcoABCDRSE* cluster, the *pcoA* (multicopper oxidase) and *pcoD* (copper inner membrane pump); two genes of *silCFBAPRSE* cluster, the *silA* (silver inner-membrane proton/cation antiporter) and *silE* (silver/copper periplasmic metal binding protein); *arsB* (arsenite transmembrane pump); *merA* (mercuric reductase) and

terF (tellurite resistance protein) (Table S2, available as Supplementary data at JAC Online). Representative isolates selected based on presence (n=38) or absence (n=18) of *silA-silE* or *pcoA-pcoD* genes, anaerobic MIC_{CuSO₄} values, serotype, PFGE-type and antibiotic resistance profiles, were further characterised for the presence or absence of *silCPRS* and/or *pcoRSE* gene clusters. Primers and conditions were described in Table S2, available as Supplementary data at JAC Online.

Representative isolates (n=166) from all serotypes and different clonal lineages, PFGE-types, sources, isolation date and antibiotic resistance phenotypes and genotypes, harbouring or lacking acquired copper/silver tolerance genes, were selected for phenotypic assays. MICs were determined in aerobic and anaerobic (GENbox jar with GENbox anaer. and an anaerobic indicator; bioMérieux) atmospheres by the agar dilution method on Mueller-Hinton II agar plates (bioMérieux) supplemented with CuSO₄ (0.25, 0.5, 1, 2, 4, 8, 12, 16, 20, 24, 28, 32, 36 mM; adjusted to pH=7.2) or AgNO₃ (0.025, 0.06, 0.08, 0.125, 0.16, 0.25, 0.32, 0.5, 0.75, 1.0, 1.5, 3 mM; adjusted to pH=7.4), as described.⁶ MICs to AgNO₃ were also determined after overnight incubation of *sil*⁺ and *sil*⁻ cultures (14-16h) in Mueller-Hinton II broth (BBL, BD, Franklin Lakes, NJ, USA) supplemented with 0.025 mM of this metal for induction of *silCFBAPRSE* cluster.⁶ Two plates (the first and last to be inoculated) of Mueller-Hinton II agar without CuSO₄ or AgNO₃ were included in all MIC assays to assess growth ability of every isolates in different atmospheres. *Enterococcus faecium* BM4105RF²⁶ or *Staphylococcus aureus* ATCC 29213²⁷ (both negative for all genes tested) and *Escherichia coli* ED8739²⁸ (plasmid pRJ1004 with *pco* genes) or *E. coli* J53²⁷ (plasmid pMG101 with *sil* genes) were used as controls in CuSO₄ or AgNO₃ tolerance assays.

Characterisation of the genetic platforms carrying metal tolerance genes

Characterisation of genetic platforms (genomic location and plasmid analysis) was carried out in isolates positive for metal tolerance genes, selected accordingly to the serotype and/or PFGE-type frequency. Plasmid analysis was based on identification of incompatibility groups by PCR-based replicon typing (PBRT) method^{29,30} and sequencing. The genetic location of metal and antibiotic resistance determinants in plasmids or chromosome was accomplished by Southern blot hybridisation of S1 nuclease (Takara Bio Inc., Shiga, Japan) and I-Ceul (New England BioLabs, Ipswich, MA, USA) digested genomic DNA with intragenic metal tolerance (*pcoD*, *silA*, *merA*, *arsB*, *terF*) and antibiotic resistance genes, *intl1*, plasmid incompatibility groups and 16S rRNA probes, by standard methods.⁶ In order to analyse the dispersion of *sil* and *pco* metal tolerance genes and their genetic context in *Salmonella* from other

serotypes and diverse epidemiological scenarios, a detailed GenBank *in silico* analysis in available genomes until October 2015 was also performed (Table S3, available as Supplementary data at JAC Online).

Conjugation assays

Transfer of genetic elements carrying metal tolerance and/or antibiotic resistance genes was assessed by standard conjugation assays.⁶ Transfer of copper/silver tolerance was analysed for isolates carrying *sil* and/or *pco* or lacking known genes, but exhibiting high MIC values to CuSO₄ (≥ 24 mM) in anaerobiosis. It was performed by filter-mating method with log phase growth cells at 1:1 donor:recipient ratio, at 37°C to known recipient strains (*Escherichia coli* K802N and *S. Typhimurium* LSP 448/02, with antibiotic and metal susceptibility shown in Table 4). Transconjugants obtained from isolates with *sil*±*pco* genes were recovered after incubation at 37°C/24h from Mueller-Hinton II agar plates supplemented with nalidixic acid (64 mg/L) plus other antibiotics [ampicillin (16 mg/L), streptomycin (32 mg/L), sulfamethoxazole (256 mg/L) or trimethoprim (8 mg/L)] in aerobic conditions. Transconjugants obtained from isolates without *sil*±*pco* genes but with high MIC_{CuSO₄} in anaerobic conditions were recovered in Mueller-Hinton II agar plates supplemented with CuSO₄ (8 mM or 12 mM, according to the recipient strain used). Confirmation of transconjugants, tolerance to Cu/Ag and co-transfer of antibiotic resistance was confirmed at phenotypic and genotypic levels as previously described.^{5,6}

RESULTS

Acquired metal tolerance genes were dispersed in diverse *Salmonella* serotypes, especially clinical relevant MDR pig-associated clones

The Cu and Ag tolerance genes *silA-silE*±*pcoA-pcoD* were detected in 37% (n=101/275) of *Salmonella* isolates from diverse sources, serotypes (n=9/49; 18%) and clones (18 PFGE-types and 10 sequence types-ST), including recently emerging ones. Most of them were identified as *S. Typhimurium* “European clone” (ST34) and *S. Rissen* (ST469) (n=80/80-100% *versus* n=21/195-11% from other serotypes/clones). They were often recovered from samples of pig origin and were MDR (n=78/101; 77%) (Table 1). The simultaneous presence of *sil* and *pco* genes was frequently found (95%; n=96/101) and always associated with the complete *sil*+*pco* gene clusters. Combinations of *sil*+*pco*+*merA*±*terF* (40%; n=40/101) were frequently detected among diverse MDR serotypes/clones (6 serotypes/6 STs; 12 PFGE-types). In contrast, single carriage of *sil* (n=4/101) or *sil*+*merA* (n=1/101) genes was detected in only 5% of the

isolates, corresponding to the *S. Typhimurium* DT104 epidemic clone (ST19) with the typical ACSSuT (A-amoxicillin, C-chloramphenicol, S-streptomycin, Su-sulfamethoxazole and T-tetracycline) antibiotic resistance profile (related with SGI-1) apart from trimethoprim resistance (Table 1).

Salmonella (63%, n=174/275) without *sil±pco* belonged to 46 serotypes, including all the isolates from several poultry/eggs-associated serotypes such as *S. Enteritidis*, *S. Hadar*, *S. Virchow*, *S. Infantis*, *S. Saintpaul*, *S. Stanley* or *S. Heidelberg*. The analysis of antibiotic resistance profiles revealed that these isolates without *sil±pco* were frequently less MDR (n=84/174; 48%) than *sil±pco*⁺ (Table S1, available as Supplementary data at JAC Online). Nonetheless, other acquired metal tolerance genes such as *merA* (26%; n=45/174) and/or *arsB+terF* (1%; n=1/174) were found in 26% (n=46/174) of these isolates, which belonged to 10 MDR serotypes (Table S1, available as Supplementary data at JAC Online).

Higher tolerance to *CuSO₄* in anaerobic atmosphere and to *AgNO₃* after prior exposure to silver was observed among the *sil+pco*⁺ or *sil*⁺ isolates

All *sil+pco*⁺ or *sil*⁺ isolates presented higher MIC_{CuSO₄} values comparing with those without these genes (MIC₅₀/MIC₉₀=28/32 mM versus 2/8 mM, respectively) in anaerobic condition tests. The MICs_{CuSO₄} in aerobiosis were higher (MIC₅₀/MIC₉₀=32/36 mM) than in anaerobiosis, with identical MICs for isolates with or without *sil±pco* genes (Table 2). Genotypic assays also confirmed the occurrence of complete *sil+pco* gene clusters in isolates with MIC_{CuSO₄}>16mM and its absence when the MICs were lower under anaerobiosis. The only exceptions were 2 isolates (*S. Essen* and *S. Il 4,5:b:-*) lacking the known *sil+pco* gene clusters, but exhibiting high MIC_{CuSO₄} values (24 and 28 mM) in anaerobic atmosphere (Table 1).

Regarding *AgNO₃*, differences between *sil*⁺ (MIC₅₀/MIC₉₀=0.25 mM in aerobiosis and 0.32/0.5 mM in anaerobiosis) and *sil*⁻ (MIC₅₀/MIC₉₀=0.125/0.16 mM in aerobiosis and 0.16/0.32 mM in anaerobiosis) isolates were observed in both atmospheres, although a significant increment was detected for *sil*⁺ isolates only after prior contact with Ag (MIC₅₀/MIC₉₀>3mM in *sil*⁺ isolates versus 0.125/0.16 mM in *sil*⁻ isolates). One exception was one *Salmonella* isolate (*S. Give*) with previous contact with *AgNO₃* and with a MIC that only increased up to 0.5 mM (Table 3).

Metal tolerance genes were located in diverse MDR genetic platforms

Among the characterised isolates, selected accordingly to the serotype and/or PFGE-type frequency, diverse genetic contexts for *sil±pco* genes were detected (Table 1).

Isolates of *S. Typhimurium* ("European clone"), *S. Rissen*, *S. Anatum*, *S. Brandenburg*, *S. Bredeney*, *S. Give* and *S. Senftenberg*, mostly resistant from 2 to 10 antibiotics, carried *sil+pco±merA±terF* genes in the chromosome. Additionally, these metal tolerance genes were often co-located in the same chromosomal I-Ceul-PFGE fragment with diverse antibiotic resistant genes in the emergent *S. Typhimurium* "European clone" [ASSuT markers-*bla*_{TEM-1}, *strA-strB*, *tet(B)* and/or *sul2*] as well as in other serotypes [e.g. *tet(A)*, *sul1*] (Table 1). In contrast, in isolates from 4 serotypes (*S. Brandenburg*, *S. Bredeney*, *S. Muenchen* and *S. Wien*) the *sil+pco±merA±terF* genes were plasmid located (190-270kb; IncHI1-n=1, IncHI2-n=2 or untypeable-n=1) along with variable combinations of typical and/or atypical *sul3*-type class 1 integrons plus other antibiotic resistance genes [e.g. *bla*_{TEM-1}, *tet(A)*, *tet(B)*]. In *S. Typhimurium* DT104 MDR clone carrying only *sil±merA* genes they were found co-located in IncFIIA (n=1) or IncN (n=3) plasmids (100-140 kb) with typical and/or atypical *sul3*-type class 1 integrons²³ plus other antibiotic resistance genes [e.g. *bla*_{TEM-1}, *tet(A)*].

Isolates of diverse clinically relevant MDR *Salmonella* serotypes without *sil±pco* carried other metal tolerance genes (*merA* and/or *arsB+terF*). They were co-located with diverse antibiotic resistance genes (several associated with class 1 integrons) either in the chromosome or in plasmids of diverse families (80-290 kb; IncFIIIs-n=5, IncI1-n=1, IncHI2-n=1, IncP-n=4, untypeable-n=2) (Table S1, available as Supplementary data at JAC Online).

Copper and silver tolerance was co-transferred along with antibiotic resistance

The conjugative transfer of plasmid located *sil+pco* (n=3/4-IncHI1, IncHI2 or untypeable; one IncHI2 not transferred) and *sil* (n=1/4-IncFIIA; three IncN not transferred) genes was achieved for 50% of the *Salmonella* strains studied (n=4/8; *S. Typhimurium*, *S. Bredeney*, *S. Muenchen* and *S. Wien*), using both *E. coli* K802N and *S. Typhimurium* LSP 448/02 recipient strains (Table 4). Acquisition of *sil+pco* genes conferred higher MIC_{CuSO₄} (20-28 mM) to transconjugants than the recipient strains *E. coli* K802N (1-2 mM) and *S. Typhimurium* LSP 448/02 (8-12 mM) in anaerobic atmosphere. Regarding MIC_{AgNO₃}, *S. Typhimurium* LSP 448/02 transconjugants also presented a high MIC_{AgNO₃} (>3 mM) after previous contact with Ag. Both *sil+pco* and antibiotic resistance genes (*sul*, *dfrA*, *aadA*, *tet*) were co-located in the same plasmid belonging to diverse families (IncFIIA, IncHI1, IncHI2 or untypeable plasmids) as in the donor strains (Table 4). None of the *sil+pco* genes located in the chromosome (n=5 tested) were transferable by conjugation assays, as well as an unknown Cu tolerance

mechanism observed in 2 *Salmonella* isolates (*S.* Essen and *S.* Il 4,5:b:-) presenting high MIC_{CuSO₄} in anaerobic conditions.

DISCUSSION

Using a large and extensively characterised *Salmonella* collection we address the potential role of metal tolerance, namely to Cu and Ag, for the emergence of particular *Salmonella* serotypes/clones associated with human infections and spread through the food chain. We also discussed the potential impact of this tolerance in the survival of such serotypes/clones in specific environments (e.g. contaminated with Cu under anaerobic conditions) as well as the diversity of genetic platforms implicated in the spread of metal tolerance genes along with antibiotic resistance among *Salmonella*.

In this study, we demonstrate that the Cu/Ag tolerance genes *sil±pco* were dispersed among diverse *Salmonella* serotypes and sources, mostly MDR, with emphasis to the clinically relevant emergent *S.* Rissen/ST469 and *S.* Typhimurium/ST34 “European clone”. These two cases along with the emergent *S.* 4,[5],12:i:- MDR European and Spanish clones, previously described as often carrying *sil±pco* genes,⁶ are strongly associated with pig production worldwide.^{1,3,4,31} The higher European Cu concentrations allowed in feed for pigs (e.g. piglets 175 mg/kg, ~2.8 mM) than for other animals,⁷ suggest the occurrence of a high selective pressure in the pig production setting. This selective pressure can potentially contribute to the emergence of such strains in pigs and piggeries related settings where Cu can rapidly accumulate, even at higher concentrations (e.g. up to 3387.6 mg/kg, 53 mM in pig manure and consequently pig amended soils).^{14-15,32-34} In contrast, *S.* Enteritidis, the most reported serotype with poultry/eggs production as the main source³⁵, showed absence of *sil±pco* genes in our study and Genbank genome analysis (Table S3, available as Supplementary data at JAC Online). These data could be partial explained by the low Cu concentrations (e.g. recommendation for species or animal categories other than piglets is 35 mg/Kg, ~0.6 mM) allowed in this setting in Europe,⁷ and/or be associated with particular features of this serotype impairing the acquisition and/or fixation of specific genetic platforms as those carrying *sil±pco*. However, the available genomes related to emerging poultry-related *Salmonella* serotypes in the USA (e.g. *S.* Heidelberg, *S.* Kentucky) carry *sil+pco* gene clusters (Table S3, available as Supplementary data at JAC Online).³⁶ This data alerts for the potential relevance of such genes in the survival and emergence of diverse *Salmonella* serotypes under different animal production practices (e.g. in the USA copper supplements are not restricted and the typical amounts range from 125-250 ppm, ~2-4 mM, in most broiler diets) or environmental factors.³⁷

A common location of *sil±pco* on the chromosome was observed in diverse *Salmonella* serotypes from our collection, especially in the emergent MDR *S. Rissen*/ST469 and in the European clone of *S. Typhimurium*/ST34. This is in agreement with data previously reported for the major clone of *S. 4,[5],12:i:-* (European clone).⁶ In other *Salmonella* serotypes/clones, *sil±pco* genes were located on large, mostly transferable, plasmids (100-270kb) of different major families (IncHI2, IncHI1, IncN, IncFIIA). These data point out to *sil±pco* widespread among diverse genetic platforms abundant in local metagenomes and described to be spread in *Enterobacteriaceae*.³⁸ In *Salmonella*, IncHI2 plasmids have been frequently associated with diverse metal tolerance genes as *sil+pco*,^{36,39} but we highlight the relevance of IncFIIA and IncN plasmids in the dispersion of metal encoding genes, namely in the epidemic *S. Typhimurium* DT104 clone.

A more detailed genomic study (our isolates, Genbank genome analysis and literature available) showed that *sil* and *pco* gene clusters are often adjacent in *Salmonella* isolates from different sources and serotypes/clonal lineages, particularly in recently emerging ones (Table 1 and Table S3, available as Supplementary data at JAC Online).^{6,19,40,41} The *in silico* analysis of recent available genome sequences pointed out, in most isolates, for a common module containing *sil+pco* gene clusters and genes coding for a hypothetical protein and a total/partial endonuclease (Table S3, available as Supplementary data at JAC Online). Such genes were described as part of a Tn7-like transposon previously identified in *Salmonella*, the Tn6230.^{19,40,42} As showed in Table S3, available as Supplementary data at JAC Online, the complete Tn6230 (common module with the typical Tn6230 proteins TnsABC, involved in transposition of the genetic element) was present in most *sil+pco*⁺ *Salmonella* isolates from diverse serotypes. Most of these Tn6230 elements were near the *yhiN* gene, previously described to be the common insertion site of this transposon.^{40,42} In contrast, when the TnsABC proteins were absent, the common module was usually located near a tRNA-Phe region or flanked by diverse insertion sequences (Table S3, available as Supplementary data at JAC Online). The absence of the TnsABC proteins could be related to recombination events, potentially associated with the fixation of the element in diverse *Salmonella* serotypes, namely in *S. 4,[5],12:i:-* and *S. Typhimurium* (Table S3, available as Supplementary data at JAC Online). These data, along with the chromosomal location of *sil+pco* genes in the emergent *S. Rissen*/ST469, *S. Typhimurium*/ST34 European clone and *S. 4,[5],12:i:-* European clone⁶ from our collection, stresses the need of further studies to elucidate the contribution of these Tn6230-like elements alone, or eventually as part of potential genomic islands often related to Tn7-like transposons,⁴² for the emergence of particular clones/serotypes.

Beside Cu and Ag tolerance genes (*sil±pco*) we frequently co-located other metal tolerance (e.g. *merA*) and antibiotic resistance genes (e.g. to tetracycline-*tetA*, *tetB*; sulphonamides-*sul1* and *sul3* class 1 integron related or *sul2*) in the same genetic platforms (plasmids or chromosome) of our isolates. In the case of the widespread European clone of *S. Typhimurium*, the chromosomal co-location of *sil+pco*, *merA* and ASSuT resistance region appears to be typical of these strains, as occurred with the European clone of *S. 4,[5],12:i:-*,⁶ features that may have relevantly contributed to its evolution and emergence. Most of these antibiotic resistance and *merA* genes are described to be widespread among diverse bacterial hosts, genetic platforms and environments,^{1,22,43,44} independently of *sil+pco* occurrence. This is also supported by our results showing several *Salmonella* serotypes carrying *sul*, *tet* and *merA* genes in diverse major plasmid families (IncP, IncFIIIs, IncI1) without *sil+pco* genes. This abundance might facilitate their involvement in different recombination events with other metal encoding genetic elements (e.g. carrying Tn6230-like) and the maintenance of MDR isolates by the use of different compounds (e.g. antibiotics or metals). Recent environmental studies in metal contaminated soils/animal manure, particularly with Cu due to anthropogenic-derived sources (e.g. animal feed additives, agriculture microbicides), have shown that metal availability, even at low level concentrations, was correlated with a high prevalence of antibiotic resistance in diverse bacteria.^{32-34,45,46} Also, an *in vitro* assay showed that the presence of sublethal levels of Cu or As were sufficient to co-select for the maintenance of MDR plasmids.⁴⁷

Independently of plasmid and chromosomal location, the occurrence of *sil* efflux system was associated with a tolerance phenotype to CuSO₄ and AgNO₃ in wild type isolates (diverse *Salmonella* serotypes/clones, sources and antibiotic resistance profiles) and their transconjugants, pointing out to an acquired adaptive feature of these strains to diverse environmental challenges, beside antibiotic resistance. However, particular methodological conditions are mandatory to be applied in order to distinguish isolates with and without *sil* genes. In the case of CuSO₄, anaerobic conditions during phenotypic assays are critical, as under aerobiosis all isolates showed MIC≥20 mM (this study)^{4,6,48} impairing the establishment of breakpoints values to distinguish isolates that have acquired Cu tolerance genes, namely *sil* or *sil-pco* gene clusters. Considering our results under anaerobiosis on an extensive collection of *Salmonella* serotypes (this study),⁶ here we proposed a CuSO₄ tolerance cut-off>16 mM. However, as we did not identify any isolate with a MIC=16 mM, the inclusion of this value in the susceptible or tolerance group is not possible and we recommend the *sil* genes identification in future cases with MIC=16 mM. This approach under anaerobiosis was also critical in our previous study to differentiate *Enterococcus* spp.

with/without the Cu tolerance *tcrYAZB* cluster, which was not accurate under aerobic conditions.⁴⁹ In the case of AgNO₃ susceptibility assays, for the detection of a clear tolerance phenotype associated with the occurrence of *sil* efflux system, a prior contact of the isolates with subinhibitory concentrations of AgNO₃ is critical, as previously reported for *S. 4,[5],12:i:-*.⁶ Using this approach, an AgNO₃ tolerance cut-off value >3 mM is proposed.

The increase tolerance of *Salmonella* isolates to CuSO₄ and AgNO₃ implies a better awareness of the potential role of these metal contaminants¹³⁻¹⁵ for an improved survival, selection and competition of *sil*+ MDR emergent serotypes/clones than *sil*-ones in diverse environments (e.g. across the food chain), especially when under specific conditions.^{4,8,9} For example, *Salmonella* isolates carrying *sil* cluster might overcome the highly toxic cuprous form Cu⁺ abundant in anaerobic environments (e.g. gut of livestock carriers, manure, waste lagoons or forage/feed, sewage and sludge)^{19,20,50} or silver salts/nanoparticles in diverse food production contexts (e.g. disinfectants used in water or surfaces in the animal production setting/environment; human and animal antiseptics).^{8,51,52} However, the relevance of *pco* gene cluster for Cu tolerance in *Salmonella* under aerobiosis was not here demonstrated, corroborating other studies.^{4,6} Additional research is needed to clarify the potential role of *pco* among *Salmonella* namely its involvement in virulence, as described for other multicopper oxidases.⁵³

In summary, we report a high occurrence and co-location of copper/silver tolerance and antibiotic resistance genes among *Salmonella* from diverse serotypes, and their particular dispersion in the emergent MDR *S. Rissen* and *S. Typhimurium* clones commonly associated with pig production. The fact that recently emerging MDR serotypes are enriched in *sil*+*pco* gene clusters also alerts for the relevance of the dispersion and fixation of these genes in other less frequent serotypes and their potential emergence in a near future. The extensive copper use as feed additive and the low O₂ levels (e.g. animal gut/manure/waste lagoons) found in certain locations of food-animal producing environments might have contributed for the maintenance and dispersion of *sil*+*pco* carrying MDR *Salmonella* and/or Tn6230-like elements along with antibiotic resistance in the animal production setting, surrounding environments and food chain. We urgently need to be aware of the role of reducing environments under multiple stressors (e.g. occurrence of the more toxic Cu⁺ or antibiotics) for an enhanced survival, selection and persistence of particular copper tolerant-MDR strains. An assessment of control measures for the use and/or accumulation of metals in diverse environments to prevent a wider expansion of MDR *S. Rissen* and *S. Typhimurium* clones or the emergence of new ones is also required.

ACKNOWLEDGMENTS

The authors wish to thank to Dr^a Maria del Rosario Rodicio (Department of Functional Biology, Area of Microbiology, University of Oviedo, Oviedo, Asturias, Spain) who kindly provided *Salmonella* Typhimurium LSP 448/02 for conjugation assays. We also wish to thank to Dr Jill Williams (Department of Genetics of the University of Melbourne, Melbourne, Victoria, Australia) and Drs Christopher P. Randall and Alex J. O'Neill (Antimicrobial Research Centre, School of Molecular and Cellular Biology, University of Leeds, Leeds, UK) who kindly provided *Escherichia coli* ED8739 (pRJ1004) and *E. coli* J53 (pMG101), respectively.

FUNDING

This work was supported by national funds [Fundação para a Ciência e a Tecnologia (FCT)] through project UID/MULTI/04378/2013. Joana Mourão and Joana Campos (grant numbers SFRH/BD/77518/2011, SFRH/BD/93091/2013 respectively; POPH-QREN) were supported by PhD fellowships from FCT. The authors are greatly indebted to all of the financing sources.

TRANSPARENCY DECLARATIONS

None to declare.

SUPPLEMENTARY DATA

Tables S1, S2 and S3 are available as Supplementary data at JAC Online (<http://jac.oxfordjournals.org/>).

REFERENCES

1. Antunes P, Mourão J, Pestana N *et al.* Leakage of emerging clinically relevant multidrug-resistant *Salmonella* clones from pig farms. *J Antimicrob Chemother* 2011; **66**: 2028-32.
2. Eng S-K, Pusparajah P, Ab Mutalib N-S *et al.* *Salmonella*: A review on pathogenesis, epidemiology and antibiotic resistance. *Front Life Sci* 2015; **8**: 284-93.
3. Gomes-Neves E, Antunes P, Manageiro V *et al.* Clinically relevant multidrug resistant *Salmonella* enterica in swine and meat handlers at the abattoir. *Vet Microbiol* 2014; **168**: 229-33.

4. Medardus JJ, Molla BZ, Nicol M *et al.* In-feed use of heavy metal micronutrients in U.S. swine production systems and its role in persistence of multidrug-resistant salmonellae. *Appl Environ Microbiol* 2014; **80**: 2317-25.
5. Mourão J, Machado J, Novais C *et al.* Characterization of the emerging clinically-relevant multidrug-resistant *Salmonella enterica* serotype 4,[5],12:i:- (monophasic variant of *S. Typhimurium*) clones. *Eur J Clin Microbiol Infect Dis* 2014; **33**: 2249-57.
6. Mourão J, Novais C, Machado J *et al.* Metal tolerance in emerging clinically relevant multidrug-resistant *Salmonella enterica* serotype 4,[5],12:i:- clones circulating in Europe. *Int J Antimicrob Agents* 2015a; **45**: 610-6.
7. List of the authorised additives in feedingstuffs (1) published in application of Article 9t (b) of Council Directive 70/524/EEC concerning additives in feedingstuffs of 25 February 2004. Official Journal of the European Union C50/1. http://ec.europa.eu/food/food/animalnutrition/feedadditives/docs/c_50_en.pdf.
8. Scientific Committee on Emerging and Newly Identified Health Risks (SCENIHR). Opinion on Nanosilver: safety, health and environmental effects and role in antimicrobial resistance. 2014. http://ec.europa.eu/health/scientific_committees/emerging/docs/scenih_r_o_039.pdf.
9. Seiler C, Berendonk TU. Heavy metal driven co-selection of antibiotic resistance in soil and water bodies impacted by agriculture and aquaculture. *Front Microbiol* 2012; **3**: 399.
10. Opinion of the Panel on Animal Feed of the Norwegian Scientific Committee for Food Safety (VKM). Zinc and copper in pig and poultry production – fate and effects in the food chain and the environment. 2014. <http://www.english.vkm.no/dav/3b1b6769dd.pdf>.
11. Yazdankhah S, Rudi K, Bernhoft A. Zinc and copper in animal feed - development of resistance and co-resistance to antimicrobial agents in bacteria of animal origin. *Microb Ecol Health Dis* 2014; **25**: 1-7.
12. European Food Safety Authority (EFSA). Opinion of the Scientific Panel on Contaminants in the Food chain on a request from the European Commission on mercury as undesirable substance in feed. *EFSA J* 2008; **654**: 1-76.
13. Bolan NS, Khan MA, Donaldson J *et al.* Distribution and bioavailability of copper in farm effluent. *Sci Total Environ* 2003; **309**: 225-36.
14. Xiong X, Yanxia L, Wei L *et al.* Copper content in animal manures and potential risk of soil copper pollution with animal manure use in agriculture. *Resour Conserv Recy* 2010; **54**: 985-90.

15. Zhang F, Li Y, Yang M *et al.* Content of heavy metals in animal feeds and manures from farms of different scales in northeast China. *Int J Environ Res Public Health* 2012; **9**: 2658-68.
16. European Food Safety Authority (EFSA) and Panel on Additives and Products or Substances used in Animal Feed (FEEDAP). Scientific Opinion on the safety and efficacy of copper compounds (E4) as feed additives for all animal species: cupric sulphate pentahydrate based on a dossier submitted by Manica S.p.A. *EFSA J* 2012; **10**: 2969.
17. Gaze WH, Krone SM, Larsson DG *et al.* Influence of humans on evolution and mobilization of environmental antibiotic resistance. *Emerg Infect Dis* 2013; **19**: e120871.
18. Baker-Austin C, Wright MS, Stepanauskas R *et al.* Co-selection of antibiotic and metal resistance. *Trends Microbiol* 2006; **14**: 176-82.
19. Hao X, Luthje FL, Qin Y *et al.* Survival in amoeba-a major selection pressure on the presence of bacterial copper and zinc resistance determinants? Identification of a "copper pathogenicity island". *Appl Microbiol Biotechnol* 2015; **99**: 5817-24.
20. Silver S. Bacterial silver resistance: molecular biology and uses and misuses of silver compounds. *FEMS Microbiol Rev* 2003; **27**: 341-53.
21. Antunes P, Coque TM, Peixe L. Emergence of an IncI_γ plasmid encoding CMY-2 β-lactamase associated with the international ST19 OXA-30-producing β-lactamase *Salmonella* Typhimurium multidrug-resistant clone. *J Antimicrob Chemother* 2010; **65**: 2097-100.
22. Antunes P, Machado J, Peixe L. Characterization of antimicrobial resistance and class 1 and 2 integrons in *Salmonella enterica* isolates from different sources in Portugal. *J Antimicrob Chemother* 2006; **58**: 297-304.
23. Antunes P, Machado J, Peixe L. Dissemination of *sul3*-containing elements linked to class 1 integrons with an unusual 3' conserved sequence region among *Salmonella* isolates. *Antimicrob Agents Chemother* 2007; **51**: 1545-8.
24. Antunes P, Mourão J, Alves T *et al.* *Salmonella enterica* serotype Bovismorbificans, a new host for CTX-M-9. *Int J Antimicrob Agents* 2013; **41**: 91-3.
25. Antunes P, Mourão J, Machado J *et al.* First description of *qnrS1*-IncN plasmid in a ST11 *Salmonella* Enteritidis clinical isolate from Portugal. *Diagn Microbiol Infect Dis* 2011b; **69**: 463-5.
26. Hasman H, Kempf I, Chidaine B *et al.* Copper resistance in *Enterococcus faecium*, mediated by the *tcrB* gene, is selected by supplementation of pig feed with copper sulfate. *Appl Environ Microbiol* 2006; **72**: 5784-9.

27. Randall CP, Oyama LB, Bostock JM *et al.* The silver cation (Ag^+): antistaphylococcal activity, mode of action and resistance studies. *J Antimicrob Chemother* 2013; **68**: 131-8.
28. Williams JR, Morgan AG, Rouch DA *et al.* Copper-resistant enteric bacteria from United Kingdom and Australian piggeries. *Appl Environ Microbiol* 1993; **59**: 2531-7.
29. Carattoli A, Bertini A, Villa L *et al.* Identification of plasmids by PCR-based replicon typing. *J Microbiol Methods* 2005; **63**: 219-28.
30. Garcia-Fernandez A, Fortini D, Veldman K *et al.* Characterization of plasmids harbouring *qnrS1*, *qnrB2* and *qnrB19* genes in *Salmonella*. *J Antimicrob Chemother* 2009; **63**: 274-81.
31. Pornsukarom S, Patchanee P, Erdman M *et al.* Comparative phenotypic and genotypic analyses of *Salmonella* Rissen that originated from food animals in Thailand and United States. *Zoonoses Public Health* 2015; **62**: 151-8.
32. Hölzel CS, Muller C, Harms KS *et al.* Heavy metals in liquid pig manure in light of bacterial antimicrobial resistance. *Environ Res* 2012; **113**: 21-7.
33. Ji X, Shen Q, Liu F *et al.* Antibiotic resistance gene abundances associated with antibiotics and heavy metals in animal manures and agricultural soils adjacent to feedlots in Shanghai; China. *J Hazard Mater* 2012; **235-236**: 178-85.
34. Zhu YG, Johnson TA, Su JQ *et al.* Diverse and abundant antibiotic resistance genes in Chinese swine farms. *Proc Natl Acad Sci U S A* 2013; **110**: 3435-40.
35. European Food Safety Authority (EFSA) and European Food Safety Authority and European Centre for Disease Prevention and Control (ECDC). The European Union summary report on trends and sources of zoonoses, zoonotic agents and food-borne outbreaks in 2013. *EFSA J* 2015; **13**: 3991.
36. Hoffmann M, Zhao S, Pettengill J *et al.* Comparative genomic analysis and virulence differences in closely related *Salmonella enterica* serotype Heidelberg isolates from humans, retail meats, and animals. *Genome Biol Evol* 2014; **6**: 1046-68.
37. Gerber P, Opio C, Steinfeld H. Poultry production and the environment – a review. In: Thieme O, Pilling D, editors. in Proceedings of the International Conference Poultry in the Twenty-First Century: Avian Influenza and Beyond. 2008, p. 379–405.
38. Carattoli A. Resistance plasmid families in Enterobacteriaceae. *Antimicrob Agents Chemother* 2009; **53**: 2227-38.
39. Garcia-Fernandez A, Carattoli A. Plasmid double locus sequence typing for IncHI2 plasmids, a subtyping scheme for the characterization of IncHI2 plasmids carrying extended-spectrum β -lactamase and quinolone resistance genes. *J Antimicrob Chemother* 2010; **65**: 1155-61.

40. Moreno Switt AI, den Bakker HC, Cummings CA *et al.* Identification and characterization of novel *Salmonella* mobile elements involved in the dissemination of genes linked to virulence and transmission. *PLoS One* 2012; **7**: e41247.
41. Qin Y, Hasman H, Aarestrup FM *et al.* Genome Sequences of Three Highly Copper-Resistant *Salmonella enterica* subsp. I Serovar Typhimurium Strains Isolated from Pigs in Denmark. *Genome Announc* 2014; **2**: e01334-14.
42. Peters JE, Fricker AD, Kapili BJ *et al.* Heteromeric transposase elements: generators of genomic islands across diverse bacteria. *Mol Microbiol* 2014; **93**: 1084-92.
43. Liebert CA, Wireman J, Smith T *et al.* Phylogeny of mercury resistance (*mer*) operons of gram-negative bacteria isolated from the fecal flora of primates. *Appl Environ Microbiol* 1997; **63**: 1066-76.
44. Boyd ES, Barkay T. The mercury resistance operon: from an origin in a geothermal environment to an efficient detoxification machine. *Front Microbiol* 2012; **3**: 349.
45. Berg J, Thorsen MK, Holm PE *et al.* Cu exposure under field conditions coselects for antibiotic resistance as determined by a novel cultivation-independent bacterial community tolerance assay. *Environ Sci Technol* 2010; **44**: 8724-8.
46. Knapp CW, McCluskey SM, Singh BK *et al.* Antibiotic resistance gene abundances correlate with metal and geochemical conditions in archived Scottish soils. *PLoS One* 2011; **6**: e27300.
47. Gullberg E, Albrecht LM, Karlsson C *et al.* Selection of a multidrug resistance plasmid by sublethal levels of antibiotics and heavy metals. *MBio* 2014; **5**: e01918-14.
48. Aarestrup FM, Hasman H. Susceptibility of different bacterial species isolated from food animals to copper sulphate, zinc chloride and antimicrobial substances used for disinfection. *Vet Microbiol* 2004; **100**: 83-9.
49. Mourão J, Rae J, Silveira E *et al.* Relevance of *tcrYAZB* operon acquisition for *Enterococcus* survival at high copper concentrations under anaerobic conditions. *J Antimicrob Chemother* 2016; **71**: 560-3.
50. Porcheron G, Garenaux A, Proulx J *et al.* Iron, copper, zinc, and manganese transport and regulation in pathogenic Enterobacteria: correlations between strains, site of infection and the relative importance of the different metal transport systems for virulence. *Front Cell Infect Microbiol* 2013; **3**: 90.
51. Maillard JY, Hartemann P. Silver as an antimicrobial: facts and gaps in knowledge. *Crit Rev Microbiol* 2013; **39**: 373-83.
52. Mijndendonckx K, Leys N, Mahillon J *et al.* Antimicrobial silver: uses, toxicity and potential for resistance. *Biomaterials* 2013; **26**: 609-21.

53. Achard ME, Tree JJ, Holden JA *et al.* The multi-copper-ion oxidase CueO of *Salmonella enterica* serovar Typhimurium is required for systemic virulence. *Infect Immun* 2010; **78**: 2312-9.

Table 1. Epidemiological features and genotypic/phenotypic characterization of *Salmonella enterica* isolates carrying acquired copper and silver tolerance genes (*sil* and *pco*)

Serotype (no. isolates)	"Clone designation" and PFGE-type (no. isolates)/ST ^a	Isolation date; region ^b	Source/sample (no. isolates) ^c	Antibiotic resistance profile/genes (no. isolates) ^d	Cu/Ag and other metal tolerance genes (no. isolates)	MIC _{range} CuSO ₄ Aerobic/anaerobic (mM) (no. isolates)	MIC _{range} AgNO ₃ Aerobic/with Ag contact (mM) (no. isolates)	Location of metal tolerance genes + co- location with antibiotic resistance genes – Chr or PL (Kb, Inc) (no. isolates) ^e
Typhimurium (n=37)	"European clone"-A (n=8), B (n=11), C (n=8), G (n=2) and F (n=1)/ST34	2003- 2004, 2007- 2013; N, C, S, I	Animal production/Pi (n=16); Food/B and U (n=2); Human/C (n=1) and H (n=11)	AMX, STR, SUL, TET, [GEN, KAN, NAL, CIP ^{DS}]/ <i>bla</i> _{TEM-1} , <i>strA</i> - <i>strB</i> , <i>sul2</i> , <i>tet</i> (B) [<i>tet</i> (A), <i>aac</i> (3)-IV, <i>aphA1</i>] (n=21) AMX, STR, SUL/ <i>bla</i> _{TEM-1} , <i>strA</i> - <i>strB</i> , <i>sul2</i> (n=1) AMX, SUL, TET/ <i>bla</i> _{TEM-1} , <i>sul2</i> , <i>tetB</i> (n=6) SUL, TET/ <i>tet</i> (B) (n=1) Susceptible (n=1)	<i>pcoA</i> - <i>pcoD</i> + <i>silA</i> - <i>silE</i> + <i>merA</i> (n=27) <i>pcoA</i> - <i>pcoD</i> + <i>silA</i> - <i>silE</i> (n=3)	28-32/20-32 (n=11)	0.125-0.5 (n=11)/>3 (n=4)	<i>pcoD</i> , <i>silA</i> , <i>merA</i> + <i>bla</i> _{TEM-1} , <i>strA</i> - <i>strB</i> , <i>sul2</i> , <i>tet</i> (B) – Chr (n=3)
Other - TB (n=2)/ST34		2012; S	Human/H (n=2)	AMX, CHL, CIP, GEN, KAN, STR, SUL, TET, TMP, NAL/ <i>bla</i> _{TEM-1} + <i>bla</i> _{OXA-30} , <i>ogxAB</i> , <i>aac</i> (6)-Ib-Cr, <i>catB3</i> , <i>cmiA1</i> , <i>flaR</i> , <i>aac</i> (3)-IV, <i>aphA1</i> , <i>aadA</i> , <i>strA</i> - <i>strB</i> , <i>sul1</i> - <i>sul2</i> - <i>sul3</i> , <i>tet</i> (B), <i>dfrA12</i> (n=2)	<i>pcoA</i> - <i>pcoD</i> + <i>silA</i> - <i>silE</i> + <i>merA</i> + <i>terF</i> (n=2)	36/24 (n=2)	0.125 (n=2)/>3 (n=1)	<i>pcoD</i> , <i>silA</i> , <i>merA</i> + <i>bla</i> _{TEM-1} , <i>strA</i> - <i>strB</i> , <i>tet</i> (B) – Chr; <i>terF</i> + <i>bla</i> _{OXA-30} , <i>ogxAB</i> , <i>aac</i> (6)- <i>Ib</i> -Cr, <i>catB3</i> , <i>cmiA1</i> , <i>flaR</i> , <i>aac</i> (3)-IV, <i>aphA1</i> , <i>aadA</i> , <i>sul1</i> - <i>sul2</i> - <i>sul3</i> , <i>dfrA12</i> – PL (180, H12) (n=2) <i>silA</i> , <i>merA</i> + <i>bla</i> _{TEM-1} , <i>sul3</i> , <i>dfrA12</i> – PL (100, F1A) (n=1) <i>silA</i> + <i>bla</i> _{TEM-1} , <i>cmiA1</i> , [<i>aadA1</i> - <i>aadA2</i>], [<i>sul1</i> - <i>sul3</i>], <i>tet</i> (A), <i>dfrA12</i> – PL (140, N) (n=3)
"DT104" – A (n=5)/ST19		2003- 2004; N	Food/P and U (n=2); Human/H (n=3)	AMX, CHL, STR, SUL, TET, TMP/ <i>bla</i> _{TEM-1} - <i>bla</i> _{SE-1} , <i>flaR</i> , <i>aadA2</i> , <i>sul1</i> - <i>sul3</i> , <i>tet</i> (G), <i>dfrA12</i> (n=1) AMX, CHL, STR, SUL, TET, TMP/ <i>bla</i> _{TEM-1} , <i>cmiA1</i> , <i>aadA2</i> - [<i>aadA1</i>], <i>sul1</i> -[<i>sul3</i>], <i>tet</i> (A), <i>dfrA12</i> (n=4)	<i>silA</i> - <i>silE</i> + <i>merA</i> (n=1) <i>silA</i> - <i>silE</i> (n=4)	28-36/24-32 (n=5)	0.125-0.25 (n=5)/>3 (n=2)	
Rissen (n=50)	N (n=49)/ST469	2002- 2004, 2006, 2007- 2013; N, C, S	Animal production/Pi (n=9); Food/ Cw, CM, P, Pt and U (n=25); Environmen/W (n=1) and WR (n=1); Human/C (n=3) and H (n=10)	AMX, STR, SUL, TET, TMP, [CHL]/ <i>bla</i> _{TEM-1} , <i>aadA2</i> , <i>sul1</i> , <i>tet</i> (A), <i>dfrA12</i> , [a <i>aadA1</i> , <i>sul3</i> , <i>catA</i> - <i>cmiA1</i>] (n=24) AMX, STR, SUL, TET, TMP/ <i>bla</i> _{TEM-1} , <i>aadA5</i> , <i>sul1</i> , <i>tet</i> (A), <i>dfrA17</i> (n=1) CHL, STR, SUL, TET, TMP/ <i>cmiA1</i> , <i>aadA2</i> - <i>aadA1</i> , <i>sul3</i> , <i>tet</i> (A), <i>dfrA12</i> (n=2) AMX, SUL, TET, TMP/ <i>bla</i> _{TEM-1} , <i>sul1</i> , <i>tet</i> (A), <i>dfrA12</i> , <i>aadA2</i> (n=1)	<i>pcoA</i> - <i>pcoD</i> + <i>silA</i> - <i>silE</i> + <i>merA</i> (n=2) <i>pcoA</i> - <i>pcoD</i> + <i>silA</i> - <i>silE</i> (n=47)	24-36/24-36 (n=33)	0.25-0.32 (n=33)/>3 (n=6)	<i>pcoD</i> , <i>silA</i> + [<i>tet</i> (A)] – Chr (n=14)

Table 1. Continued

Serotype (no. isolates)	"Clone designation" and PFGE-type (no. isolates)/ST ^a	Isolation date; region ^b	Source/sample (no. isolates) ^c	Antibiotic resistance profile/genes (no. isolates) ^d	Cu/Ag and other metal tolerance genes (no. isolates)	MIC _{range} CuSO ₄ Aerobic/anaerobic (mM) (no. isolates)	MIC _{range} AgNO ₃ Aerobic/with Ag contact (mM) (no. isolates)	Location of metal tolerance genes + co- location with antibiotic resistance genes – Chr or PL (Kb, Inc) (no. isolates) ^e
				AMX, STR, TET, TMP/ <i>bla</i> _{TEM-1} , <i>aadA2</i> , <i>tetA</i> , <i>dfrA12</i> (n=1) AMX, TET, CHL, STR/ <i>bla</i> _{TEM-1} , <i>tetA</i> (n=1) SUL, TET, TMP, [STR]/ <i>sul1</i> , <i>tetA</i> , <i>dfrA12</i> , <i>aadA2</i> (n=4) AMX, SUL, TMP/ <i>bla</i> _{TEM-1} , <i>sul1</i> , <i>dfrA12</i> , <i>aadA2</i> (n=1) TET, [STR]/ <i>tetA</i> , [a <i>aadA2</i>] (n=11) Susceptible (n=3)				
	Z (n=1)/ST469	2013; S	Human/H (n=1)	AMX, STR, SUL, TET, TMP/ <i>bla</i> _{TEM-1} , <i>aadA2</i> , <i>sul1</i> , <i>tetA</i> , <i>dfrA12</i> (n=1)	<i>pcoA-pcoD</i> + <i>silA-silE</i> (n=1)	36/32 (n=1)	0.25/>3 (n=1)	<i>pcoD</i> , <i>silA</i> + <i>bla</i> _{TEM-1} - Chr (n=1)
Anatum (n=2)	M (n=1)/ST64	2004; N	Human/C (n=1)	AMX, CHL, STR, SUL, TET, TMP/ <i>bla</i> _{TEM-1} , <i>aadA5</i> , <i>sul1</i> , <i>tetA</i> , <i>dfrA17</i> (n=1) SUL, TET, TMP/ <i>sul1</i> , <i>tetA</i> , <i>dfrA17</i> (n=1) AMX, CHL, STR, SUL, TET, TMP/ <i>bla</i> _{TEM-1} , <i>aadA1</i> , <i>sul1</i> , <i>tetA</i> , <i>dfrA1</i> (n=1)	<i>pcoA-pcoD</i> + <i>silA-silE</i> + <i>merA</i> (n=2) <i>pcoA-pcoD</i> + <i>silA-silE</i> + <i>merA</i> + <i>terF</i> (n=1)	28-36/24-32 (n=2)	0.25-0.5/>3 (n=2)	<i>pcoD</i> , <i>silA</i> , <i>merA</i> + <i>aadA5</i> , <i>sul1</i> , <i>dfrA17</i> - Chr (n=1) <i>pcoD</i> , <i>silA</i> , <i>merA</i> + <i>sul1</i> , <i>tetA</i> - Chr (n=1) <i>pcoD</i> , <i>silA</i> , <i>merA</i> , <i>terF</i> + <i>bla</i> _{TEM-1} , <i>aadA1</i> , <i>sul1</i> , <i>dfrA1</i> - PL (260, H12) (n=1) <i>pcoD</i> , <i>silA</i> , <i>merA</i> + <i>sul1</i> , <i>dfrA1</i> , <i>aadA1</i> - Chr (n=1)
Brandenburg (n=2)	Y (n=1)/ST64 G (n=1)/ST334	2002; N 2003; N	Food/P (n=1) Food/P (n=1)	SUL, TET, TMP/ <i>sul1</i> , <i>tetA</i> , <i>dfrA17</i> (n=1) AMX, CHL, STR, SUL, TET, TMP/ <i>bla</i> _{TEM-1} , <i>aadA1</i> , <i>sul1</i> , <i>tetA</i> , <i>dfrA1</i> (n=1)	<i>pcoA-pcoD</i> + <i>silA-silE</i> + <i>merA</i> + <i>terF</i> (n=1)	32/24 (n=2)	0.16/>3 (n=2)	
	V (n=1)/ST334	2004; N	Human/H (n=1)	SUL, TET, TMP/ <i>sul1</i> , <i>tetA</i> , <i>dfrA1</i> , <i>aadA1</i> (n=1)	<i>pcoA-pcoD</i> + <i>silA-silE</i> + <i>merA</i> (n=1) <i>pcoA-pcoD</i> + <i>silA-silE</i> + <i>merA</i> + <i>terF</i> (n=2) <i>pcoA-pcoD</i> + <i>silA-silE</i> + <i>merA</i> (n=2)	28-36/28-32 (n=4)	0.125-0.25 (n=4)/>3 (n=1)	<i>pcoD</i> , <i>silA</i> , <i>merA</i> + <i>sul1</i> , <i>dfrA1</i> , <i>aadA1</i> - Chr (n=1)
Bredeney (n=4)	L (n=4)/ST306	2002, 2004, 2010; N	Food/OT (n=1); Human/H (n=3)	SUL, TET, TMP/ <i>sul1</i> , <i>tetA</i> , <i>dfrA17</i> , <i>aadA5</i> (n=1) TET, TMP/ <i>tetA</i> , <i>dfrA17</i> , <i>sul1</i> , <i>aadA5</i> (n=2)	<i>pcoA-pcoD</i> + <i>silA-silE</i> + <i>merA</i> (n=1) <i>pcoA-pcoD</i> + <i>silA-silE</i> + <i>merA</i> (n=2)	28-36/28-32 (n=4)	0.125-0.25 (n=4)/>3 (n=1)	<i>pcoD</i> , <i>silA</i> , <i>merA</i> + <i>sul1</i> , <i>tetA</i> , <i>dfrA17</i> , <i>aadA5</i> - PL (195, UN); <i>terF</i> - Chr (n=1) <i>pcoD</i> , <i>silA</i> , <i>merA</i> , [i <i>terF</i>] + <i>sul1</i> , <i>dfrA17</i> , [a <i>aadA5</i>] - Chr (n=3)
Give (n=1)	U (n=1)/STNew	2010; C	Food/U (n=1)	SUL, TMP/ <i>sul1</i> , <i>dfrA17</i> (n=1) Susceptible (n=1)	<i>pcoA-pcoD</i> + <i>silA-silE</i> (n=1)	36/32 (n=1)	0.25/0.5 (n=1)	<i>pcoD</i> , <i>silA</i> - Chr (n=1)
Muenchen (n=1)	D (n=1)/ST82	2003; N	Human/H (n=1)	CHL, STR, SUL, TMP/ <i>catA</i> , <i>aadA1</i> , <i>sul1</i> , <i>dfrA1</i> (n=1)	<i>pcoA-pcoD</i> + <i>silA-silE</i> + <i>merA</i> + <i>terF</i> (n=1)	36/32 (n=1)	0.32/>3 (n=1)	<i>pcoD</i> , <i>silA</i> , <i>merA</i> , <i>terF</i> + <i>catA</i> , <i>aadA1</i> , <i>sul1</i> , <i>dfrA1</i> - PL (270, H12) (n=1) <i>pcoD</i> , <i>silA</i> - Chr (n=3)
Senftenberg (n=3)	SA (n=3)/ST14	2002- 2003; N, C	Animal production/Pi (n=1); Food/Pt (n=2)	NAL, CIP ^{DS} /ND (n=1) Susceptible (n=2)	<i>pcoA-pcoD</i> + <i>silA-silE</i> (n=3)	32-36/32 (n=3)	0.25-0.32 (n=3)/>3 (n=1)	

Table 1. Continued

Serotype (no. isolates)	"Clone designation" and PFGE-type (no. isolates)/ST ^a	Isolation date; region ^b	Source/sample (no. isolates) ^c	Antibiotic resistance profile/genes (no. isolates) ^d	Cu/Ag and other metal tolerance genes (no. isolates)	MIC _{range} CuSO ₄ Aerobic/anaerobic (mM) (no. isolates)	MIC _{range} AgNO ₃ Aerobic/with Ag contact (mM) (no. isolates)	Location of metal tolerance genes + co- location with antibiotic resistance genes – Chr or PL (Kb, Inc) (no. isolates) ^e
Wien (n=1)	J (n=1)/ST102	2003; S	Food/P (n=1)	AMX, CHL, KAN, STR, SUL, TET, TMP/bla _{TEM-1} , cmlA1-catA, aphA1, aadA2-aadA1, sul1-sul3, tet(B), dfrA1 (n=1)	pcoA-pcoD + silA-silE (n=1)	32/28 (n=1)	0.25/>3 (n=1)	pcoD, silA + bla _{TEM-1} , cmlA1- catA, aphA1, aadA2-aadA1, sul1-sul3, tet(B), dfrA1 – PL (250, H1) (n=1)

AMX, amoxicillin; CHL, chloramphenicol; CIP, ciprofloxacin; CIP^{DS}, decreased susceptibility to ciprofloxacin, GEN, gentamicin; KAN, kanamycin; NAL, nalidixic acid; STR, streptomycin; SUL, sulfamethoxazole; TET, tetracycline; TMP, trimethoprim; Chr, chromosome; PL, plasmid; ND, not determined; UN, untypable.

^a"Clone designation" was adopted only when has been previously published. PFGE-types were designated by capital letters, including previously described^{1,22,23}, or firstly designated in this study.

^bN, north; C, centre; S, south; I, islands (Azores).

^cB, beef; C, community; CM, cooked meal; Cw, cow; H, hospitals patients; OT, other type of food; P, pork; Pi, pigs and piggeries environment; Pt, poultry; U, unknown; W, bathing or drinking water; WR, river water.

^dVariable antibiotic resistance phenotypes and genotypes were presented between square brackets. Isolates resistant to three or more antibiotics from different families were considered multidrug-resistant.

^eVariable metal tolerance genes were presented between square brackets. Underlined metal tolerance or antibiotic resistance genes were also detected in transconjugants.

^fIn particular serotypes/clonal lineages *silA*±*pcoD*, *merA* or *terF* were in the same genetic platform that the genes belonging to conventional class 1 integrons [1000bp (*aadA1*) in *S. Muenchen*; 1000bp (*aadA2*) in *S. Typhimurium* "DT104" clone; 1700bp (*dfrA1-aadA1*) in *S. Brandenburg* and *S. Wien* and 1800bp (*dfrA17-aadA5*) in *S. Bredeney* and *S. Anatum*] or to atypical *sul3* [type I integron 7085bp (*dfrA12-orfF-aadA2-cmlA1-aadA1-qacH-IS440-sul3*) in *S. Typhimurium* "DT104" clone and type III integron 7304bp (*estX-psp-aadA2-cmlA1-aadA1-qacH-IS440-sul3*)] in *S. Wien*].^{22,23}

Table 2. Minimum inhibitory concentrations of CuSO₄ for *Salmonella enterica* isolates

Serotype	Environment	silA- silE	pcoA- pcoE	No. of Isolates ^a	MIC (mM) ^b																MIC ₅₀ (mM)	MIC ₉₀ (mM)	
					0.5	1	2	4	8	12	16	20	24	28	32	36							
Typhimurium	Aerobic	+	+	13													4	7	2	32	36		
		+	-	5													2	2	1	32	36		
		-	-	32									1				10	20	1	32	32		
		+	+	13									1				10	1	1	24	28		
		+	-	5									1				1	3	1	28	32		
Rissen	Aerobic	-	-	32		1	18	13												4	8		
	Anaerobic	+	+	34										1			1	21	12	32	36		
Enteritidis	Aerobic	+	+	34									1	19	13	1		13	1	28	32		
	Anaerobic	-	-	14													10	4		28	32		
Other	Aerobic	-	-	14		4	8	2												2	4		
		+	+	14													2	5	7	32	36		
	Anaerobic	-	-	54																	36		
		+	+	14													3	4	7	28	32		
		-	-	54		13	33	2	3	1							1 ^c	1 ^c		2	8		
TOTAL	Aerobic	+	+	or -	66												1	8	35	22	32	36	
		-	-	100													1	25	50	24	32	36	
	Anaerobic	+	+	or -	66												1	15	27	22	1	28	32
		-	-	100		17	42	22	16	1							1 ^c	1 ^c		2	8		

^aThe number of isolates included representatives of all the serotypes: *Salmonella enterica* Typhimurium (n=50/96), *Salmonella enterica* Rissen (n=34/50), *Salmonella enterica* Enteritidis (n=14/32) and other *Salmonella enterica* serotypes (n=68/97).

^b*Escherichia coli* ED8739 (plasmid pRJ1004 with *pco* genes; MIC_{CuSO₄} = 32-36 mM, aerobiosis)²⁸ and *Enterococcus faecium* BM4105RF (negative for all genes tested; MIC_{CuSO₄} = 8-12 mM, aerobiosis)²⁶ were used as control strains in Cu assays; Those strains were also used in anaerobic assays: *E. coli* ED8739 (plasmid pRJ1004 with *pco* genes; MIC_{CuSO₄} = 16-20 mM) and *E. faecium* BM4105RF (MIC_{CuSO₄} = 2-4 mM).⁶ A PCR approach confirmed the presence or absence of *silCPRS* and *pcoRSE* gene clusters in representative isolates of all anaerobic MIC.

Table 2. *Continued*

The corresponding CuSO4 mg/L values for the mM-values tested are: 0.25 mM-39.90 mg/L, 0.5 mM-79.80 mg/L, 1 mM-159.61 mg/L, 2 mM-319.22 mg/L, 4 mM-638.44 mg/L, 8 mM-1276.87 mg/L, 12 mM-1915.308 mg/L, 16 mM-2553.74 mg/L, 20 mM-3192.18 mg/L, 24 mM-3830.62 mg/L, 28 mM-4469.05 mg/L, 32 mM-5107.49 mg/L, 36 mM-5745.92 mg/L.

^aIsolates without amplification of *silCPRS* and *pcoRSE* gene clusters by PCR.

Table 3. Minimum inhibitory concentrations of AgNO₃ for *Salmonella enterica* isolates

Serotype	Environment	siIA-silE	No. of Isolates ^a	MIC (mM) ^b										MIC ₅₀ (mM)	MIC ₉₀ (mM)
				0.08	0.125	0.16	0.25	0.32	0.5	0.75	1	1.5	>3		
Typhimurium	Aerobic	+	18	5	2	9	2							0.25	0.5
		-	32	13	19									0.16	0.16
	Anaerobic	+	18	2	1	2	13							0.5	0.5
		-	32	11	2	18	1							0.25	0.25
Rissen	Aerobic/previous contact with 0.025 mM AgNO ₃	+	7									7	>3		>3
		-	4	4										0.125	0.125
	Aerobic	+	34			33	1							0.25	0.25
	Anaerobic	+	34			11	11							0.32	0.5
Enteritidis	Aerobic/previous contact with 0.025 mM AgNO ₃	+	7									7	>3		>3
		-	14												
	Aerobic	-	14	12	2									0.125	0.16
	Anaerobic	-	14		1	13								0.25	0.25
Other	Aerobic/previous contact with 0.025 mM AgNO ₃	-	6	5	1									0.125	0.16
		+	14												
	Aerobic	+	14	1	2	8	2	1						0.25	0.32
		-	54	33	16	5								0.125	0.16
TOTAL	Anaerobic	+	14		3	3	7	1						0.5	0.5
		-	54	24	5	10	2	8	5					0.125	0.32
	Aerobic/previous contact with 0.025 mM AgNO ₃	+	9						1			8	>3		>3
		-	5	3	2									0.125	0.16
TOTAL	Aerobic	+	66	6	4	50	3	3	3					0.25	0.25
		-	100	58	37	5								0.125	0.16
	Anaerobic	+	66	2	4	16	11	31	2					0.32	0.5
		-	100	24	16	13	8	6						0.16	0.32
TOTAL	Aerobic/previous contact with 0.025 mM AgNO ₃	+	23						1			22	>3		>3
		-	15	12	3									0.125	0.16

Table 3. Continued

^aThe number of isolates included representatives of all the serotypes: *Salmonella enterica* Typhimurium (n=50/96), *Salmonella enterica* Rissen (n=34/50), *Salmonella enterica* Enteritidis (n=14/32) and other *Salmonella enterica* serotypes (n=68/97).

^b*Escherichia coli* J53 (plasmid pMG101 with *sil* genes; MIC_{AgNO₃}>3 mM, aerobiosis)²⁷ and *Staphylococcus aureus* ATCC 29213 [negative for all genes tested; MIC_{AgNO₃}=0.16-0.25 mM, aerobiosis]²⁷ were used as control strains in Ag assays. Those strains were also used in anaerobic assays: *E. coli* J53 (plasmid pMG101 with *sil* genes; MIC_{AgNO₃}>3 mM) and *S. aureus* ATCC 29213 (MIC_{AgNO₃}=0.32-0.5 mM).⁶ A PCR approach confirmed the presence or absence of *silCPRS* gene cluster in representative isolates of all MIC after previous contact with Ag. The corresponding mg/L values for the mM-values tested are: 0.08 mM-13.59 mg/L, 0.125 mM-21.23 mg/L, 0.16 mM-27.18 mg/L, 0.25 mM-42.47 mg/L, 0.32 mM-54.36 mg/L, 0.5 mM-84.94 mg/L, 0.75 mM-127.40 mg/L, 1 mM-169.87 mg/L, 1.5 mM-254.81 mg/L, >3 mM-509.62 mg/L.

Table 4. Co-transfer of copper and silver tolerance with antibiotic resistance

<i>Salmonella</i> serotypes (no. of positive mating assays)	TRANSCONJUGANTS									
	Compounds used as selective agents	Recipient strains used ^a	MIC _{CuSO4} (mM)			MIC _{AgNO3} (mM)	Aerobiosis with Ag contact	Cu/Ag and other metal tolerance genes	Antibiotic resistance phenotype	Location of metal tolerance genes + co-location with antibiotic resistance genes – PL (Kb, Inc)
			Aerobiosis	Anaerobiosis	Aerobiosis					
Typhimurium (1/3)	SUL (256 mg/L) TMP	<i>E. coli</i> K802N	24	20	0.125	^b		<i>silA</i> + <i>merA</i>	AMX, SUL, TMP, NAL	<i>silA</i> , <i>merA</i> + <i>bla</i> _{TEM} , <i>sul3</i> , <i>dfrA12</i> – PL (100, FIIA)
Bredeney (1/1)	(8 mg/L) SUL	<i>S. enterica</i> LSP 448/02	>36	28	0.25	>3		<i>silA</i> + <i>merA</i>	AMX, SUL, TMP, NAL	
	(256 mg/L) TMP	<i>E. coli</i> K802N	24	20	0.08	^b		<i>pcoD</i> + <i>silA</i> + <i>merA</i>	SUL, TET, TMP, NAL	<i>pcoD</i> , <i>silA</i> , <i>merA</i> + <i>sul1</i> , <i>tet(A)</i> , <i>dfrA17</i> , <i>aadA5</i> ^c – PL (195, UN)
Muenchen (1/1)	(8 mg/L) SUL	<i>S. enterica</i> LSP 448/02	>36	28	0.25	>3		<i>pcoD</i> + <i>silA</i> + <i>merA</i>	SUL, TET, TMP, NAL	
	(256 mg/L) STR	<i>E. coli</i> K802N	24	20	0.08	^b		<i>pcoD</i> + <i>silA</i> + <i>merA</i> + <i>terF</i>	CHL, STR, SUL, TMP, NAL	<i>pcoD</i> , <i>silA</i> , <i>merA</i> , <i>terF</i> + <i>catA</i> , <i>aadA1</i> , <i>sul1</i> , <i>dfrA1</i> ^c – PL (270, HI2)
Wien (1/1)	(32 mg/L) TMP	<i>S. enterica</i> LSP 448/02	>36	28	0.25	>3		<i>pcoD</i> + <i>silA</i> + <i>merA</i> + <i>terF</i>	CHL, STR, SUL, TMP, NAL	
	(8 mg/L) SUL	<i>S. enterica</i> LSP 448/02	>36	28	0.25	>3		<i>pcoD</i> + <i>silA</i> + <i>merA</i> + <i>terF</i>	CHL, STR, SUL, TMP, NAL	
Wien (1/1)	(256 mg/L) SUL	<i>E. coli</i> K802N	24	20	0.08	^b		<i>pcoD</i> + <i>silA</i>	AMX, CHL, KAN, STR, SUL, TET, TMP, NAL	<i>pcoD</i> , <i>silA</i> + <i>bla</i> _{TEM} , <i>cmiA1</i> - <i>catA</i> , <i>aphA1</i> , <i>aadA2</i> - <i>aadA1</i> , <i>sul1</i> - <i>sul3</i> , <i>tet(B)</i> , <i>dfrA1</i> ^c – PL (250, HI1)
	(16 mg/L) AMX	<i>S. enterica</i> LSP 448/02	>36	28	0.25	>3		<i>pcoD</i> + <i>silA</i>	AMX, CHL, KAN, STR, SUL, TET, TMP, NAL	

AMX, amoxicillin; CHL, chloramphenicol; KAN, kanamycin; NAL, nalidixic acid; STR, streptomycin; SUL, sulfamethoxazole; TET, tetracycline; TMP, trimethoprim; PL, plasmid; UN, untypeable.

Table 4. Continued

^aAntibiotic resistance and metal tolerance of the recipient strains used in conjugation assays: *Escherichia coli* K802N (rifampicin and nalidixic acid resistant; MIC_{CuSO₄}=24-28 mM, aerobiosis and MIC_{CuSO₄}=1-2 mM, anaerobiosis; MIC_{AgNO₃}=0.08-0.125 mM, aerobiosis); *Salmonella* Typhimurium LSP 448/02 (nalidixic acid resistant; MIC_{CuSO₄}=32-36 mM, aerobiosis and MIC_{CuSO₄}=8-12 mM, anaerobiosis; MIC_{AgNO₃}=0.16-0.25 mM, aerobiosis and MIC_{Ag contact}=0.125-0.25 mM).

^bThe described Ag protocol assay was designed for testing *Salmonella* isolates.

^cIn particular serotypes/clonal lineages *slfA±pcoD*, *merA* or *terF* were in the same genetic platform that the genes belonging to conventional class 1 integrons [*aadA1*] in *S. Muenchen*; 1700bp (*dfrA1-aadA1*) in *S. Wien* and 1800bp (*dfrA17-aadA5*) in *S. Bredeney*] or to atypical *su13* [type III integron 7304bp (*estX-psp-aadA2-cmIA1-aadA1-qacH-IS440-sul3*)] in *S. Wien*.^{22,23}

Table S1. Epidemiological features and genotypic/phenotypic characterization of *Salmonella enterica* isolates with absence of acquired copper and silver tolerance genes (*sil* and *pco*)

Serotype (no. isolates)	"Clone designation" and PFGE-type (no. isolates)/ST ^a	Isolation date; region ^b	Source/sample (no. isolates) ^c	Antibiotic resistance profile/genes (no. isolates) ^d	Other metal tolerance genes + co-location with antibiotic resistance genes - Chr or PL (Kb, Inc) (no. isolates) ^e	MIC _{range} CuSO ₄ Aerobic/anaerobic (mM) (no. isolates)	MIC _{range} AgNO ₃ Aerobic/with Ag contact (mM) (no. isolates)
Typhimurium (n=59)	"DT104"-A (n=28)/ST19	2002-2004, 2007-2014; N, C, S, I	Animal production/Pi (n=9); Food/P or U (n=4); Human/ C (n=2) and H (n=13)	AMX, CHL, STR, SUL, TET, [GEN, NAL, CIP ^{DS}]/ <i>bla</i> _{PSE-1} , <i>floR</i> , <i>aadA2</i> , <i>sul1</i> , <i>tet</i> (G), [<i>aac</i> (3)-IV, <i>sul2</i>] (n=22) AMX, CHL, STR, SUL/ <i>bla</i> _{PSE-1} , <i>floR</i> , <i>aadA2</i> , <i>sul1</i> (n=1) AMX, STR, SUL, [NAL]/ <i>bla</i> _{PSE-1} , <i>aadA2</i> , <i>sul1</i> (n=2) AMX, SUL/ <i>bla</i> _{PSE-1} , <i>sul1</i> (n=1) STR, SUL/ <i>aadA2</i> , <i>sul1</i> (n=2) CHL, KAN, SUL, TMP/ <i>cmiA1</i> , <i>aphA1</i> , <i>sul3</i> , <i>dfra12</i> , <i>aadA2</i> - <i>aadA1</i> (n=1) AMX, CHL, STR, SUL, TET, [GEN]/ <i>bla</i> _{OXA-30} , <i>catA</i> , <i>aadA1</i> , <i>sul1</i> , <i>tet</i> (B), [<i>bla</i> _{CMY-2} , <i>floR</i> , <i>sul2</i> , <i>aac</i> (3)- IV] (n=15) AMX, CHL, STR, SUL/ <i>bla</i> _{OXA-30} , <i>catA</i> , <i>aadA1</i> , <i>sul1</i> (n=1) AMX, STR, SUL, TET/ <i>bla</i> _{OXA-30} , <i>aadA1</i> , <i>sul1</i> , <i>tet</i> (B) (n=1) AMX, STR, SUL/ <i>bla</i> _{OXA-30} , <i>aadA1</i> , <i>sul1</i> (n=1)	(-) (n=28)	28-36/4-8 (n=14)	0.125-0.16 (n=14)/0.125 (n=2)
	"DT104"-Q (n=1)/ST19	2000; N	Human/H (n=1)		<i>merA</i> + <i>cmiA1</i> , <i>aphA1</i> , <i>sul3</i> , <i>dfra12</i> , <i>aadA2</i> - <i>aadA1</i> – PL (100, Flis) (n=1)	32/2 (n=1)	0.16/ND (n=1)
	"OXA-30- producing"-T (n=18)/ST19	2002-2004, 2008-2013; N, C, S	Food/P or Pt (n=4); Environment/U (n=1); Human/H (n=13)		<i>merA</i> + <i>bla</i> _{OXA-30} , <i>catA</i> , <i>aadA1</i> , <i>sul1</i> , <i>tet</i> (B) – PL (100-140, Flis) (n=4) <i>merA</i> – ND (n=14)	28-32/4-8 (n=9)	0.125-0.16 (n=9)/0.125 (n=1)

Serotype (no. isolates)	"Clone designation" and PFGE-type (no. isolates)/ST ^a	Isolation date; region ^b	Source/sample (no. isolates) ^c	Antibiotic resistance profile/genes (no. isolates) ^d	Other metal tolerance genes + co-location with antibiotic resistance genes - Chr or PL (Kb, Inc) (no. isolates) ^e	MIC _{range} CuSO ₄ Aerobic/anaerobic (mM) (no. isolates)	MIC _{range} AgNO ₃ Aerobic/with Ag contact (mM) (no. isolates)
	"Other" - Z (n=2)/ST19, AT/ND (n=1), AM/ND (n=1) and ND (n=8)	2002-2004, 2010, 2012, 2013; N, S	Food/Cw, P or Q (n=3); Environment/W (n=2); Human/H (n=7)	AMX, CHL, STR, SUL, TET, TMP/ <i>bla</i> _{TEM-1} , <i>cmiA1</i> , <i>aadA2</i> - <i>aadA1</i> , <i>sul3</i> , <i>tet(A)</i> , <i>dfrA12</i> , [<i>sul2</i>] (n=2)	(-) (n=12)	20-32/4-8 (n=8)	0.125-0.16 (n=8)/0.125 (n=1)
				AMX, STR, SUL, TET/ <i>bla</i> _{TEM-1} , <i>strA-strB</i> , <i>sul2</i> , <i>tet(A)</i> (n=2)			
				AMX, SUL, TET, TMP/ <i>bla</i> _{TEM-1} , <i>sul2</i> , <i>tet(A)</i> (n=2)			
				AMX, TET/ <i>bla</i> _{TEM-1} , <i>tet(A)</i> (n=1)			
				TET/ <i>tet(A)</i> (n=1)			
				AMX/ <i>bla</i> _{TEM-1} (n=1)			
				NAL, CIP ^{DS} /ND (n=2)			
				Susceptible (n=1)			
				SUL, TET, TMP, [NAL, CIP ^{DS}]/ <i>sul1</i> , <i>tet(A)</i> , <i>dfrA1</i> , <i>aadA1</i> (n=6)			
				AMX, STR, SUL, TMP/ <i>bla</i> _{TEM-1} , <i>aadA1</i> , <i>sul1-sul2</i> , <i>dfrA1</i> (n=1)			
Enteritidis (n=32)	C (n=10)/ST11	2002-2004, 2008; N, S, Unknown	Animal production/Pt (n=1); Food/Pt or U (n=3); Human/H (n=6)	AMX, STR, SUL, NAL, CIP ^{DS} / <i>bla</i> _{TEM-1} , <i>aadA1</i> , <i>sul1</i> (n=1)	<u><i>merA</i> + <i>sul1</i>, <i>tet(A)</i>, <i>dfrA1</i></u> <u><i>aadA1</i> - PL (80, P)</u> (n=1) <i>merA</i> - ND (n=3) (-) (n=3)	28-32/1-2 (n=4)	0.125-0.16 (n=4)/0.125 (n=3)
				NAL, TMP, CIP ^{DS} / <i>qnrS1</i> , <i>dfrA14</i> (n=1)			
				NAL, CIP ^{DS} /ND (n=1)			
				SUL, TET/ <i>sul1</i> , <i>tet(A)</i> , <i>dfrA1</i> , <i>aadA1</i> (n=1)			
				AMX, SUL, TET, TMP/ <i>bla</i> _{TEM-1} , <i>sul2</i> , <i>tet(A)</i> (n=1)			
				STR, SUL, TET, NAL, CIP ^{DS} / <i>strA</i> - <i>strB</i> , <i>sul2</i> , <i>tet(A)</i> (n=1)			
				AMX, GEN/ <i>bla</i> _{TEM-1} , <i>aac(3)-IV</i> (n=1)			
				AMX, [NAL, CIP ^{DS}]/ <i>bla</i> _{TEM-1} (n=4)			
				CHL, NAL, CIP ^{DS} /(-) (n=1)			
				TET, [NAL, CIP ^{DS}]/ <i>tet(A)</i> (n=2)			
Adelaide (n=1) Agona (n=1)	AA (n=1)/ST440 ND (n=1)	2012; N 2010; S	Human/H (n=1) Human/H (n=1)	Susceptible (n=5)	(-) (n=1) (-) (n=1)	28/2 (n=1) 32/2 (n=1)	0.25/0.125 (n=1) 0.125/ND (n=1)
				Susceptible (n=1)			

Serotype (no. isolates)	"Clone designation" and PFGE-type (no. isolates)/ST ^a	Isolation date; region ^b	Source/sample (no. isolates) ^c	Antibiotic resistance profile/genes (no. isolates) ^d	Other metal tolerance genes + co-location with antibiotic resistance genes - Chr or PL (Kb, Inc) (no. isolates) ^e	MIC _{range} CuSO ₄ Aerobic/anaerobic (mM) (no. isolates)	MIC _{range} AgNO ₃ Aerobic/with Ag contact (mM) (no. isolates)
Anatum (n=1)	S (n=1)/ST64	2003; S	Human/H (n=1)	CHL, STR, SUL, TMP/ <i>cmiA1</i> , <i>aadA2-aadA1</i> , <i>sul3</i> , <i>dfrA12</i> (n=1)	<i>merA</i> + <i>cmiA1</i> , <i>aadA2-aadA1</i> , <i>sul3</i> , <i>dfrA12</i> - PL (120, 11) (n=1)	32/1 (n=1)	0.16/0.16 (n=1)
Bovismorbificans (n=1)	W (n=1)/ST142	2011; N	Human/H (n=1)	AMX, CTX, STR, SUL, GEN, KAN/ <i>bla</i> _{CTX-M-9} , <i>aadB-aadA2</i> , <i>sul1</i> (n=1)	<i>arsB</i> + <i>terF</i> + <i>bla</i> _{CTX-M-9} , <i>aadB</i> , <i>aadA2</i> , <i>sul1</i> - PL (240, H12) (n=1)	32/8 (n=1)	0.16/0.125 (n=1)
Brandenburg (n=1)	ND (n=1)	2003; S	Food/P (n=1)	STR, SUL, TET/ <i>strA-strB</i> , <i>sul2</i> , <i>tet(A)</i> (n=1)	(-) (n=1)	36/2 (n=1)	0.16/ND (n=1)
Bredeney (n=1)	ND (n=1)	2003; N	Food/P (n=1)	Susceptible (n=1)	(-) (n=1)	32/2 (n=1)	0.16/ND (n=1)
Brikama (n=1)	ND (n=1)	2011; S	Human/H (n=1)	Susceptible (n=1)	(-) (n=1)	36/2 (n=1)	0.125/ND (n=1)
Budapest (n=2)	BA (n=1)/ST1672	2011; N	Animal production/A (n=1)	STR(-) (n=1)	(-) (n=1)	36/2 (n=1)	0.125/ND (n=1)
Derby (n=16)	ND (n=1) B (n=14)/ST40	2011; N 2003-2004, 2007-2008; N, S	Human/H (n=1) Animal production/Pi (n=11); Food/P (n=2); Human/H (n=1)	Susceptible (n=1) AMX, STR, SUL, TET, GEN, KAN/ <i>bla</i> _{TEM-1} , <i>aadA2</i> , <i>sul1</i> , <i>tet(A)</i> (n=1) STR, SUL, TET/ <i>aadA2</i> , <i>sul1</i> , <i>tet(A)</i> (n=8) Susceptible (n=5)	(-) (n=1) <i>merA</i> + <i>aadA2</i> , <i>sul1</i> , <i>tet(A)</i> - Chr (n=1) <i>merA</i> - ND (n=8) (-) (n=5)	36/2 (n=1) 32-36/1-2 (n=3)	0.125/ND (n=1) 0.16-0.25/ND (n=3)
Essen (n=1)	ND (n=2)	2002, 2010; S	Food/P (n=1); Human/H (n=1)	STR, SUL/ <i>aadA2</i> , <i>sul1</i> (n=1) TET/(-) (n=1)	(-) (n=2)		
Farsta (n=1)	ND (n=1)	2003; C	Environment/W (n=1)	Susceptible (n=1)	(-) (n=1)	32/24 (n=1)	0.125/ND (n=1)
Freiburg (n=1)	ND (n=1)	2002; S	Food/Pt (n=1)	NAL, CIP ^{DS} /ND (n=1)	(-) (n=1)	36/2 (n=1)	0.125/ND (n=1)
Gaillac (n=1)	ND (n=1)	2010; N	Human/C (n=1)	Susceptible (n=1)	(-) (n=1)	36/2 (n=1)	0.125/ND (n=1)
Give (n=4)	ND (n=2)	2002; S 2002-2003; N	Food/OT (n=1) Food/P (n=2)	Susceptible (n=1) Susceptible (n=2)	(-) (n=1) (-) (n=2)	36/2 (n=1) 36/2 (n=1)	0.125/ND (n=1) 0.16/ND (n=1)
Gloucester (n=1)	GA (n=2)	2008; N	Animal production/Pi (n=2)	Susceptible (n=2)	(-) (n=2)		
Goldcoast (n=2)	ND (n=1) R (n=1)/ST358	2004; S 2003; N	Food/P (n=1) Human/H (n=1)	Susceptible (n=1) AMX, CHL, GEN, STR, SUL, TET, TMP/ <i>bla</i> _{TEM-1} , <i>catA</i> , <i>aadA2</i> , <i>sul1</i> - <i>sul2</i> , <i>tet(A)</i> , <i>dfrA12</i> (n=1)	(-) (n=1) <i>merA</i> + <i>bla</i> _{TEM-1} , <i>aadA2</i> , <i>sul1</i> - <i>sul2</i> , <i>tet(A)</i> , <i>dfrA12</i> - PL (95, UN) (n=1)	32/2 (n=1) 28-36/2 (n=2)	0.125/ND (n=1) 0.16-0.25 (n=2)/0.16 (n=1)
Guerin (n=1)	ND (n=1)/ST358 GB (n=1)/ST508	2002; N 2011; N	Food/P (n=1) Animal production/A (n=1)	Susceptible (n=1) Susceptible (n=1)	(-) (n=1) (-) (n=1)	36/2 (n=1)	0.125/ND (n=1)
Hadar (n=4)	ND (n=4)	2002; N, C, S	Food/Pt (n=4)	AMX, STR, TET, NAL, CIP ^{DS} / <i>bla</i> _{TEM-1} , <i>strA</i> , <i>tet(A)</i> (n=1) TET, STR, NAL, CIP ^{DS} / <i>tet(A)</i> , <i>strA</i> -	(-) (n=4)	32/8 (n=1)	0.16/ND (n=1)

Serotype (no. isolates)	"Clone designation" and PFGE-type (no. isolates)/ST ^a	Isolation date; region ^b	Source/sample (no. isolates) ^c	Antibiotic resistance profile/genes (no. isolates) ^d	Other metal tolerance genes + co-location with antibiotic resistance genes - Chr or PL (Kb, Inc) (no. isolates) ^e	MIC _{range} CuSO ₄ Aerobic/anaerobic (mM) (no. isolates)	MIC _{range} AgNO ₃ Aerobic/with Ag contact (mM) (no. isolates)
				<i>strB</i> (n=1)			
Heidelberg (n=2)	F (n=1)/ST15	2002; S	Food/Pt (n=1)	AMX, NAL/ <i>bla</i> _{TEM-1} (n=1) NAL, CIP ^{DS} /ND (n=1)	<i>merA</i> + <i>bla</i> _{TEM-1} , <i>sul1</i> , <i>tet(A)</i> , <i>dhfrA1</i> , <i>aadA1</i> ^f - <u>PL</u> (120, P) (n=1)	28-32/1-2 (n=2)	0.16-0.25/ND (n=2)
Houston (n=2)	ND (n=1) P (n=2)/ST48	2003; N 2003-2004; N	Human/H (n=1) Food/B (n=1); Human/C (n=1)	Susceptible (n=1) AMX, KAN, STR, SUL, TET, TMP/ <i>bla</i> _{TEM-1} , <i>aphA1</i> , <i>aadA2</i> , <i>sul1</i> - <i>sul2</i> , <i>tet(A)</i> , <i>dhfrA12</i> (n=2)	(-) (n=1) <i>merA</i> + <i>bla</i> _{TEM-1} , <i>aphA1</i> , <i>aadA2</i> , <i>sul1</i> - <i>sul2</i> , <i>tet(A)</i> , <i>dhfrA12</i> - <u>PL</u> (95, UN) (n=1) <i>merA</i> - ND (n=1)	32-36/1-4 (n=2)	0.16 (n=2)/0.125 (n=1)
Indiana (n=1) Infantis (n=3)	ND (n=1) E (n=1)/ST32	2011; S 2003; S	Environment/W (n=1) Food/P (n=1)	Susceptible (n=1) STR, SUL, TET, NAL, CIP ^{DS} / <i>aadA1</i> , <i>sul1</i> , <i>tet(A)</i> (n=1)	(-) (n=1) <i>merA</i> + <i>aadA1</i> , <i>sul1</i> , <i>tet(A)</i> ^f - <u>PL</u> (290, P) (n=1)	32/2 (n=1) 32-36/1-4 (n=2)	0.125/ND (n=1) 0.16/ND (n=2)
Istanbul (n=1) Kubacha (n=1)	ND (n=2) ND (n=1) ND (n=1)	2002-2003; N 2002; S 2002; Unknown	Human/C (n=1) and H (n=1) Food/Pt (n=1) Food/P (n=1)	Susceptible (n=2) NAL, CIP ^{DS} /ND (n=1) Susceptible (n=1)	(-) (n=2)	32/2 (n=1) 36/2 (n=1)	0.125/ND (n=1) 0.125/ND (n=1)
Linguiere (n=1)	LA (n=1)/ST508	2011; C	Animal production/A (n=1)	Susceptible (n=1)	(-) (n=1)	32/1 (n=1)	0.125/ND (n=1)
Livingstone (n=1) Marshall (n=1) Mbandaka (n=5)	ND (n=1) ND (n=1) ND (n=2)	2011; S 2010; S 2002, 2010; C	Food/U (n=1) Human/H (n=1) Food/Pt or U (n=2)	Susceptible (n=1) Susceptible (n=1) Susceptible (n=1)	(-) (n=1) (-) (n=1) (-) (n=2)	32/2 (n=1) 36/2 (n=1) 32/2 (n=1)	0.125/ND (n=1) 0.125/ND (n=1) 0.125/ND (n=1)
Muenchen (n=3)	MA (n=1) MB (n=2) A (n=1)/ST118	2008; N 2008; N 2011; C	Animal production/Pi (n=1) Animal production /Pi (n=2) Animal production/A (n=1)	STR/ <i>strA-strB</i> (n=1) Susceptible (n=2)	(-) (n=1) (-) (n=2)	32/2 (n=1) 32/2 (n=1)	0.125/ND (n=1) 0.125/ND (n=1)
Newport (n=1) Ohio (n=1)	ND (n=1) ND (n=1)	2003, 2010; S 2003; N 2011; S	Human/H (n=2) Human/H (n=1) Human/H (n=1)	Susceptible (n=2) Susceptible (n=1) AMX, KAN, STR, SUL, TET, NAL, CIP ^{DS} / <i>bla</i> _{TEM} , <i>aphA1</i> , <i>strA</i> , <i>sul2</i> , <i>tet(A)</i> (n=1)	(-) (n=2) (-) (n=1) (-) (n=1)	36/2 (n=1) 36/2 (n=1)	0.125/ND (n=1) 0.125/ND (n=1)
Ramsey (n=1)	ND (n=1)	2011; S	Human/H (n=1)	Susceptible (n=1)	(-) (n=1)	36/8 (n=1)	0.125/ND (n=1)

Serotype (no. isolates)	"Clone designation" and PFGE-type (no. isolates) ^a	Isolation date; region ^b	Source/sample (no. isolates) ^c	Antibiotic resistance profile/genes (no. isolates) ^d	Other metal tolerance genes + co-location with antibiotic resistance genes - Chr or PL (Kb, Inc) (no. isolates) ^e	MIC _{range} CuSO ₄ Aerobic/anaerobic (mM) (no. isolates)	MIC _{range} AgNO ₃ Aerobic/with Ag contact (mM) (no. isolates)
Saintpaul (n=3)	I (n=1)/ST27	2002; S, Unknown	Food/Pt (n=3)	AMX, CHL, STR, SUL, TET, TMP/ <i>bla</i> _{TEM-1} , <i>catA</i> , <i>aadA1</i> , <i>sul1</i> - <i>sul2</i> , <i>tet(A)</i> , <i>dhfrA1</i> (n=3)	<i>merA</i> - ND (n=3)	32/1 (n=3)	0.16 (n=3)/0.125 (n=1)
Stanley (n=2)	H (n=2)/ST27						
	ND (n=2)	2011; N	Human/H (n=2)	Susceptible (n=2)	(-) (n=2)	36/2 (n=1)	0.125/ND (n=1)
Teddington (n=1)	ND (n=1)	2002; N	Food/Pt (n=1)	Susceptible (n=1)	(-) (n=1)	32/2 (n=1)	0.125/ND (n=1)
Tompson (n=1)	ND (n=1)	2010; S	Human/H (n=1)	STR/(-) (n=1)	(-) (n=1)	28/1 (n=1)	0.125/ND (n=1)
Uganda (n=1)	ND (n=1)	2002; N	Food/Pt (n=1)	Susceptible (n=1)	(-) (n=1)	32/2 (n=1)	0.125/ND (n=1)
Urbana (n=1)	ND (n=1)	2010; S	Human/H (n=1)	Susceptible (n=1)	(-) (n=1)	36/2 (n=1)	0.125/ND (n=1)
Virchow (n=4)	ND (n=4)	2002-2003; N	Animal production/Pt (n=3); Human/H (n=1)	STR/(-) (n=1) NAL, CIP ^{DS} /ND (n=1) Susceptible (n=2)	(-) (n=4)	36/1 (n=1)	0.125/ND (n=1)
II 4,5:b:- (n=1)	ND (n=1)	2010; S	Food/U (n=1)	STR/(-) (n=1)	(-) (n=1)	32/28 (n=1)	0.125/ND (n=1)
II 42:b:enzx15 (n=1)	ND (n=1)	2003; N	Environment/WR (n=1)	Susceptible (n=1)	(-) (n=1)	32/12 (n=1)	0.125/ND (n=1)
II O4,12:z:1,7 (n=1)	ND (n=1)	2003; N	Food/U (n=1)	Susceptible (n=1)	(-) (n=1)	36/2 (n=1)	0.125/ND (n=1)
IIIa 48:z4 (n=1)	ND (n=1)	2011; S	Environment/W (n=1)	Susceptible (n=1)	(-) (n=1)	36/1 (n=1)	0.125/ND (n=1)
IIIb 50:i:z53 (n=1)	ND (n=1)	2010; S	Human/H (n=1)	STR/(-) (n=1)	(-) (n=1)	28/1 (n=1)	0.125/ND (n=1)

AMX, amoxicillin; CTX, cefotaxime; CHL, chloramphenicol; CIP, ciprofloxacin; CIP^{DS}, decreased susceptibility to ciprofloxacin; GEN, gentamicin; KAN, kanamycin; NAL, nalidixic acid; STR, streptomycin; SUL, sulfamethoxazole; TET, tetracycline; TMP, trimethoprim; Chr, chromosome; PL, plasmid; ND, not determined; UN, untypable.

^a"Clone designation" was adopted only when has been previously published. PFGE-types were designated by capital letters, including previously described^{1,21-24} or firstly designated in this study.

^bN, north; C, center; S, south; I, islands (Azores).

^cA, aquacultures environment; B, beef; C, community; Cw, cow; H, hospitals patients; OT, other type of food; P, pork; Pt, poultry; Pi, pigs and piggeries environment; Pt, poultry; Q, quail; U, unknown; W, bathing or drinking water; WR, river water.

Table S1. Continued

^d(-), Absence of all antibiotic resistance genes searched for. Isolates resistant to three or more antibiotics from different families were considered multidrug-resistant.

^e(-), Absence of all metal tolerance genes searched for. Underlined metal tolerance or antibiotic resistance genes were also detected in transconjugants.

^fIn particular serotypes/clonal lineages *merA*, *arsB* and/or *terF* were in the same genetic platform that the genes belonging to conventional class 1 integrons [1000bp (*aadA1*) in *S. Infantis*; 1000bp (*aadA2*) in *S. Derby*; 1600bp (*aadB-aadA2*) in *S. Bovismorbificans*; 1700bp (*dfrA1-aadA1*) in *S. Enteritidis* and *S. Heidelberg*; 2000bp (*bla_{OXA-30}-aadA1*) in *S. Typhimurium* “OXA-30-producing” clone; 2000bp (*dfrA12-orfF-aadA2*) in *S. Goldcoast* and *S. Houston*] or to atypical *sul3*-type I integrons 7085bp (5’CS-*dfrA12-orfF-aadA2-cmlA1-aadA1-qacH-IS440-sul3*) in *S. Typhimurium* “DT104”/Q clone and *S. Anatum*].²¹⁻²⁴

Table S2. Primers used in PCR assays to search acquired metal tolerance genes

Operon or cluster	Target Gene	Gene coding for	Primer	Primer sequence (5'-3') ^a	Amplicon size (bp)	Annealing temperature (°C)	Nucleotide position (GenBank accession no.)	Reference
<i>sil</i>	<i>silA</i>	Silver inner-membrane proton/cation antiporter	<i>silA_Fw</i> <i>silA_Rv</i>	GCAAGACCGGTAAAGCAGAG CCTGCCAGTACAGGAACCAT	936	59	8936-8955 9871-9852 (<u>AF067954.1</u>)	[6]
	<i>silC</i>	Outer membrane protein	<i>silC_4090_F</i> <i>silC_4470_R</i>	CGGGCTGGCGHAMCTTTTTTG CCAGTTGCTGRTGAAATARC	381	60	4090-4110 4470-4450 (<u>BX664015.1</u>)	This study
	<i>silE</i>	Silver/copper periplasmic binding protein	<i>silE_1105_F</i> <i>silE_1368_R</i>	GTTCGTCATGGTYTCATGAGC GTACTYCCCCGGACATCACTAATT	264	62	1105-1125 1368-1345 (<u>AF067954.1</u>)	[6]
	<i>silP</i>	P-type cation ATPase	<i>silP_11882_F</i> <i>silP_12405_R</i>	GGCGATAAGCTCCGCATCAGA TCCACTTTTTCAAGACGCTCA	524	60	11882-11902 12405-12385 (<u>BX664015.1</u>)	[54]
	<i>silR</i>	Transcriptional regulator responder	<i>silR_3244_F</i> <i>silR_3572_R</i>	CCCTGATGGCGAAGCAAGAA AACGGCTGGGATATCRTCSCG	329	62	3244-3263 3572-3553 (<u>BX664015.1</u>)	[54]
	<i>silS</i>	Membrane kinase sensor	<i>silS_2279_F</i> <i>silS_3019_R</i>	GCGGGTAAARACATCCTCAAT CTGACCTTTTTATCAGCCTG	741	60	2279-2299 3019-2999 (<u>BX664015.1</u>)	[54]
<i>pco</i>	<i>pcoA</i>	Multicopper oxidase	<i>pcoA_978_F</i> <i>pcoA_1481_R</i>	CTCGCGGATGTCAGTGGCTACACCT ATCCGGAAGGTCAGCACCGTCCATAGAC	504	60	978-1002 1481-1454 (<u>X83541.1</u>)	[6]
	<i>pcoD</i>	Copper inner membrane pump	<i>pcoD_F</i> <i>pcoD_R</i>	CTGGCCACACTTGCCTGGGG CAGCTACGGCGCCAGAAAT	500	55	3801-3820 4300-4281 (<u>X83541.1</u>)	[6]
	<i>pcoE</i>	Periplasmic chaperone	<i>pcoE_6585_F</i> <i>pcoE_7057_R</i>	CCTGGTTCTCGAGTGATGA TGACCATATTTCGCCCTTCT	473	58	6585-6604 7057-7039 (<u>X83541.1</u>)	This study
	<i>pcoR</i>	Regulator of <i>pcoABCD</i> cluster	<i>pcoR_4460_F</i> <i>pcoR_4928_R</i>	AGGCTATCAGGCCGATCTCT CCAGACCAGGGACGAGATAA	469	58	4460-4478 4928-4909 (<u>X83541.1</u>)	This study

Table S2. Continued

Operon or cluster	Target Gene	Gene coding for	Primer	Primer sequence (5'-3') ^a	Amplicon size (bp)	Annealing temperature (°C)	Nucleotide position (GenBank accession no.)	Reference
	<i>pcoS</i>	Sensor of <i>pcoABCD</i> cluster	pcoS_5182_F pcoS_6236_R	TAATCAGGACCGCGATTTC CACTGTCAAGCTCAAGGTGT	1055	63	5182-5201 6236-6217 (X83541.1)	This study
<i>mer</i>	<i>merA</i>	Mercuric reductase	merA_1F merA_5R	ACCATCGGCGGCACCTGCGT ACCATCGTCAGGTAGGGGAAC	1238	65	2140-2159 3377-3357 (K03089.1)	[43]
<i>ars</i>	<i>arsB</i>	Arsenite transmembrane pump	arsB_Fw arsB_Rv	AGTGAAGACAGACGAAGCG GGCAGATAGTGTGGAATGCG	1136	60	159735-159754 160870-160851 (BX664015.1)	[55]
<i>ter</i>	<i>terF</i>	Tellurite resistance protein	terF_Fw1 terF_Rv	ATAGCACTGGATCGTGTTCC TTCATCGATCCACGGTCTG	990	60	80174-80193 81163-81145 (BX664015.1)	This study

^aH = A, C or T; K = G or T; M = A or C; R = A or G; S = G or C; Y = C or T.

Additional references:

54. Kremer AN, Hoffmann H. Subtractive hybridization yields a silver resistance determinant unique to nosocomial pathogens in the *Enterobacter cloacae* complex. *J Clin Microbiol* 2012; **50**: 3249-57.
55. Garcia Fernandez A, Cloeckaert A, Bertini A *et al.* Comparative analysis of IncHI2 plasmids carrying *bla*_{CTX-M-2} or *bla*_{CTX-M-9} from *Escherichia coli* and *Salmonella enterica* strains isolated from poultry and humans. *Antimicrob Agents Chemother* 2007; **51**: 4177-80.

Table S3. Dispersion of *sil±pco* gene clusters and their genetic environments in *Salmonella enterica* genomes available at GenBank database

Cu/Ag cluster ^a	Serotype (no. isolates)	Strain name(s)/ST (no. isolates) ^b	Country/Year(s) of isolation	Source	Genetic element associated with <i>sil±pco</i> clusters ^{a,d}		Insertion site related with the genetic element (no. isolates) ^e	GenBank accession number or reference
					Tn6230	Tn6230-partial		
<i>sil</i>	Typhimurium (n=5)	34502, YU39, CH9, CH44/ST213 (n=2)	NR; Canada/2000-2001; Mexico/2005	Swine, children blood	+		Other (n=4)	AUQT01000028.1, CP011429.1, CQHB01000038.1, CQHS01000024.1
		35423/ST213 (n=1)	NR	Cattle			NA (n=1)	(^f)
		CVM N48691, CVM N50445, CVM N51240, CVM N51270, CVM N51287, CVM N51306	USA/2013	Pork chops, ground turkey	+		tRNA-Phe (n=7)	LHKM01000014.1, LHLF01000008.1, LHMA01000018.1, LHMFO1000007.1, LHNH01000009.1, LHNW01000011.1, LHOM01000029.1
	4,12:i:- (n=8)	STM3910	Italy/2011	Human faeces			NA (n=1)	(^f)
<i>sil±pco</i>	I 8,20:-:z6 (n=1)	CVM N43471	USA/2013	Chicken breast			NA (n=1)	(^f)
	Agona (n=2)	CVM N47726	USA/2013	Ground turkey	+		<i>yhiN</i> gene (n=1)	LHJT01000022.1
		21.H.10/ST13 (n=1)	Germany/2010	Human faeces			NA (n=1)	(^f)
	Anatum (n=3)	CFSAN024764	Chile/2009	Human	+		<i>yhiN</i> gene (n=1)	JWQO01000019.1
<i>sil±pco</i>	CVM N44698	CVM N44698	USA/2013	Ground turkey	+		Other (n=1)	LHFG01000020.1
	CVM N44699	CVM N44699	USA/2013	Ground turkey			NA (n=1)	(^f)

Cu/Ag cluster ^a	Serotype (no. isolates)	Strain name(s)/ST (no. isolates) ^b	Country/Year(s) of isolation	Source	Genetic element associated with <i>sil</i> / <i>pco</i> clusters ^{a,d}		Insertion site related with the genetic element (no. isolates) ^e	GenBank accession number or reference
					Tn6230	Tn6230-partial		
Bovismorbificans (n=1)		499208 (n=1)	UK/1970–1980	Alpaca	+		Other (n=1)	CSTH01000028.1
	Brandenburg (n=1)	CVM N45949 (n=1)	USA/2013	Chicken breast	+		Other (n=1)	LHHJ01000009.1
	Cubana (n=4)	CFSAN002050, CFSAN001083, CVM42234/ST286 (n=2)	Philippines/1972; USA/2012	Fresh alfalfa sprouts, dessicated coconut, chick feed		+	tRNA-Phe (n=3)	CP006055.1, APAG01000439.1, ATEU01000022.1
		76814	USA/2004	Swine			NA (n=1)	(^f)
	Derby (n=1)	CVM N51280	USA/2013	Pork chops			NA (n=1)	(^f)
Havana (n=1)	CFSAN024771	Chile/2012	Garuma gull			NA (n=1)	(^f)	
Heidelberg (n=27)		SH111_227, 24393, SARA 39, CVM N44695, CVM N44710, CVM N45397, CVM N45951, CVM N45956, CVM N46812, CVM N48578, CVM N48699, CVM N50450, CVM N51244	USA/2001, 2003, 2013	Bovine, turkey intestines, human, chicken wings, pork chops, ground turkey, ground beef, chicken breast	+		Other (n=13)	[36, 56], LHFD01000022.1, LHFQ01000012.1, LHGB01000032.1, LHHL01000045.1, LHHQ01000047.1, LHHU01000028.1, LHJY01000019.1, LHKU01000049.1, LHME01000058.1, LHMI01000035.1
		CFSAN002064, CFSAN002069, CVM N45958, CVM N46811,	USA/2012, 2013	Human stool, chicken sample, chicken breast,	+		<i>yhiN</i> gene (n=5)	CP005995.1, CP005390.2, LHHS01000023.1,

Cu/Ag cluster ^a	Serotype (no. isolates)	Strain name(s)/ST (no. isolates) ^b	Country/Year(s) of isolation	Source	Genetic element associated with <i>sil</i> / <i>pc</i> clusters ^{a,d}		Insertion site related with the genetic element (no. isolates) ^e	GenBank accession number or reference
					Tn6230	Tn6230-partial		
		CVM N50449/ST15 (n=1)		chicken wings				LHHT01000004.1, LHMD01000005.1
		20752, 32507, 24359, N15757, N30678, 41565, CVM N44707, CVM N51243, CVM N51984	USA/2002, 2003, 2007, 2011, 2013	Turkey cecal tonsil, ground turkey, turkey intestines, chicken breast, pork chops			NA (n=9)	(^f)
	Infantis (n=3)	CVM N44705, Sal147, Sal280	Portugal/2011, 2012; USA/2013	Chicken breast, food sample			NA (n=3)	(^f)
	Kentucky (n=21)	CVM N47723, CVM N45934, CVM N45939, CVM N45944, CVM N46849, CVM N48705, CVM N48710	USA/2013	Ground turkey, chicken breast	+		Other (n=1), NA (n=6)	LHJQ01000057.1, LHGX01000055.1, LHHC01000025.1, LHHH01000045.1, LHIU01000057.1, LINU01000071.1, LHLB01000019.1
		N312, 20793, CVM N43447, CVM N47720, CVM N50429, CVM N51981/ST152 (n=2)	USA/2002, 2004, 2013	Chicken breast		+	Other (n=6)	AUQI01000016.1, AUQJ01000021.1, LHDL01000020.1, LHJO01000031.1, LHLN01000030.1, LHOT01000028.1
		SALC-205-3, CVM N444693, CVM N45412, CVM N45937, CVM N47729, CVM N47730, CVM N48687, CVM N51241	USA/2013; Canada/2004	Chicken breast, chicken fecal material, chicken wings			NA (n=8)	(^f)

Cu/Ag cluster ^a	Serotype (no. isolates)	Strain name(s)/ST (no. isolates) ^b	Country/Year(s) of isolation	Source	Genetic element associated with <i>sil</i> / <i>pc</i> clusters ^{a,d}		Insertion site related with the genetic element (no. isolates) ^e	GenBank accession number or reference
					Tn6230	Tn6230-partial		
Liverpool (n=1)		CVM N50444	USA/2013	Ground turkey	+		Other (n=1)	LHLZ01000022.1
Mbandaka (n=1)		CVM N51302	USA/2013	Chicken breast	+		<i>yhiN</i> gene (n=1)	LHOI01000026.1
Montevideo (n=45)		S5-403, 315996572, 609458-1, 495297-1, 495297-3, 495297-4, 515920-1, 515920-2, 609460, NC_MB110209-0054, OH_2009072675, CASC_09SCPH15965, 19N, 81038-01, MD_MDA09249507, 414877, 366867, 413180, 446600, 556152 , 556150-1, MB101509-0077, MB102109-0047, MB111609-0052, 2009083312, 2009085258, 315731156, IA_2009159199, IA_2010008282, IA_2010008283, IA_2010008284, IA_2010008287, CT_02035278, CT_02035318, 80959-06, CT_02035320, CT_02035321, IA_2010008286, 507440-20, MB110209-0055,	USA/NR, 2006, 2007, 2009, 2010, 2013	Pistachio ^c , black peper ^c , human stool, chicken ^c , hearts of romaine ^c , human sample, mozzarella ^c , perch ^c , sea trout ^c , king fish ^c , red pepper flakes ^c , lunch meat ^c , environmental isolate ^c , environmental swab ^c , red pepper, chcken breast	+		[40], AESH01000033.1, AETF01000076.1, AESI01000042.1, AESJ01000015.1, AESK01000007.1, AESL01000025.1, AESM01000016.1, AETH01000090.1, AESO01000068.1, AESP01000033.1, AESQ01000058.1, AESV01000032.1, AESZ01000024.1, AETA01000018.1, AETB01000005.1, AETC01000011.1, AETD01000010.1, AETE01000043.1, AETJ01000070.1, AETG01000080.1, AETK01000018.1, AETL01000127.1, AETN01000032.1, AETO01000025.1, AETP01000034.1, AETQ01000024.1,	

Cu/Ag cluster ^a	Serotype (no. isolates)	Strain name(s)/ST (no. isolates) ^b	Country/Year(s) of isolation	Source	Genetic element associated with <i>slt</i> / <i>pc</i> clusters ^{a,d}		Insertion site related with the genetic element (no. isolates) ^e	GenBank accession number or reference
					Tn6230	Tn6230-partial		
		FCC0146, CVM						AETR01000099.1,
		N51288/ST316 (n=38);						AETS01000072.1,
		4441 H/ST138 (n=1)						AETT01000046.1,
								AETU01000051.1,
								AETW01000065.1,
								AHHS01000110.1,
								AHHT01000019.1,
								AHHR01000012.1,
								AHHU01000003.1,
								AHHR01000012.1,
								AHIK01000009.1,
								CP007530.1,
								AETM01000157.1,
								JSWT01000001.1,
								LHNX01000007.1,
								AESY01000016.1
		IA_2010008285, CT_02035327/ST316 (n=2)	USA/NR, 2010	Lunch meat ^c , environmental isolate ^c			NA (n=2)	(ⁱ)
Ohio (n=1)		CFSAN001079/ST329 (n=1)	USA/NR	Pork sausage	+		tRNA-Phe (n=1)	APAK01000322.1
Rissen (n=1)		150	USA/2009	Food from white pepper outbreak	+		Other (n=1)	AHUI01000058.1
Schwarzengrund (n=15)		CFSAN003382, CVM N48682, CVM N51311	USA/2011, 2013	Creek water, ground turkey, chicken breast	+		<i>yhiN</i> gene (n=3)	LARZ01000012.1, LHKE01000010.1, LHOP01000021.1
		CVM N43459, CVM	USA/2013	Chicken wings,	+		Other (n=6)	LHDW01000030.1,

Cu/Ag cluster ^a	Serotype (no. isolates)	Strain name(s)/ST (no. isolates) ^b	Country/Year(s) of isolation	Source	Genetic element associated with <i>sil</i> / <i>apco</i> clusters ^{a,d}		Insertion site related with the genetic element (no. isolates) ^e	GenBank accession number or reference
					Tn6230	Tn6230-partial		
	Senftenberg (n=21)	N43479, CVM N44700, CVM N50434, CVM N51257, CVM N51289		chicken breast				LHEJ01000002.1, LHF010000042.1, LHLQ01000036.1, LHMV01000011.1, LHN01000009.1
		CVM N45952, CVM N47711, CVM N47712, CVM N51259, CVM N51267	USA/2013	Ground turkey, ground breast, chicken breast		+	tRNA-Phe (n=5)	LHMM01000008.1, LHJG01000015.1, LINN01000004.1, LHM01000017.1, LHN01000001.1
		CVM N47722	USA/2013	Ground turkey	+		Other (n=1)	LINO01000043.1
		A4-543, SS209, NC_MB012510-0038, 604314, 423984-2, CFSAN024718, CFSAN024719, CFSAN024720, CFSAN024721, CFSAN024722, CFSAN024723, CFSAN024724, NCTC10384, CVM N43462, CVM N51274, CVM N51312/ST14 (n=4)	NR/1964; USA/NR, 2005, 2006, 2010, 2013; France/NR; Chile/2009-2013	Cattle, gastroenteritis, human stool, black pepper, salame-soprasetta, kelp gull, human sample, faeces, chicken wings, ground turkey	+		<i>yhiN</i> gene (n=16)	[40], CAGQ01000054.1, AYEJ01000033.1, AOXW01000018.1, AOYP01000072.1, JWRE01000006.1, JWRD01000007.1, JWRC01000002.1, JWRB01000009.1, JWRA01000005.1, JWQZ01000006.1, JWQY01000030.1, LN868943.1, LHDZ01000008.1, LHNK01000004.1, LHOQ01000023.1

Cu/Ag cluster ^a	Serotype (no. isolates)	Strain name(s)/ST (no. isolates) ^b	Country/Year(s) of isolation	Source	Genetic element associated with <i>sil±pco</i> clusters ^{a,d}		Insertion site related with the genetic element (no. isolates) ^e	GenBank accession number or reference
					Tn6230	Tn6230-partial		
		CVM N51281	USA/2013	Ground turkey	+		Other (n=1)	LHNQ01000022.1
		ATCC 8400/ST290 (n=1)	USA/NR	NR		+	tRNA-Phe (n=1)	AOXU01000003.1
		423984-1, 316235162, 361154004/ST14 (n=3)	USA/NR, 2009	Environmental swab ^c , pistachios, shelled pistachios			NA (n=3)	(ⁱ)
	Typhimurium (n=10)	S7, S15, S23, ST4020, ST4024, Sal368	Denmark/NR; Hong Kong/2007; Portugal/ 2012	Food, copper feed pigs, human blood		+	tRNA-Phe (n=6)	[41], LFGX01000049.1, JRYU01000005.1, JRYU01000005.1
		MG101, ST2533, ST2286, FSAN236CA	USA/1973; Hong Kong/2006; Colombia/2012	Human burn, human stool; animal feed, raw chicken carcass at slaughterhouse	+		Other (n=2), NA (n=2)	[57, 58], JRZP01000030.1, JRZR01000027.1, LKIP01000046.1
	Tennessee (n=2)	TXSC_TXSC08-19/ST319 (n=1)	USA/2004	Fishmeal	+		<i>yhiN</i> gene (n=1)	CP007505.1
		TXSC_TXSC08-21/ST319 (n=1)	New Zealand/2004	Meat and bone meal, lamb-based			NA (n=1)	(ⁱ)
	Worthington (n=4)	CVM N45406, CVM N45931, CVM N46835, CVM N51304	USA/2013	Chicken breast, ground turkey		+	tRNA-Phe (n=4)	LHGK01000008.1, LHGV01000004.1, LHIL01000001.1, LHOK01000015.1

NR, not reported

Table S3. Continued

^aAnalysis of *sil*/ \pm *pco* gene clusters dispersion as well as their genetic environments was performed in completed *Salmonella* genomes, plasmid sequences or contigs (when visualization of the genetic environment was possible) available for Microbial Genome BLAST[®] (<http://blast.ncbi.nlm.nih.gov/Blast.cgi>; accessed 19/10/2015). The sequences were searched using amino acidic sequences (blastp) of *pcoA* and *pcoD* from pRJ1004 (accession no. CAA58528.1)²⁸ and *silA* and *silE* from pMG101 (accession no. AAD11749.1)⁵⁸ as query, and those with a minimum of 97% amino acidic identity were selected. What we consider in the manuscript as a common *sil*/ \pm *pco* genetic module corresponds to part of Tn6230 (*sil*/ \pm *pco* gene clusters and genes coding for a hypothetical protein and a total/partial endonuclease; e.g. GenBank accession number AOXW01000018.1 from 180790 bp to 157014 bp), was based on the synteny of those genes and not in the differences of the intergenic regions (variability observed between \approx 21 and 24 Kb), or small nucleotide variability within each gene.

^bNumber of isolates with sequence types (STs) previously assigned, according to the *Salmonella* MLST scheme (<http://mlst.warwick.ac.uk/mlst/dbs/Senterica>) and available in NCBI (<http://www.ncbi.nlm.nih.gov/>) or PATRIC (<https://www.patricbrc.org/portal/portal/patric/Home>) databases.

^cIsolates with different source information among databases. The information here considered was that available in BioSample NCBI (<http://www.ncbi.nlm.nih.gov/biosample/>).

^dTwo types of genetic environments were found associated with the common *sil*/ \pm *pco* module and marked with “+” when they are present in the analysed strains (n=174): “Tn6230”, when the typical proteins (TnsABC) involved in transposition of these genetic elements are present or “Tn6230-partial” corresponding to the common module but without the TnsABC proteins. The absence of “+” means that it was not possible to evaluate if the genetic element was complete or partial due to the interruption of the contig.

^eInsertion sites related to Tn6230 or Tn6230-partial genetic elements: “*yhiN* gene” – common insertion site of Tn6230; “tRNA-phe” – consists of the common tRNA-Phe + *xerD* recombinase + relaxase + DNA helicase region proximal to Tn6230-partial element; “Others” – Tn6230 or Tn6230-partial were found in other insertion sites beside *yhiN* gene or tRNA-phe region; “NA”, not applicable when it was not possible to evaluate the insertion site of the genetic element due to the interruption of the contig.

^fGenBank accession numbers of isolates with interrupted contigs. *Salmonella* serotypes with *sil* cluster: Typhimurium (AUQU01000026.1) and with *sil*/ \pm *pco* clusters: 4,12:i:- (LJJK01000035.1), I 8,20:-:z6 (LHEE01000083.1), Agona (CASU01000164.1), Anatum (LHFH01000016.1), Cubana (AZGR01000012.1), Derby (LHNP01000052.1), Havana (JWQI01000012.1), Heidelberg

Table S3. *Continued*

(AMNM01000010.1, AMNL01000012.1, AMNP01000117.1, AMND01000026.1, AMNK01000100.1, AJHA01000116.1, LHFN01000150.1, LIOA01000049.1, LHOW01000029.1), Infantis (LHFL01000049.1, LFGU01000369.1, LFGW01000003.1), Kentucky (Dhanani *et al.*, 2015, LIMQ01000067.1, LHGN01000037.1, LHHA01000088.1, LHJV01000024.1, LHJW01000021.1, LINR01000077.1, LHMG01000006.1), Montevideo (AETV01000009.1, AHHW01000039.1), Senftenberg (AYDP01000247.1, AOYU01000007.1, AYDO01000054.1), Tennessee (AOXQ01000059.1, AOXQ01000257.1).

Additional references:

56. Han J, Lynne AM, David DE *et al.* DNA sequence analysis of plasmids from multidrug resistant *Salmonella enterica* serotype Heidelberg isolates. *PLoS One* 2012; **7**: e51160.
57. Gupta A, Matsui K, Lo JF *et al.* Molecular basis for resistance to silver cations in *Salmonella*. *Nat Med* 1999; **5**: 183-8.
58. Randall CP, Gupta A, Jackson N *et al.* Silver resistance in Gram-negative bacteria: a dissection of endogenous and exogenous mechanisms. *J Antimicrob Chemother* 2015; **70**: 1037-46.

***Salmonella enterica* serotype Bovismorbificans, a new host for
CTX-M-9**

Patrícia Antunes^{1,2}, Joana Mourão², Tatiana Alves², Joana Campos², Carla Novais², Ângela
Novais² and Luísa Peixe^{2*}

¹ Faculdade de Ciências da Nutrição e Alimentação, Universidade do Porto, Porto, Portugal; ² REQUIMTE, Laboratório de Microbiologia, Faculdade de Farmácia, Universidade do Porto, Porto, Portugal.

International Journal of Antimicrobial Agents 2013 January; 41(1):91-93

Elsevier has authorized the reproduction of the final published PDF version of this paper in this thesis through the License Agreement Number 3791960645341 on January 18, 2016.

Table 1
Features of the carbapenem-resistant isolates and patient characteristics.

Patient	Hospitalisation unit	Isolate	Site of isolation	Carbapenemase	Associated β -lactamases	MLST type	MICs of carbapenems		
							IPM	MER	ERT
1	Oncology	<i>Klebsiella pneumoniae</i>	Stool	OXA-48	None	ST14	2	0.75	1.5
2	Outpatient	<i>Escherichia coli</i>	Pus	OXA-48	CTX-M-14, TEM-1, SHV-12	ST38	0.75	0.25	1.5
3	Oncology	<i>E. coli</i>	Blood	OXA-48+VIM-1	None	ST101	32	>32	>32
4	Bone marrow transplantation unit	<i>K. pneumoniae</i>	Throat swab	VIM-1	CTX-M-14, TEM, SHV-12	ST17	1.5	0.5	0.75
5	Oncology	<i>K. pneumoniae</i>	Urine	VIM-1	CTX-M-15	ST101	32	>32	>32
6	Oncology	<i>Enterobacter cloacae</i>	Pus	VIM-1	None	N/D	8	0.5	0.25
7	Paediatric oncology	<i>E. cloacae</i>	Peritoneal fluid	VIM-1	SHV-12	N/D	32	32	16
8	Oncology	<i>E. cloacae</i>	Pus	VIM-1	None	N/D	1	0.5	1
9	Oncology	<i>E. coli</i>	Pus	VIM-1	TEM-1, SHV-12	ST23	1	0.5	0.5
10	Paediatric oncology	<i>E. cloacae</i>	Blood	VIM-1	CTX-M-9, TEM-1, SHV-11	N/D	1.5	0.5	0.5

MLST, multilocus sequencing typing; MIC, minimum inhibitory concentration; IPM, imipenem; MER, meropenem; ERT, ertapenem; N/D, not determinable.

References

- [1] Nordmann P, Naas T, Poirel L. Global spread of carbapenemase-producing Enterobacteriaceae. *Emerg Infect Dis* 2011;17:1791–8.
- [2] Paterson DL, Bonomo RA. Extended-spectrum β -lactamases: a clinical update. *Clin Microbiol Rev* 2005;18:657–86.
- [3] Diancourt L, Passet V, Verhoef J, Grimont PA, Brisse S. Multilocus sequence typing of *Klebsiella pneumoniae* nosocomial isolates. *J Clin Microbiol* 2005;43:4178–82.
- [4] Poirel L, Bonnin RA, Nordmann P. Genetic features of the widespread plasmid coding for the carbapenemase OXA-48. *Antimicrob Agents Chemother* 2012;56:559–62.
- [5] Carrère A, Poirel L, Yilmaz M, Akan OA, Feriha C, Cuzon G, et al. Spread of OXA-48-encoding plasmid in Turkey and beyond. *Antimicrob Agent Chemother* 2010;54:1369–73.

Laurent Poirel*

Service de Bactériologie-Virologie, INSERM U914 'Emerging Resistance to Antibiotics', Hôpital de Bicêtre, Assistance Publique/Hôpitaux de Paris, Faculté de Médecine Paris Sud, K.-Bicêtre, France

Mohammed O. Abdelaziz

Department of Microbiology and Immunology, Faculty of Pharmacy, Helwan University, Cairo, Egypt

Sandrine Bernabeu

Patrice Nordmann

Service de Bactériologie-Virologie, INSERM U914 'Emerging Resistance to Antibiotics', Hôpital de Bicêtre, Assistance Publique/Hôpitaux de Paris, Faculté de Médecine Paris Sud, K.-Bicêtre, France

*Corresponding author. Present address: Service de Bactériologie-Virologie, Hôpital de Bicêtre, 78 rue du Général Leclerc, 94275 Le Kremlin-Bicêtre Cedex, France.
Tel.: +33 1 45 21 3632; fax: +33 1 45 21 63 40.
E-mail address: laurent.poirel@bct.aphp.fr (L. Poirel)

24 August 2012

doi:10.1016/j.ijantimicag.2012.08.015

***Salmonella enterica* serotype Bovismorbificans, a new host for CTX-M-9**

Sir,

The spread of non-typhoidal *Salmonella* resistant to broad-spectrum cephalosporins constitutes an emerging threat worldwide, although their distribution is uneven in different geographical regions and ecological niches [1]. Although production of AmpC-type enzymes, mainly CMY-2, is rising in Europe, extended-spectrum β -lactamases (ESBLs) are still the most associated with resistance to cephalosporins [1]. Different food-producing animals and food carry ESBL-producing bacteria, with CTX-M being the most frequent family [1]. The CTX-M-9 enzyme is one of the most frequently found in *Salmonella* both of human and animal origin [1], linked to the clonal spread of a particular *Salmonella* serotype (*Salmonella enterica* serotype Virchow) and food-producing animals (poultry) in neighbouring countries (Spain and France) [1,2]. In this study, we describe the molecular characterisation of the first ESBL-producing *Salmonella* in Portugal, consisting of *bla*_{CTX-M-9} located in an In60-like structure and within IncHI2 plasmid, carried by an emerging human *Salmonella* serotype, *S. enterica* serotype Bovismorbificans.

Susceptibility to antibiotics as well as β -lactamase production was assayed by disc diffusion and/or Etest [Clinical and Laboratory Standards Institute (CLSI) documents M2-A10/M100-21]. Screening of genes encoding resistance to antibiotics or biocides found in the animal setting (feed, disinfectants and environmental pollution) was performed by PCR and sequencing as described




Fig. 1. (a) Genetic context of *bla*_{CTX-M-9} within an In60 variant identified in a *Salmonella enterica* serotype Bovismorbificans isolate (HP507391). Differences compared with the In60 of *S. enterica* serotype Virchow [2] are the presence of *int11* (A), the presence of *aadB* instead of *dfrA16* (B) and the absence of insertion sequence IS3000 (C). The discontinued arrows indicate genes not found. The grey arrows indicate genes not determined. (b) Antimicrobial susceptibility profiles and resistance genotypes of *S. Bovismorbificans* carrying *bla*_{CTX-M-9} (HP507391) and the corresponding transconjugant (conjugation assays using *Escherichia coli* K802N) (HP507391.T). TGC, tigecycline; CIP, ciprofloxacin; AMX, amoxicillin; AMC, amoxicillin/clavulanic acid; TIC, ticarcillin; TIM, ticarcillin/clavulanic acid; CEF, cefalothin; FOX, cefoxitin; CTX, cefotaxime; CAZ, ceftazidime; FEP, cefepime; CR, ceftipime; IPM, imipenem; MEM, meropenem; ERT, ertapenem; ATM, aztreonam; GEN, gentamicin; KAN, kanamycin; STR, streptomycin; SUL, sulfamethoxazole; TOB, tobramycin; NAL, nalidixic acid.

previously [3]. Transferability of *bla* genes and characterisation of integron and plasmid backbones was performed by PCR, hybridisation (I-CeuI/S1 nuclease) and/or sequencing [4]. The genetic environment of *bla*_{CTX-M-9} (class 1 integron, Tn402 derivatives) was characterised by PCR mapping based on known sequences [4]. Clonal identification was performed by multilocus sequence typing (MLST) (<http://mlst.ucc.ie/mlst/dbs/Senterica>).

An *S. enterica* isolate carrying the *bla*_{CTX-M-9} gene was isolated from the faeces of a 4-year-old boy admitted to the emergency ward of a Portuguese hospital with gastrointestinal infection in February 2011. The isolate was identified as *S. Bovismorbificans* and was assigned to sequence type 142 (ST142), previously associated with food-borne and human European isolates (<http://mlst.ucc.ie/mlst/dbs/Senterica>). *Salmonella* Bovismorbificans is an unusual serotype but it is increasingly reported in Northern European countries where it has been linked to severe outbreaks traced mostly to vegetables [5]. However, it was not possible to associate the isolate reported here with consumption of a particular food products or animal contact or foreign travel. The isolate was resistant to ampicillin, cefalothin and cefotaxime, but not to ceftazidime, cefepime, aztreonam or carbapenems (Fig. 1). Resistance to gentamicin, kanamycin, tobramycin, streptomycin and sulfamethoxazole was also detected, but it was not expressed in the transconjugant (with the exception of sulfamethoxazole) irrespective of the presence of the corresponding genes (Fig. 1).

The *bla*_{CTX-M-9} gene was associated with an unusual class 1 integron containing *aadB*–*aadA2* gene cassettes and with the insertion sequence *ISCR1* upstream of *bla* but lacking *IS3000* downstream (GenBank accession no JX026665; Fig. 1). This novel In60 variant differs from the previously described In60 integrons in the gene cassette array and also in the absence of *IS3000* and Tn402 derivatives sequences downstream of *bla*_{CTX-M-9}. In Europe, the *bla*_{CTX-M-9} gene was identified in a variety of *ISCR1*-bearing class 1 integrons among Enterobacteriaceae [2,4], although in *Salmonella* it was usually linked to a conserved In60 backbone carrying *dfrA16*–*aadA2* disseminated among humans and poultry and associated with a particular clone of *S. Virchow* [2,3].

The genetic platform bearing *bla*_{CTX-M-9} was located on a transferable 240 kb IncHI2 plasmid carrying a *rep* gene with 99%

identity with that of the prototype R478 plasmid (GenBank accession no. NC005211). The spread of CTX-M-9-producing bacteria both of human and animal origin has also been linked to highly related R478–IncHI2 multidrug-resistant (antibiotic/biocides) plasmids [3,4]. As in the case of R478, *arsB* (arsenic) and *terF* (tellurium) genes were detected, but not *merA* (mercury), *silA* (silver/copper) and *copD* (copper) resistance genes. The variability detected in the plasmid content and In60 platform compared with those previously described suggests the occurrence of multiple recombination events as reported previously [3,4]. The concomitant presence of genes coding for resistance to antibiotics and metals found in the animal setting could contribute to the successful spread of the IncHI2 plasmid family by different events of co-selection. In fact, these multidrug-resistant plasmids are the fifth most common plasmid family occurring in Enterobacteriaceae and have been associated with the dissemination of diverse clinically relevant genes, hence facilitating their increasing prevalence and persistence in different ecological niches.

This study corresponds to the first description of an ESBL-producing *Salmonella* strain in Portugal and the first description of a CTX-M-9 enzyme in the emergent serotype, *S. Bovismorbificans*. The emergence of *bla*_{CTX-M-9} in a similar genetic environment (In60 variant, IncHI2 plasmid) to those circulating in other *Salmonella* serotypes and *Escherichia coli* from humans and animals in Europe suggests an ongoing spread and evolution of this plasmid in different/new hosts and probably under different selective pressures, requiring surveillance to contain its further dissemination.

Acknowledgments

The authors are very grateful to Dr Fernando Fonseca (Centro Hospitalar Póvoa de Varzim/Vila do Conde, Póvoa de Varzim, Portugal) and to Centro Nacional de *Salmonella* (Lisbon, Portugal) for the serotyping.

Funding: This work was supported by Fundação para a Ciência e a Tecnologia (FCT) (grant no. PEst-C/EQB/LA0006/2011). J.M. was supported by a Fellowship from FCT (SFRH/BD/77518/2011), and A.N. by a Marie Curie Intra European Fellowship within the 7th

European Community Framework Programme (ref. PIEF-GA-2009-255512).

Competing interests: None declared.

Ethical approval: Not required.

References

- [1] EFSA Panel on Biological Hazards (BIOHAZ). Scientific opinion on the public health risks of bacterial strains producing extended-spectrum β -lactamases and/or AmpC β -lactamases in food and food-producing animals. EFSA J 2011;9:2322.
- [2] Riaño I, García-Campello M, Sáenz Y, Alvarez P, Vinué L, Lantero M, et al. Occurrence of extended-spectrum β -lactamase-producing *Salmonella enterica* in northern Spain with evidence of CTX-M-9 clonal spread among animals and humans. Clin Microbiol Infect 2009;15:292–5.
- [3] García Fernández A, Cloeckaert A, Bertini A, Praud K, Doublet B, Weill FX, et al. Comparative analysis of IncHI2 plasmids carrying *bla*_{CTX-M-2} or *bla*_{CTX-M-9} from *Escherichia coli* and *Salmonella enterica* strains isolated from poultry and humans. Antimicrob Agents Chemother 2007;51:4177–80.
- [4] Novais A, Cantón R, Valverde A, Machado E, Galán JC, Peixe L, et al. Dissemination and persistence of *bla*_{CTX-M-9} are linked to class 1 integrons containing CR1 associated with defective transposon derivatives from Tn402 located in early antibiotic resistance plasmids of IncHI2, IncP1- α , and IncFI groups. Antimicrob Agents Chemother 2006;50:2741–50.
- [5] Rimhanen-Finne R, Niskanen T, Lienemann T, Johansson T, Sjöman M, Korhonen T, et al. A nationwide outbreak of *Salmonella* Bovismorbificans associated with sprouted alfalfa seeds in Finland, 2009. Zoonoses Public Health 2011;58:589–96.

Patrícia Antunes^{a,b,*}

^a Faculdade de Ciências da Nutrição e Alimentação, Universidade do Porto, Rua Dr Roberto Frias, 4200 Porto, Portugal

^b REQUIMTE, Laboratório de Microbiologia, Faculdade de Farmácia, Universidade do Porto, Rua Jorge Viterbo Ferreira no 228, 4050-313 Porto, Portugal

Joana Mourão

Tatiana Alves

Joana Campos

Carla Novais

Ângela Novais

Luísa Peixe

REQUIMTE, Laboratório de Microbiologia, Faculdade de Farmácia, Universidade do Porto, Rua Jorge Viterbo Ferreira no 228, 4050-313 Porto, Portugal

*Corresponding author. Tel.: +351 22 507 4320;
fax: +351 22 507 4329.

E-mail addresses: patriciaantunes@fcna.up.pt,
patantunes@gmail.com (P. Antunes)

13 July 2012

doi:10.1016/j.ijantimicag.2012.09.009

First report of *bla*_{PER-3} in *Acinetobacter baumannii*

Sir,

Acinetobacter baumannii is an important opportunistic nosocomial pathogen with a remarkable ability to survive and disseminate. In addition, *A. baumannii* can acquire new antibiotic resistance genes and upregulate them, making it a problematic pathogen causing numerous outbreaks worldwide [1,2]. Patients at high risk for developing *A. baumannii* infections are those with prolonged hospitalisation and immune suppression, such as cancer patients [2]. Antibiotic selective pressure also contributes to the rising incidence of resistance [1].

The mechanisms of resistance in *A. baumannii* that cause most concern are the β -lactamases, and all the Ambler class enzymes (A–D) have been found in this species. Class D and class B β -lactamases mainly confer resistance to the carbapenems, whilst

class A are mainly extended-spectrum β -lactamases (ESBLs) [1]. The first report of an ESBL in *A. baumannii* was *bla*_{PER-1} in France [3], with PER-like variants in *A. baumannii* now increasingly reported worldwide. PER-3 was initially characterised from an *Aeromonas punctata* in France (GenBank accession no. AY740681.1) and recently in a medical centre in Taiwan [3], sharing 99% homology to *bla*_{PER-1}. Here we report the first identification of a *bla*_{PER-3}-producing *A. baumannii* isolate.

Isolate AB-15094 was obtained from a 9-year-old patient with osteosarcoma in 2010 at the Children's Cancer Hospital 57357 (Cairo, Egypt). Phenotypic identification was followed by sequencing of the *bla*_{OXA-51-like} gene and restriction analysis of the 16S–23S rRNA spacer sequence [2,4].

Minimum inhibitory concentrations (MICs) were determined and interpreted according to British Society for Antimicrobial Chemotherapy (BSAC) guidelines for imipenem, meropenem, ceftazidime and rifampicin. Detection of the OXA-type carbapenemases *bla*_{OXA-23}, *bla*_{OXA-24}, *bla*_{OXA-51} and *bla*_{OXA-58} as well as class A β -lactamase genes *bla*_{PER-like}, *bla*_{VEB}, *bla*_{GES} and *bla*_{TEM} and the class C β -lactamase gene *bla*_{ADC} was done by standard PCR [3,4]. The class 1 integron harbouring the *bla*_{PER-3} gene was analysed by standard PCR and sequencing methods [4]. S1 nuclease digestion was performed to detect and size plasmids [3]. To investigate whether *bla*_{PER-3} was plasmid-borne, a plasmid curing experiment was conducted. Susceptibility to ceftazidime was examined after curing and was compared with the initial susceptibility of the uncured strain.

Isolate AB-15094 was confirmed as *A. baumannii* harbouring the *bla*_{OXA-65} gene. It was sensitive to imipenem and meropenem but resistant to ceftazidime and rifampicin, with MICs of 64 mg/L and 256 mg/L respectively. The isolate was positive for *bla*_{PER-3} as revealed by sequencing. The strain was negative for all other ESBLs tested as well as for the acquired class D carbapenemases. No insertion sequence (IS) element was detected upstream of the *bla*_{ADC} gene, suggesting no overexpression of the gene [4].

There was no hybridisation of *bla*_{PER-3}-specific probes with any of the isolate's three plasmids (60, 70 and 145.5 kb, respectively), suggesting chromosomal localisation of the gene. This was supported by the lack of change in ceftazidime susceptibility after 10 days of plasmid curing. Furthermore, attempted conjugal transfer was also negative. Sequencing upstream of *bla*_{PER-3} revealed a complex class 1 integron structure harbouring *sul1*, *qacE Δ 1* in the variable 3'CS and ORF513 directly upstream of the *bla*_{PER-3} gene, serving as the transcriptional promoter. Fig. 1 shows the schematic representation of the genetic environment of *bla*_{PER-3}.

This isolate was obtained from a paediatric cancer patient in Egypt and is the first identification of *bla*_{PER-3} in a clinical *A. baumannii* isolate. Previous reports of this gene have been from *A. punctata* in France and Taiwan where the *bla*_{PER-3} gene was reported to be localised both on plasmids and on the chromosome [3]; however, in this study it was located on the chromosome alone. The genetic environment upstream of *bla*_{PER-3} was similar to the *A. punctata* isolate identified in France (GenBank accession no. AY740681.1), being encompassed in a complex class 1 integron. The upstream regions of *bla*_{PER-3} and *bla*_{PER-1} differ significantly despite their structural homology. *bla*_{PER-1} was part of transposon-related structure Tn1213 consisting of ISPa12 and ISPa13 flanking the gene [1]. The *bla*_{PER-3} in this study lacks these IS elements and harbours ISCR1, which is closely associated with class A ESBLs and facilitates the movement of these resistance genes [3,5].

PER-3 is a point-mutant derivative of PER-1 [3] harbouring the change Histidine134Leucine. This mutation is positioned in the H4 helix of the secondary structure of the Ω -loop region, which is associated with the active site of class A β -lactamases [5]. The observed mutation could explain the reduced resistance to ceftazidime (MIC = 64 mg/L) in isolate AB-15094 harbouring *bla*_{PER-3}

Clinical *Salmonella* Typhimurium ST34 with metal tolerance genes and an IncHI2 plasmid carrying *oqxAB-aac(6')-Ib-cr* from Europe

Joana Campos^{1†}, Joana Mourão^{1†}, Sara Marçal², Jorge Machado³, Carla Novais¹, Luísa Peixe¹ and Patrícia Antunes^{1,2*}

¹ UCIBIO/REQUIMTE, Departamento de Ciências Biológicas, Laboratório de Microbiologia, Faculdade de Farmácia, Universidade do Porto, Porto, Portugal; ² Faculdade de Ciências da Nutrição e Alimentação, Universidade do Porto, Porto, Portugal; ³ Laboratório Nacional de Referência de Infecções Gastrointestinais, Departamento de Doenças Infeciosas, Instituto Nacional de Saúde Dr. Ricardo Jorge, Lisboa, Portugal.

†These authors contributed equally to the experimental work.

Journal of Antimicrobial Chemotherapy 2015 December; [Epub ahead of print]

Oxford University Press has authorized the reproduction of the final published PDF version of this paper in this thesis through the License Agreement Number 3787960688972 on January 15, 2016.

J Antimicrob Chemother
doi:10.1093/jac/dkv409

Clinical *Salmonella* Typhimurium ST34 with metal tolerance genes and an IncHI2 plasmid carrying *oqxAB-aac(6')-Ib-cr* from Europe

Joana Campos^{1†}, Joana Mourão^{1†}, Sara Marçal², Jorge Machado³, Carla Novais¹, Luísa Peixe¹ and Patrícia Antunes^{1,2*}

¹UCIBIO/REQUIMTE, Departamento de Ciências Biológicas, Laboratório de Microbiologia, Faculdade de Farmácia, Universidade do Porto, Rua Jorge Viterbo Ferreira no. 228, 4050-313 Porto, Portugal; ²Faculdade de Ciências da Nutrição e Alimentação, Universidade do Porto, Rua Dr Roberto Frias, 4200-465 Porto, Portugal; ³Laboratório Nacional de Referência de Infecções Gastrointestinais, Departamento de Doenças Infecciosas, Instituto Nacional de Saúde Dr Ricardo Jorge, Avenida Padre Cruz, 1649-016 Lisboa, Portugal

*Corresponding author. Tel: +351-22-5074320; Fax: +351-22-5074329; E-mail: patriciaantunes@fcna.up.pt

†These authors contributed equally to the experimental work.

Sir,
Fluoroquinolones are critical antibiotics for treating severe *Salmonella* infections, and the widespread of resistant isolates included in diverse epidemiological scenarios and carrying plasmid-mediated quinolone resistance (PMQR) is a global threat.^{1,2} Among PMQR mechanisms, those encoded by *oqxAB* and *aac(6')-Ib-cr* genes are of special concern as they also confer reduced susceptibility to other antibiotics (*oqxAB*: chloramphenicol, trimethoprim, olaquinox; *aac(6')-Ib-cr*: aminoglycosides) and biocides [*oqxAB*: quaternary ammonium compounds (QACs)].^{2,3} Although *oqxAB* ± *aac(6')-Ib-cr* are prevalent and widespread in Asia, where olaquinox is still widely used in animal production, they remain scarce in Europe.^{1,2,4–6} Here we describe the molecular characterization of clinical ciprofloxacin-resistant *Salmonella enterica* Typhimurium with concomitant presence of *oqxAB* and *aac(6')-Ib-cr* recovered for the first time in Europe.

Two ciprofloxacin-resistant (MIC 2 mg/L) *S. enterica* Typhimurium isolates carrying *oqxAB* and *aac(6')-Ib-cr* genes were isolated from faeces of two young children (1 and 2 years old) admitted to a Portuguese hospital in September 2012. Further characterization included the study of susceptibility to 11 other antimicrobial agents and β-lactamase and carbapenemase production by disc diffusion, Etest and/or microdilution methods (EUCAST/CLSI guidelines) and the Blue-Carba test. Screening of genes encoding resistance to antibiotics (including quinolone resistance-determining region mutations and PMQR) or tolerance to biocides/metals found in the animal setting (e.g. feed/disinfectants) was performed by PCR and/or sequencing.^{7–9}

Determination of transferability of PMQR genes (conjugation at 25/30/37°C and transformation) and characterization of integrons and plasmid backbones were performed by PCR, plasmid-based replicon typing, plasmid MLST (pMLST; <http://pubmlst.org/plasmid/>), hybridization (I-Ceul/S1-PFGE nuclease) and/or sequencing.^{1,7,8} The *oqxAB-aac(6')-Ib-cr* genetic environment was characterized by PCR mapping based on known sequences.¹ Clonal analysis was assessed by XbaI-PFGE⁸ and MLST (<http://mlst.warwick.ac.uk/mlst/dbs/Senterica>).


The two *Salmonella* Typhimurium isolates were assigned to the widespread ST34 (<http://mlst.warwick.ac.uk/mlst/dbs/Senterica>) and presented indistinguishable PFGE patterns (Figure S1, available as Supplementary data at JAC Online). Clonal expansion of *Salmonella* Typhimurium, including the ST34 clone, with PFGE profiles closely related to those of our isolates, has been associated with the spread of *oqxAB-aac(6')-Ib-cr* genes in Asian human clinical and food-producing animal isolates.^{1,6} However, the isolates reported here could not be related to any outbreak of foodborne infection or to particular food products/animal contact and foreign travel. Additionally, a single *gyrA* mutation (D87N) found in our *Salmonella* Typhimurium strain was identical to that detected in most *oqxAB*-positive *Salmonella* Typhimurium isolates from Asia.^{1,6} The strain was co-resistant to amoxicillin, chloramphenicol, gentamicin, kanamycin, nalidixic acid, streptomycin, sulfamethoxazole, tetracycline and trimethoprim (Table 1), and carried diverse metal tolerance genes (*merA*, mercury; *silA*, silver/copper; *pcoD*, copper; *terF*, tellurite). Transferability of *oqxAB-aac(6')-Ib-cr* was achieved by transformation, with an 8-fold increase in ciprofloxacin MIC and acquisition of resistance to other antibiotics (amoxicillin, chloramphenicol, gentamicin, kanamycin, sulfamethoxazole and trimethoprim) (Table 1). A 2-fold increase in benzalkonium chloride MIC was observed, highlighting the role of the *OqxAB* efflux pump in the decreasing susceptibility to diverse compounds with antimicrobial activity, including QACs.³ The concomitant presence of genes encoding resistance to antibiotics (e.g. quinolones, quinoxalines, florfenicol) and biocides/metals (e.g. QACs, copper) widely used in farm animals could contribute to the successful spread of *Salmonella* carrying *oqxAB-aac(6')-Ib-cr* genes by different events of co-selection.

The *oqxAB* genes were linked to an incomplete class 1 integron containing the *aac(6')-Ib-cr-bla_{OXA-1}-catB3-arr3-qacEΔ1-sul1* gene cassette array and flanked by two IS26 (Figure 1), widely spread mobilizing elements responsible for the dissemination of *oqxAB*⁹ and other clinically relevant antibiotic resistance genes.² The *oqxAB* genetic environment was identical to that found in Asian *Salmonella* Typhimurium isolates from food-producing animals.¹ In other bacteria (including different *Salmonella* serotypes) similar genetic platforms were observed, although differing in the linkage of *oqxAB* with *bla_{CTX-M}* alleles instead of the gene cassette array.^{2,4,5} This IS26-*oqxAB*-IS26-*aac(6')-Ib-cr-bla_{OXA-1}-catB3-arr3-qacEΔ1-sul1*-IS26 was located on a 180 kb IncHI2 plasmid carrying a *rep* gene 100% identical to Asian *Salmonella* Typhimurium pHXY0908 bearing *oqxAB-aac(6')-Ib-cr* genes (GenBank accession number KM877269), but with only 95% identity to that of the prototype R478 (GenBank accession number BX664015) and prototype pAPEC-01-R (GenBank accession number DQ517526).

Research letter

Table 1. Antimicrobial susceptibility phenotypes and genotypes of *Salmonella* Typhimurium isolates (234 and 248), the selected transformant (248_T) and the recipient (*Escherichia coli* DH5α)

Isolate	MIC (mg/L)			Resistance phenotype ^c /genotype to other antimicrobial agents	Metal tolerance genes
	CIP ^a	NAL ^b	BZK		
234	2	>256	32	AMX, CHL, GEN, KAN, STR, SUL, TET, TMP/ <i>bla</i> _{OXA-1} - <i>bla</i> _{TEM} , <i>catB3-cmlA-floR</i> , <i>aac(3)-IV</i> , <i>aphA1</i> , <i>aadA-strA-strB</i> , <i>sul1-sul2-sul3</i> , <i>tet(B)</i> , <i>dfrA12</i>	<i>merA</i> , <i>silA</i> , <i>pcoD</i> , <i>terF</i>
248	2	>256	32	AMX, CHL, GEN, KAN, STR, SUL, TET, TMP/ <i>bla</i> _{OXA-1} - <i>bla</i> _{TEM} , <i>catB3-cmlA-floR</i> , <i>aac(3)-IV</i> , <i>aphA1</i> , <i>aadA-strA-strB</i> , <i>sul1-sul2-sul3</i> , <i>tet(B)</i> , <i>dfrA12</i>	<i>merA</i> , <i>silA</i> , <i>pcoD</i> , <i>terF</i>
248_T	0.25	128	16	AMX, CHL, GEN, KAN, SUL, TMP/ <i>bla</i> _{OXA-1} , <i>catB3-cmlA-floR</i> , <i>aac(3)-IV</i> , <i>aphA1</i> , <i>aadA</i> , <i>sul1-sul2-sul3</i> , <i>dfrA12</i>	<i>terF</i>
<i>E. coli</i> DH5α	0.03	64	8		

^aPMQR genes screened: *qnrA*, *qnrB*, *qnrC*, *qnrD*, *qnrS*, *qepA*, *aac(6')-Ib-cr* and *oqxAB*.^b*Salmonella* isolates presented a single *gyrA* mutation D87N.^cSusceptibility to amoxicillin (AMX), benzalkonium chloride (BZK), ciprofloxacin (CIP), chloramphenicol (CHL), gentamicin (GEN), imipenem, kanamycin (KAN), nalidixic acid (NAL), streptomycin (STR), sulfamethoxazole (SUL), tetracycline (TET) and trimethoprim (TMP) was determined.**Figure 1.** Genetic environment (12 110 bp) of *oqxAB-aac(6')-Ib-cr* genes in an IncHI2 plasmid from an ST34 *Salmonella* Typhimurium strain. Horizontal arrows represent the positions and transcriptional directions of the ORFs. The IS26 elements are shown as light grey boxes and dotted outlines indicate incomplete sequences.

plasmids. Also, among the metal tolerance genes found, our IncHI2 plasmid and pHX0908 carried only *terF*, contrasting with the prototypes R478 and pAPEC-01-R with multiple metal tolerance genes (e.g. *merA*, *arsB*, *silA* or *pcoD*). The spread of *oqxAB*±*aac(6')-Ib-cr*-producing bacteria of both human and animal origin in Asia has been linked mostly to IncHI2 MDR plasmids^{1,4-6} belonging to ST1/ST2 by pMLST,⁴ the two major plasmid variants associated with the dissemination of diverse clinically relevant genes (e.g. *bla*_{ESBL}).¹⁰ Our MDR IncHI2 plasmid was non-typeable by pMLST (lacking allele *smr0199/smr0018*=allele number 3), suggesting the occurrence of multiple recombination events and of a new variant, considering those circulating in Europe (<http://pubmlst.org/plasmid/>).¹⁰

To the best of our knowledge, we describe for the first time in Europe an MDR *Salmonella* strain harbouring both *oqxAB* and *aac(6')-Ib-cr* PMQR genes within an IncHI2 plasmid. The spread of these bacteria and/or MDR plasmids, as is occurring in Asia in human/animal settings, is of concern since they may contribute to amplification of *oqxAB-aac(6')-Ib-cr* among Enterobacteriaceae in Europe, probably under different selective pressures (e.g. antibiotics/biocides/metals). Continuous surveillance to contain further transmission of these PMQR genes for new hosts or different settings, facilitated by international human and animal/food travel, is urgently required to preserve critical compounds with antimicrobial activity, including fluoroquinolones.

Acknowledgements

We wish to thank to Dr Lina Cavaco (Division of Microbiology and Risk Assessment, DTU Food, Technical University of Denmark, National Food Institute, Denmark) who kindly provided PMQR control strains.

Funding

This work was supported by a research grant from the Fundação para a Ciência e a Tecnologia (grant number UID/Multi/04378/2013). J. Campos and J. Mourão were supported by PhD fellowships from Fundação para a Ciência e a Tecnologia (grant numbers SFRH/BD/93091/2013 and SFRH/BD/77518/2011, respectively).

Transparency declarations

None to declare.

Supplementary data

Figure S1 is available as Supplementary data at JAC Online (<http://jac.oxfordjournals.org/>).

Research letter

References

- 1 Li L, Liao X, Yang Y et al. Spread of *oqxAB* in *Salmonella enterica* serotype Typhimurium predominantly by IncHI2 plasmids. *J Antimicrob Chemother* 2013; **68**: 2263–8.
- 2 Jacoby GA, Strahilevitz J, Hooper DC. Plasmid-mediated quinolone resistance. *Microbiol Spectr* 2014; **2**: doi:10.1128/microbiolspec.PLAS-0006-2013.
- 3 Hansen LH, Jensen LB, Sørensen HI et al. Substrate specificity of the OqxAB multidrug resistance pump in *Escherichia coli* and selected enteric bacteria. *J Antimicrob Chemother* 2007; **60**: 145–7.
- 4 Li L, Liao X, Liu Z et al. Co-spread of *oqxAB* and *bla_{CTX-M-9G}* in non-Typhi *Salmonella enterica* isolates mediated by ST2-IncHI2 plasmids. *Int J Antimicrob Agents* 2014; **44**: 263–8.
- 5 Liu B-T, Yang Q-E, Li L et al. Dissemination and characterization of plasmids carrying *oqxAB-bla_{CTX-M}* genes in *Escherichia coli* isolates from food-producing animals. *PLoS One* 2013; **8**: e73947.
- 6 Wong MHY, Yan M, Chan EWC et al. Emergence of clinical *Salmonella enterica* serovar Typhimurium isolates with concurrent resistance to ciprofloxacin, ceftriaxone, and azithromycin. *Antimicrob Agents Chemother* 2014; **58**: 3752–6.
- 7 Antunes P, Mourão J, Machado J et al. First description of *qnrS1*-IncN plasmid in a ST11 *Salmonella* Enteritidis clinical isolate from Portugal. *Diagn Microbiol Infect Dis* 2011; **69**: 463–5.
- 8 Mourão J, Novais C, Machado J et al. Metal tolerance in emerging clinically relevant multidrug-resistant *Salmonella enterica* serotype 4,[5],12:i:- clones circulating in Europe. *Int J Antimicrob Agents* 2015; **45**: 610–6.
- 9 Kim HB, Wang M, Park CH et al. *oqxAB* encoding a multidrug efflux pump in human clinical isolates of Enterobacteriaceae. *Antimicrob Agents Chemother* 2009; **53**: 3582–4.
- 10 García-Fernández A, Carattoli A. Plasmid double locus sequence typing for IncHI2 plasmids, a subtyping scheme for the characterization of IncHI2 plasmids carrying extended-spectrum β -lactamase and quinolone resistance genes. *J Antimicrob Chemother* 2010; **65**: 1155–61.

Figure S1 is available as Supplementary data at JAC online




Figure S1. PFGE patterns of clinical ciprofloxacin-resistant *S. enterica* isolates carrying *oqxAB* and *aac-6'-Ib-cr* genes recovered from two young children admitted in a Portuguese hospital (2012). The *Salmonella* serotype and multilocus sequence type (MLST) are also indicated.



Chapter | 4

Conclusions

“Anyone who has never made a mistake has never tried anything new.”

Albert Einstein

Overall, the results of this thesis conducted to the following **general CONCLUSION**:

The acquisition of metal tolerance genes, particularly to copper and silver, confers an advantage to particular clinically relevant multidrug-resistant *Salmonella* serotypes/clones by facilitating their better survival and persistence in environments contaminated with those metals across the food chain.

Further conclusions are summarized as follows:

1) Pig production setting is an important source of clinically relevant *Salmonella* serotypes/clones

- Clinically relevant MDR *Salmonella* serotypes/clones are particularly associated with pig setting in Portugal (2002-2014), namely to piggeries environment and pork products. These MDR *Salmonella* serotypes/clones were mainly identified as *S. Typhimurium* (European clone), *S. 4,[5],12:i:-* (European, Spanish and Southern European clones) and *S. Rissen/ST469*. They are a public health issue emergent in diverse European regions and now well characterized (e.g. clonality, MDR profiles and associated genetic platforms, sources) in our country.

- The presence of typical molecular patterns in clones of the worldwide epidemic *S. 4,[5],12:i:-* as well as *S. Typhimurium* European clone and *S. Rissen* highlights the ability of those serotypes/clones to acquire and maintain different adaptive features (e.g. MDR) with potential impact in their emergence. This was demonstrated for the most prevalent *S. 4,[5],12:i:-* European (ASSuT phenotype, RR1-RR2 resistance regions, ST34) and Spanish clones (ACGSSuTTm phenotype, *sul3*-type III integron within IncA/C plasmids, ST34) but also for the less frequent Southern European clone (CSSuTTm phenotype, *sul3*-type I integron within IncR plasmids, ST19), both in isolates from human clinical samples, food, environment and piggeries. Also, *S. Rissen* (ASSuTTm phenotype, ST469) and *S. Typhimurium* European clone (ASSuT phenotype, ST34) carried particular features.

2) Epidemiology of metal tolerance in *Salmonella*

Acquired metal tolerance genes were dispersed in diverse *Salmonella* genetic backgrounds, especially in clinically relevant MDR pig-associated serotypes or clones

- Genes coding for copper (*pcoA-pcoD/silA-silE*), silver (*silA-silE*) and mercury (*merA*) tolerance were detected in isolates from diverse sources, clones and serotypes and were more frequently observed than those coding for arsenic (*arsB*) or tellurite (*terF*) tolerance, only found in few isolates/serotypes. The dispersion of those genes in *Salmonella* from different serotypes and clones suggests a genetic exchange between members sharing the same ecological communities and hosts potentially subjected to similar metal environmental stresses.

- *Salmonella* of emerging pig-associated serotypes/clones are more enriched in genes coding for copper and silver (*silA-silE±pcoA-pcoD*) than *Salmonella* serotypes/clones associated with other food-animal settings, which is possibly explained by variable selection contexts of animal production practices. This is supported by the allowed use of higher copper amounts in feed for pigs comparing with other animals, which might have contributed for the selection and persistence of emerging *Salmonella sil±pco*⁺, particularly the MDR serotypes/clones *S.* 4,[5],12:i:- European clone/ST34, *S.* Typhimurium European clone/ST34 and *S.* Rissen/ST469. In contrast, *sil* and *pco* genes were absent in several poultry/eggs-associated serotypes (e.g. *S.* Enteritidis, *S.* Infantis, *S.* Heidelberg), a setting where only low Cu concentrations are allowed in Europe. Nevertheless, an often occurrence of *sil±pco* in available genomes of emerging poultry-associated serotypes (e.g. Heidelberg, Kentucky), suggests that different production practices (e.g. in USA copper supplements are not restricted by animal type) might contribute for the selection and emergence of different *Salmonella* serotypes or strains.

- *Salmonella* isolates carrying *sil±pco* genes coding for copper/silver tolerance were more resistant to antibiotics than those without these genes clusters. This data highlights the importance of carrying different genetic features involved in an improved survival capacity to environments subject to different stressors (e.g. metals or antibiotics), which might account for the expansion of MDR *Salmonella* serotypes/clones across the food-animal setting.

- The combination of *sil±pco* genes can be used as additional molecular biomarkers of current emerging clones in Europe. This is supported by the occurrence of the *sil±pco* genes in almost all *S.* Rissen, *S.* 4,[5],12:i:- and *S.* Typhimurium European clones and the *sil* gene alone in *S.* 4,[5],12:i:- Spanish and Southern European clones.

3) Metal tolerance and antibiotic resistance: characterization of genetic elements and horizontal gene transfer

Variable genetic platforms contribute for the co-dispersion of metal tolerance and antibiotic resistance genes

- Metal tolerance and antibiotic resistance genes are often co-located in the same genetic platforms, sharing either the same plasmid or chromosomal regions in diverse *Salmonella* serotypes. Diverse metal tolerance genes were located in the chromosome (*sil±pco*, *merA*, *terF*) or in large plasmids (*sil±pco*, *merA*, *terF*, *arsB*) along with those coding for resistance to other antimicrobial agents as biocides (e.g. *oqxAB*-quaternary ammonium compounds) or antibiotics (e.g. ampicillin-*bla*_{TEM}, tetracycline-*tet* or sulphonamides-*sul* genes) including some relevant in the treatment of *Salmonella* infections (e.g. *bla*_{CTX-M-9}-extended-spectrum cephalosporins, or *aac(6')-Ib-cr/oqxAB*-fluoroquinolones). Also, the combination of particular metal (*sil±pco* and/or *merA*) and antibiotic resistance (e.g. *tet* genes, typical-*sul1* and/or atypical *sul3*-type class 1 integrons) genes found within these genetic platforms might also reflect their abundance in the animal setting and thus the probability to recombine and accumulate in the same genetic platforms.

- Large plasmids carrying metal tolerance genes belong to families widely identified in *Enterobacteriaceae* and often involved in antibiotic resistance spread. This is supported by the detection of metal tolerance genes (*sil±pco*, *merA*, *terF*, *arsB*) in major plasmid families such as IncA/C, IncR, IncFIIA, IncHI1, IncHI2, IncN, IncP and IncI1, some of them transferable. Although some of those plasmids (e.g. IncHI2) have been frequently associated with diverse metal tolerance genes, we unveiled the relevance of IncFIIA and IncN plasmids in the dispersion of such genes. These data also alerts for the importance of single horizontal transfer events in the co-dispersion of both metal tolerance and clinically relevant antibiotic resistance genes [e.g. *aac(6')-Ib-cr/oqxAB*, *bla*_{CTX-M-9}] among bacteria from different species sharing the same ecosystems and subjected to diverse selective pressures.

- Chromosomal location of the *sil±pco* genes coding for copper/silver tolerance and/or *merA* were mainly observed in the MDR European clone of *S. Typhimurium* and *S. 4,[5],12:i:-* and in the *S. Rissen*. These data along with the inability of transfer chromosomal genetic platforms in conjugation assays highlight the importance of clonal spread for the maintenance of *sil±pco* in *Salmonella* recovered from the food-animal setting. However, the need of critical and particular conditions for the mobilization of

chromosomal genetic platforms carrying *sil±pco* in laboratory assays could not be discarded.

- The *sil* and *pco* gene clusters are adjacent in the same genetic region and are inserted in a Tn7-like element, the Tn6230, both in plasmids and the chromosome, particularly in recently emergent *Salmonella* serotypes/clones. This was supported by our data and/or by a detailed *in silico* genomic analysis of available *Salmonella* genomes from different sources and serotypes. In some cases, the absence of transposition proteins of Tn6230-like elements could be related to recombination events potentially associated with the fixation of these elements in diverse *Salmonella* serotypes. Further studies are needed to clarify the contribution of Tn6230-like elements in the emergence of particular *Salmonella* serotypes/clones as *S. Typhimurium* and *S. 4,[5],12:i:-*.

4) Copper and silver tolerance phenotypes in *Salmonella*

Higher tolerance to CuSO₄ in anaerobic atmosphere and to AgNO₃ after prior exposure to silver was observed among the *sil+pco*⁺ or *sil*⁺ isolates

- Wild type *Salmonella* isolates and transconjugants carrying the *sil* efflux system, independently of the chromosomal or plasmid location, were associated with higher tolerance to CuSO₄ in anaerobiosis and to AgNO₃ after prior exposure to silver in aerobiosis. These data contributed to demonstrate the role of the *sil* gene cluster in *Salmonella* adaption to metal contaminated environments, namely to overcome the more toxic Cu⁺ occurring in reduced settings potentially present in food-animal productions (e.g. animal gut of livestock carriers, manure, waste lagoons or forage/feed, sewage and sludge).

- Further studies are needed to understand the "biocidome" associated with non-typhoidal *Salmonella*: i) to clarify the potential role of the *pco* gene cluster in other important adaptive features (e.g. virulence); ii) to unveil other genetic determinants associated with copper tolerance, which could explain the presence of high levels of aerobic CuSO₄ tolerance in *Salmonella* isolates, carrying or not *pco* genes, or the occurrence of high anaerobic CuSO₄ tolerance in isolates lacking the known *sil±pco* gene clusters.

Modified methodological approaches in Cu/Ag tolerance assays are critical to identify isolates carrying *sil±pco* gene clusters

- The use of anaerobiosis to determine *Salmonella* CuSO₄ tolerance and the prior contact with subinhibitory concentrations of AgNO₃ is mandatory for the detection of the higher MICs to these compounds determined by *sil±pco* acquisition, independently of the *Salmonella* serotype/clone. The inclusion of anaerobiosis, here firstly applied to the standard method using aerobiosis, was considered because *silE* binds to the more toxic form Cu⁺ in anaerobiosis. In the case of AgNO₃, despite of *sil* gene cluster had been reported as being inducible in the presence of silver, few studies used this approach (applied a prior contact with silver before MIC determination) or the methodological details were poorly described, impairing its reproducibility. Thus, well-described protocols allowing reproducible results are published and available in this thesis (Chapter 5 - Appendix) for general use.

- Here we have proposed tolerance cut-offs to CuSO₄/AgNO₃ that can be used in further studies in order to distinguish *Salmonella* isolates with and without *sil* genes, which will have a significant impact in data comparison from different studies. This is critical, as currently the use of different methodologies is associated with an unclear link between phenotypic and genotypic data related with *sil* acquisition, which hinders the understanding of copper and silver tolerance in *Salmonella*. The application of these cut-offs is only suitable when the modified methodological approaches proposed are applied.

CLOSING REMARKS

Different gaps in the field of metal and biocide tolerance knowledge have been pointed out by different entities such as the European Scientific Committee on Emerging and Newly Identified Health Risks (SCENIHR). The lack of information on this topic has hindered the understanding of the impact of the use of biocides and metals in public health. Among other issues, it is referred the lack of monitoring programs, which limits the gathering of robust epidemiological data and knowledge about the selection and spread of antibiotic resistance genes associated with the use of metals and biocides. It is also denoted the absence of standardized methods for the evaluation of bacterial susceptibility to these compounds.

In *Salmonella*, a foodborne zoonotic pathogen of clinical importance with ability to genetically interact with different microorganisms in diverse biological communities, the results of this study contributed to:

- a) Know the epidemiology of tolerance to copper, silver, mercury, arsenic and tellurium using, for the first time, a large collection of *Salmonella* isolates (n=406) representative of different epidemiological and genetic backgrounds (niches/sources, time frames, serotypes and clones). It was showed that the pig production setting seems to be a relevant reservoir of successful and worldwide emerging MDR *Salmonella* serotypes/clones enriched in different adaptive features (besides antibiotic resistance or virulence genes), including tolerance to metals currently used in food-animal setting.
- b) Understand the potential role of metals, particularly copper and silver, widely present in different anthropogenic contexts, in the selection and maintenance of antibiotic-resistant *Salmonella*, through the characterization of the genetic elements harbouring metal tolerance and antibiotic resistance genes.
- c) Develop more efficient methodological approaches for detecting copper and silver tolerance and to establish tolerance cut-offs to CuSO₄/AgNO₃, which will allow the comparison of data in further studies.

The new information provided by this study might help to support the implementation of more efficient intervening measures related to the use and/or accumulation of metals in diverse environments, in order to prevent a wider expansion of foodborne zoonotic pathogens such as MDR *Salmonella* serotypes/clones or the emergence of new ones.



Appendix

“If you want to live a happy life, tie it to a goal, not to people or objects.”

Albert Einstein

I. Review paper – “Salmonellosis: the role of poultry meat”

Salmonellosis: the role of poultry meat

CMI Themed Issue “Poultry and human infections”

Patrícia Antunes^{1,2}, Joana Mourão², Joana Campos² and Luísa Peixe^{2*}

¹ Faculdade de Ciências da Nutrição e Alimentação, Universidade do Porto, Rua Dr. Roberto Frias, 4200 Porto, Portugal; ² UCIBIO/REQUIMTE, Laboratório de Microbiologia, Faculdade de Farmácia, Universidade do Porto, Rua Aníbal Cunha, no. 164, 4050-047 Porto, Portugal.

Clinical Microbiology and Infection 2016 [Epub ahead of print]

Elsevier has authorized the reproduction of the final published PDF version of this paper in this thesis through the License Agreement Number 3800171308082 on February 01, 2016.

ARTICLE IN PRESS

REVIEW

Salmonellosis: the role of poultry meat

P. Antunes^{1,2}, J. Mourão², J. Campos² and L. Peixe²

1) Faculdade de Ciências da Nutrição e Alimentação, Universidade do Porto and 2) UCIBIO/REQUIMTE. Laboratório de Microbiologia, Faculdade de Farmácia, Universidade do Porto, Porto, Portugal

Abstract

Salmonellosis remains one of the most frequent food-borne zoonoses, constituting a worldwide major public health concern. Currently, at a global level, the main sources of infection for humans include meat products, including the consumption of contaminated poultry meat, in spite of the success of *Salmonella* control measures implemented in food-animal production of industrialized countries. In recent years, a shift in *Salmonella* serotypes related to poultry and poultry production has been reported in diverse geographical regions, being particularly associated with the spread of certain well-adapted clones. Moreover, antimicrobial resistance in non-typhoidal *Salmonella* is considered one of the major public health threats related with food-animal production, including the poultry production chain and poultry meat, which is an additional concern in the management of salmonellosis. The circulation of the same multidrug-resistant *Salmonella* clones and/or identical mobile genetic elements encoding antibiotic resistance genes from poultry to humans highlights this scenario. The purpose of this review was to provide an overview of the role of poultry meat on salmonellosis at a global scale and the main problems that could hinder the success of *Salmonella* control measures at animal production level. With the increasing globalization of foodstuffs like poultry meat, new problems and challenges might arise regarding salmonellosis control, making new integrated intervention strategies necessary along the food chain.

Clinical Microbiology and Infection © 2015 European Society of Clinical Microbiology and Infectious Diseases. Published by Elsevier Ltd. All rights reserved.

Keywords: AmpC, extended-spectrum β -lactamases, multidrug-resistance, plasmid-mediated quinolone resistance, *Salmonella*, Enteritidis, Infantis, Kentucky, Heidelberg, Typhimurium, Virchow

Article published online: XXX

Corresponding author: L. Peixe, Laboratório de Microbiologia, Faculdade de Farmácia, Universidade do Porto, Rua Jorge Viterbo Ferreira n° 228, 4050-313 Porto, Portugal
E-mail: lpeixe@ff.up.pt

Introduction

Salmonella infections are a worldwide major public health concern; Salmonellosis is caused by non-typhoidal *Salmonella enterica* serotypes (serotypes other than *S. Typhi* and *S. Paratyphi*) and is typically characterized by a self-limiting gastro-enteritis syndrome (manifested as diarrhoea, fever and abdominal pain), with an incubation period between 4 and 72 h and mortality being rare [1–3]. In healthy humans, the

infectious dose is generally 10^6 to 10^8 , but lower bacterial counts can cause disease in certain conditions, as well as in infants and the elderly [2]. Although uncommon, life-threatening invasive infections with bacteraemia (5%–10% of infected persons) and/or other extra-intestinal infections may occur, affecting especially the risk groups (infants, young children, older people and immunocompromised patients) [1–3]. In severe cases, effective antimicrobial agents are essential, so the emergence of *Salmonella* that are resistant to critical antibiotics is of concern [1,2].

In industrialized countries, the main reservoir of non-typhoidal *Salmonella* is the intestinal tract of food-producing animals, which readily leads to contamination of diverse foodstuffs [3–5]. Therefore, despite other sources (e.g. contact with animals/reptiles, environment or person-to-person), food-borne salmonellosis is the most relevant source with a high

global impact in human health. It was estimated that non-typhoidal *Salmonella* causes around 93.8 million illnesses and 155 000 deaths each year worldwide [6]. In the USA, more than 1 million annual cases of food-borne salmonellosis were estimated by the CDC; they were associated with the largest number of hospitalizations and deaths compared with other food-borne microbial agents [7]. In Europe, salmonellosis has been the second most common zoonosis (82 694 confirmed cases and 20.4 cases per 100 000 population in 2013) and the most frequent cause of food-borne outbreaks, in spite the reported decreasing trend that has resulted from *Salmonella* control programmes [8].

Although different serotypes have been associated with salmonellosis, a limited number are responsible for most human infections; *S. enterica* Enteritidis being the most frequent one in the EU (39.5% in 2013) and USA (14.5% in 2012) followed by *S. enterica* Typhimurium (including its monophasic variant) (28.8% in EU, 2013; 11.6% in USA) [8,9]. *Salmonella* Enteritidis is commonly associated with poultry and products thereof, whereas *S. Typhimurium* has a wider species range, including pigs and cattle as well as poultry [10]. Therefore, foods of animal origin, in particular contaminated poultry products (eggs and poultry meat) have been considered the main vehicles of *Salmonella* infection and clearly associated with the worldwide epidemic of *S. Enteritidis* [4,5,11,12]. Moreover, diverse epidemiological studies have supported the great contribution of poultry foodstuffs to the salmonellosis burden [4,5,12,13].

In recent years, with the implementation of *Salmonella* control programmes mainly in poultry production (e.g. in the EU and USA), changing trends in food-borne salmonellosis and associated serotypes were observed, with the expansion of previously less common serotypes, frequently resistant to antibiotics. With the increasing globalization of foodstuffs like poultry meat, which is one of the most consumed and increasingly globally traded meat products, new problems might arise regarding salmonellosis control. An overview of the role of poultry meat in salmonellosis at a global scale is provided in this review.

Non-typhoidal *Salmonella* and poultry meat

Poultry populations, in particular chicken and turkey, are frequently colonized with *Salmonella* without detectable symptoms (sub-clinical infections/healthy carriers) by horizontal and vertical transmission at primary production level [4,5]. The presence of *Salmonella* in healthy poultry animals is suggested as the main risk factor, by allowing bacteria to easily transmit in table eggs and poultry meat to humans [10]. In fact, in Europe it is assumed that the observed reduction in salmonellosis cases

(32% between 2008 and 2012) is mainly due to successful *Salmonella* control measures (involving surveillance, biosecurity and vaccination) implemented in poultry/egg production and focused on particular serotypes (e.g. *S. Enteritidis* and *S. Typhimurium*) that are considered of public health significance [4,8,10,12]. These measures led to the achievement of reduction targets for poultry populations in most EU countries and lower non-compliance regarding *Salmonella* in poultry products [4,8]. Moreover, decreasing contamination rates in raw poultry products are in agreement with those recently observed in industrialized countries from other geographical regions with pathogen reduction programmes, such as the USA [12,14,15]. It should be noted that by the 2000s a high incidence of *Salmonella* in poultry products was reported in the EU, with rates >50% for several countries [16]. In the 2013 zoonosis EFSA/ECDC report involving data from European countries, as in previous years, *Salmonella* was most frequently reported, although at low levels, in fresh turkey (5.4%) and fresh broiler meat (3.5%), in comparison with eggs (0.1%) or fresh pig meat (0.7%) [8]. Despite the highest incidence being detected in poultry meat, eggs still remain the most important source of food-borne *Salmonella* outbreaks [8]. In fact, using quantitative source attribution models the higher number of human salmonellosis cases in Europe was attributable to eggs (65% in 2011 and 17% in 2012) and pigs (28% in 2011 and 56.8% in 2012) compared with broilers (2.4% in 2011 and 10.6% in 2012) and turkey meat (2.6% in 2011 and 4.5% in 2012) [17,18]. However, diverse surveys targeted to detect *Salmonella* in poultry products in developing countries, some with expansion of the poultry industry, still detected high percentages of positive samples, ranging from ~13% to 39% in South America [19,20], ~35% in Africa [21,22] and ~35% to 50% in Asia [23–25]. Those differences possibly reflect diverse poultry production husbandry practices and absence of control measures along the food chain, highlighting the importance for *Salmonella* spread of the extensive international trade in animals and their products [4].

Worldwide data about *Salmonella* serotype prevalence in humans and in the diverse range of foodstuffs have contributed to establish an epidemiological link between salmonellosis and poultry products, with diverse serotypes overlapping between humans and poultry meat (chicken and turkey) (Fig. 1). In the EU, recent changes in the frequency of *Salmonella* serotypes causing human infections were reported, which in some cases were in line with those occurring in poultry (Fig. 1). Nevertheless, interpretation of these data should be cautious, owing to limitations in the number of poultry isolates serotyped each year. Of particular relevance is the decrease in *S. Enteritidis* human cases (19% reduction between 2011 and 2013 in the EU), a serotype typically associated with poultry meat and egg




FIG. 1. Distribution of the major serotypes of non-typhoidal *Salmonella* associated with human cases (salmonellosis) and poultry meat in EU, 2011 to 2013. Data were obtained from EFSA reports: humans (2011–2013) and turkey/broiler meat (2013) [8]; turkey/broiler meat (2011–2012) (<http://www.efsa.europa.eu/en/efsajournal/pub/3129> and <http://www.efsa.europa.eu/en/efsajournal/pub/3547>) (last accessed 1 September 2015). The percentages were calculated based on the total number of serotyped isolates per type of meat or human salmonellosis cases.

consumption, in line with a decreasing trend in poultry and poultry products pointed by different studies [8,12,26–29]. Increasing occurrences of other serotypes implicated in human infections (e.g. *S. Infantis*, *S. Stanley*, *S. Kentucky*) related with poultry meat (chicken and turkey) have been reported in the EU (Fig. 1). However, one major difference in the serotype pattern between humans and poultry is related to *S. Typhimurium* and its monophasic variant, both frequently associated with human salmonellosis, but less common in poultry meat, pigs and pig meat being the main source [8]. In USA and Canada, other serotypes, like *S. Heidelberg* and *S. Kentucky*, have emerged as predominant serotypes in poultry and have also been implicated in human salmonellosis, beyond *S. Enteritidis* [9,12,14,15,28,30–32].

The shift in *Salmonella* serotypes related to poultry and poultry production has been associated with the spread of certain clones. For instance, the *S. Infantis* increase (26.5% in human cases between 2011 and 2013 accounting for the fourth most common serotype and the most reported in broilers and broiler meat) [8], has been associated with the spread of several clones of broiler origin in diverse European countries, including the dominant Hungarian clone [33–35]. Also, *S. Stanley* showed an increase since 2011 with a peak in 2012, being the sixth most common human serotype and one of the three most common

in turkey meat, together with *S. Derby* and *S. Kentucky* in 2013 [8]. A large *S. Stanley* outbreak caused by a new clone (novel Pulsed Field Gel Electrophoresis-type), was linked with the consumption of turkey meat, and is still circulating in the European food market (at least since 2011), with a considerable risk of becoming endemic in the poultry production chain in Europe [36–38]. In the USA, *S. Heidelberg* in particular has been identified as one of the top human and poultry serotypes, with several clones implicated in diverse large multistate outbreaks resulting from the consumption of contaminated chicken or turkey products [12,31]. The spread and the global persistence of serotype *S. Kentucky* reflect other particular situations related to the increased globalization of travel and the food/animal trade in different geographical regions. This serotype has been associated with a worldwide (Europe, Africa and Asia) spread of a particular epidemic clone (*S. Kentucky* ST198-X1), recovered from several livestock reservoirs, particularly poultry farms, with chicken and turkey implicated as the potential major human infection vehicles [39–43]. These and other examples of multi-country/multi-state outbreaks or clonal expansion of *Salmonella* infections linked to poultry meat (Table 1) serve as a reminder of the importance of acting upon any *Salmonella* contamination in the food chain and monitoring to detect the emergence of any serotype or new clone. This

TABLE 1. *Salmonella* outbreaks and emerging clones linked with poultry meat products (2002–ongoing)

Serotype (outbreak/clone designation) ^a	Source	Year(s) ^b	Geographical region	No. of cases ^c	Reference
Enteritidis (outbreak)	Chicken	2015–	USA	3	CDC, 2015 ^d
Enteritidis (outbreak)	Chicken	2015–	USA	9	CDC, 2015 ^d
Hadar (outbreak)	Turkey	2010–2011	USA	12	CDC, 2015 ^d
Heidelberg (outbreak)	Chicken	2013–2014	USA, Puerto Rico	634	CDC, 2015 ^d
Heidelberg (outbreak)	Chicken	2013	USA	9	CDC, 2015 ^d
Heidelberg (outbreak)	Chicken	2012–2013	USA	134	CDC, 2015 ^d
Heidelberg (outbreak)	Chicken	2011	USA	190	CDC, 2015 ^d
Heidelberg (outbreak)	Turkey	2011	USA	136	CDC, 2015 ^d
Infantis (Israel clone)	Broiler chickens	2007–2009	Israel	NA	[26]
Infantis (Hungarian clone)	Broilers and chicken	2004–2009	Europe	NA	[33–35]
Kentucky (ST198-X1)	Chicken and turkey	2002–2013	Europe, Africa, Asia	NA	[39,40,42]
Stanley (outbreak)	Turkey	2011–2013	Europe	710	[36,37]

^aST, sequence type.^byear- include ongoing reported outbreaks.^cEstimated number of cases only when outbreaks were reported. NA, not applicable in emerging clones.^dCDC, 2015. *Salmonella*. Reports of selected *Salmonella* Outbreak Investigations. Available at: <http://www.cdc.gov/salmonella/outbreaks.html> [accessed September 2015].

scenario also remains alert for the need for inclusion of those serotypes, or particular clones, considered to be of public health significance, in control and surveillance programmes [10,12,38].

Salmonella serotypes and clones associated with human infections and with an enhanced ability to colonize several food animals, able to persist along the food chain (e.g. primary production on-farm, slaughter operations, equipment and meat handlers, retail meat) with efficient transmission and rapid spread are of public health relevance [11,34,43–45]. Although understanding the exact mechanisms of their persistence and spread in poultry production are still largely unknown, recent studies focusing on emergent poultry-associated *Salmonella* strains unveiled specific features that could provide a significant advantage both in the environment and in the host (poultry/human) [12]. For example, in Israel, an *S. Infantis* emergent clone possessed a megaplasmid, which increased its tolerance to stress factors (e.g. mercury and oxidative stress) and its virulence/pathogenicity (e.g. enhanced biofilm formation, adhesion and invasion into avian and mammalian host cells) [46]. Also, a genomic study of several predominant *Salmonella* serotypes from Canadian broiler chickens showed the presence of multiple features related with pathogenicity (e.g. genes encoding adhesins, flagellar proteins, iron acquisition systems, type III secretion system) and stress tolerance (e.g. metal and antiseptic tolerance genes; better acid-stress response) [32]. *S. Heidelberg*, including ground turkey outbreak isolates, carried phages and plasmids with diverse virulence factors (e.g. P2-like phage-*sopE1* gene, IncX-type IV secretion system), which could play a role in their virulence (a serotype highly related to invasive infections), colonization and persistence (a poultry-associated serotype) [12,31]. In *S. Kentucky*, the acquisition of an *E. coli* ColV virulence plasmid was also associated with enhanced colonization ability in chicken, particularly in a dominant avian clonal type [12]. These features can play a role

in the successful spread of emergent and virulent serotypes/clones that could contribute in a short time to replacing other *Salmonella*. Moreover, those emergent *Salmonella* serotypes are usually enriched with antimicrobial resistance determinants conferring multidrug-resistance [12,31,32,46], which is currently one of the major public health concerns (discussed in the next section).

Antimicrobial-resistant non-typhoidal *Salmonella* and the poultry linkage

The emergence and spread of *Salmonella* isolates presenting resistance to several antibiotics is of concern because these medicines are crucial to the successful treatment of invasive infections [1,11]. Since resistance to older antibiotics (e.g. ampicillin, chloramphenicol and trimethoprim-sulfamethoxazole) has been increasing for many years, recommended treatment options for salmonellosis included fluoroquinolones (ciprofloxacin) and extended-spectrum cephalosporins [1,2]. However, resistance to those ‘critically important antibiotics for human health’ is emerging, leading to increased severity, morbidity and mortality of diseases and the need for the use of last-line antimicrobials (e.g. carbapenems) in therapy [47].

For several decades, the contribution of the food-animal as a reservoir of antimicrobial resistance with impacts on human health has been controversial, but accumulating evidence linking particularly poultry production with human disease have been reported, namely involving non-typhoidal *Salmonella* [48–50]. The first evidence is the close association between the use of antimicrobial agents and the occurrence of resistance. In fact, regular use of antibiotics (mainly for animal health and in some countries as growth promoters), a practice associated with modern intensive food-animal/poultry production, has been

considered the main driver for the development of antibiotic resistance in zoonotic bacteria, such as *Salmonella* [48,50,51]. For example, licensing of the fluoroquinolone enrofloxacin for animal use, especially in poultry, in the 1990s led to increased rates of decreased susceptibility to ciprofloxacin in *S. Typhimurium* DT104 recovered from animal/food (particularly poultry) and humans [5]. Other findings points to a link between resistance to nitrofurans in human *Salmonella* and the food chain, suggesting that its illegal use in the poultry industry might have contributed to the selection and persistence of *S. Enteritidis* in poultry, and consequently to human salmonellosis in Portugal [52]. More recently, a voluntary withdrawal of ceftiofur by the poultry producers in Canada was correlated with a decreasing occurrence of ceftiofur-resistant *S. Heidelberg* (one of the most common serotypes associated with salmonellosis in this country) from both human infections and retail poultry, with an increase of the resistance levels after reintroduction of use [30].

Further evidence for the impact of poultry production on human health problems associated with antimicrobial resistance in *Salmonella* is the correlation between different reservoirs (humans and food-animals) obtained from systematic surveillance data. In the last EU report, *Salmonella* resistant (R) or multidrug-resistant (MDR) to commonly used antimicrobials was frequently detected in humans (R = 50%; MDR = 31.8%) and animals, especially broilers (MDR = 56%) and turkeys (MDR = 73%), but also in derived meat products [43]. Also, in the USA, high levels of resistance and MDR in chicken (R = 60%; MDR = 26%) and turkey (R = 77%; MDR = 39.6%) seems to greatly contribute to levels in humans (R = 19%; MDR = 10%) [53]. Those data about MDR are extremely worrying because of the possible role of diverse antibiotics in the co-selection of *Salmonella* strains resistant to clinically relevant antibiotics, such as fluoroquinolones and extended-spectrum cephalosporins. In the EU, relatively low levels of *Salmonella* non-susceptible ('clinically' resistant and 'intermediate' resistant categories combined) to ciprofloxacin (3.8%) and 'microbiological' resistant (non-wild type by epidemiological cut-off values) to cefotaxime (1.4%) were observed in humans. Moreover, the highest levels of 'microbiological' resistance to these critically important antimicrobials were detected in broiler meat (68% ciprofloxacin and 10.1% cefotaxime) and turkey meat (73.4% ciprofloxacin and 4.7% cefotaxime), suggesting an important contribution of the poultry production chain to the human burden [43]. In addition, it was reported that the highest levels of resistance to ciprofloxacin were more common in *S. Enteritidis*, *S. Infantis* and *S. Kentucky*, three serotypes commonly associated with poultry meat [43]. Also, in the USA, in spite of a decreasing trend since 2009, cephalosporin's resistance levels are low in humans (2.5% ceftriaxone),

but still high in turkey (9%) and retail chicken (20%), especially in *S. Heidelberg* a poultry-associated serotype [53].

In middle-income countries such as in the Asiatic continent, resistance to ciprofloxacin and extended-spectrum cephalosporins is currently a growing problem, with poultry products playing a potential role in its emergence [2,3]. High levels of *Salmonella* non-susceptible to ciprofloxacin (15–48%) and cephalosporins (38% ceftriaxone) were observed in humans in Asian countries [54,55]. These data are in agreement with several studies that documented a high prevalence of resistance to fluoroquinolones (>22.5% ciprofloxacin) and cephalosporins (12.5% to >23.4% ceftriaxone and 26.6% ceftazidime) in poultry meat from South Korea and China [24,25]. In these countries the high rates of antimicrobial resistance detected reflect the levels of antimicrobial consumption in livestock, including for growth promotion, an area that remains largely unregulated [25,55,56]. Recent global estimates indicate that consumption of antibiotics in livestock (especially in poultry), which outranks the consumption by humans, will rise especially in middle-income countries. In these countries extensive farming systems will be replaced by large-scale intensive husbandry systems that routinely use antibiotics, to respond to the increasing demand for animal protein [56]. Therefore, international trade of contaminated breeding animals (poultry production depends on a pyramid-like breeding system), feed and poultry products with antibiotic-resistant *Salmonella* may contribute to the rapid worldwide spread of these bacteria, with impacts on human health.

A third indication of the significant impact on human health of antibiotic-resistant *Salmonella* contaminating these foodstuffs, is provided by diverse reports demonstrating their association with plasmid-mediated resistance determinants to cephalosporins and fluoroquinolones [49,51,57]. Moreover, several studies provide further evidence of transmission from poultry to humans of those clinically relevant MDR *Salmonella* clones [11,49,51,57]. A wide range of *Salmonella* serotypes (e.g. *Enteritidis*, *Heidelberg*, *Infantis*, *Kentucky*, *Typhimurium*, *Virchow*) frequently recovered from poultry (animal and food) have been associated with the worldwide dissemination of extended-spectrum β -lactamases, plasmid-encoded AmpC β -lactamases (AmpC), or plasmid-mediated quinolone resistance genes (Tables 2 and 3). By far the most common genes, found in poultry and/or poultry meat samples, associated with resistance to extended-spectrum cephalosporins, were those coding for extended-spectrum β -lactamases, CTX-M (e.g. CTX-M-1, -2, -9 and -15) and TEM-52 enzymes followed by AmpC-type CMY-2 (Table 2). For example, in several European countries, spread of *S. Virchow* producing CTX-M-2 or CTX-M-9 and *S. Infantis* producing TEM-52 has been found among poultry and humans [58–66]. Instead, *S. Heidelberg*, one of the

TABLE 2. *Salmonella* serotypes/clones carrying extended-spectrum and AmpC β -lactamases recovered from poultry and poultry products

Salmonella serotype	ESBL or AmpC ^a enzymes (no. of isolates)	Source	Concomitant presence in humans ^b	Country(ies)/year(s)	Non- β -lactam resistance phenotype ^c	Genetic element PL – Inc group	Reference(s)
Agona	CTX-M-1 (n=1) CTX-M-2 (n=2)	Poultry (caecum) Poultry		The Netherlands/2006	SUL-TET	PL – II	[65,66]
		Turkey carcass		Brazil/2011	NR	PL – NR	[71]
				Brazil/2008	CIP-NAL-STR-SXT-TET	PL – P	[72]
				Brazil/2008, 2010	NR	PL – NR	[71]
				Belgium/2001, 2004	(–)	PL – II	[62]
				Germany/2005	(–) or NAL	PL – II	[73]
Bareilly	CMY-2 (n=5) ACC-1 (n=6)	Poultry	Yes	The Netherlands/2001–02	(–)	PL – NR	[74]
Blockley	TEM-52 (n=6)	Poultry, poultry meat	Yes	The Netherlands/2001–02	(–) or SPT-SUL-TMP	PL – NR	[74]
Brackenridge	CTX-M-14 (n=1)	Poultry		Brazil/2011	NR	PL – NR	[71]
Braenderup	ACC-1 (n=2)	Broilers		The Netherlands/2006	(–)	PL – NT	[65]
		Broiler		The Netherlands/2001	(–)	PL – NR	[74]
				Italy/2006	(–)	PL – NR	[75]
Bredeney	SHV-12 (n=1)	Poultry		USA/2010	NR	PL – N	[76]
	CTX-M-1 (n=1)	Turkey		Canada/1999	SUL-TET	PL – A/C	[77]
	CMY-2 (n=1)	Turkey		Belgium/2001	(–)	PL – II	[62]
Derby	TEM-52 (n=1)	Poultry		Brazil/2004	NR	NR	[71]
Enteritidis	CTX-M-2 (n=1)	Broiler		Spain/2000–04	STR-SUL-SXT-NAL-TET	NR	[60]
	CTX-M-9 (n=1)	Broiler		China/2004	NAL	PL – NR	[78]
	CTX-M-14 (n=5)	Imported chicken meat		China/2012–13	CHL-CIP-ENR-FFC-NAL-OLA-SXT-(AMK-GEN-LVX-TET)	PL – HI2, F, N or B/O	[70]
	CTX-M-15 (n=34)	Chicken meat, faeces or disease	Yes	Korea/2009	STR-SUL-TET-(GEN-NEO)	PL – FII	[79]
		Chicken		South Korea/2009–10	NR or GEN-STR-SUL-TET	PL – FII	[80]
		Broiler		Portugal/2012–13	CHL-CIP-SUL-NAL-TET	NR	[81]
		Poultry		Italy/2006	TET	NR	[75]
Emek	CMY-2 (n=1)	Poultry		Brazil/2011	NR	NR	[71]
Essen	CTX-M-15 (n=1)	Diseased chicken		Korea/2009	GEN-STR-SUL-TET	PL – FII	[79]
Gaminara	CTX-M-14 (n=1)	Poultry		Brazil/2009	NR	NR	[71]
Give	CTX-M-2 (n=1)	Poultry		Brazil/2009	NR	NR	[71]
Hadar	CTX-M-14 (n=1)	Poultry		Brazil/2005	NR	NR	[71]
Havana	CTX-M-1 (n=1)	Broiler		Portugal/2012–13	SUL-TET	NR	[81]
	CMY-2 (n=2)	Broiler		Portugal/2012–13	(–)	NR	[81]
Heidelberg	CTX-M-1 (n=1)	Broiler carcass		Portugal/2012–13	SUL-TMP	NR	[81]
	CTX-M-2 (n=14)	Poultry		Brazil/2009, 2011	NR	NR	[71]
		Chicken		Venezuela/2005–07, 2008	CIP-GEN	PL – NR	[82]
	CTX-M-14 (n=1)	Poultry		Brazil/2011	NR	NR	[71]
	SHV-2 (n=1)	Retail chicken meat	Yes	Canada/2007	GEN-STR-SUL	PL – II	[83]
	CMY-2 (n=25)	Chicken or chicken retail		Canada/2001–04	CHL-GEN-KAN-STR-SUL-SXT-TET	PL – A/C, II	[67]
		Chicken food		USA/2002, 2004, 2010	(CHL-GEN-KAN-STR-SUL-TET)	PL – A/C, II	[31]
Indiana	CTX-M-14 (n=21)	Poultry		Brazil/2011	NR	NR	[71]
		Poultry		China/2012–13	ENR-NAL-SXT-(AMK-CHL-CIP-FFC-GEN-LVX-OLA-TET)	PL – HI2, A/C, F, N, P-1 α , B/O	[70]
	CTX-M-24 (n=29)	Chicken		China/2008–09	AMK-CHL-FFC-GEN-NAL-OLA-RIF-STR-SXT-(TET)	PL – HI2	[84]
	CTX-M-65 (n=3)	Poultry		China/2012–13	CHL-NAL-OLA-SXT-(CIP-ENR-FFC-GEN-LVX-TET)	PL – HI2, F, N	[70]
	TEM-52 (n=1)	Broilers		The Netherlands/2006	(–)	PL – NT	[65]
	DHA-1 (n=3)	Chicken faeces		South Korea/2006–07	SUL-TMP-(APM-NAL-NEO-STR-TET)	PL – NR	[85]
Infantis	CTX-M-1 (n=2)	Poultry (caecum or unknown)		The Netherlands/2006	SUL-TMP	PL – II	[65,66]
	CTX-M-2 (n=3)	Poultry		Brazil/2005	NR	NR	[71]
		Retail Chicken Products		Japan/2002-03	(–)	NR	[86]
		Retail Chicken Products		Japan/1997, 2003	(–)	NR	[86]
		Poultry	Yes	Belgium/2001–05	(–)	PL – II	[62]
		Poultry (caecum or unknown)		The Netherlands/2006	(–)	PL – II	[65,66]
		Broiler		Belgium/2004	(–)	PL – II	[63]
		Chicken		Japan/2004–06	(–)	PL – NR	[87]
		Retail Chicken Products		Japan/2000, 2003	(–)	NR	[86]
		Retail Chicken Products		Japan/1997, 1999–2003	(–)	PL – NR	[86]
Kentucky	CTX-M-25/OXA-21 (n=1)	Turkey		Poland/2009	CIP-NAL	PL – A/C	[42]
	SHV-12 (n=3)	Whole chicken and chicken neck skin		Ireland/2008–09	CHL-SUL-TET	PL – NR	[88]
	CMY-2 (n=3)	Whole chicken		Ireland/2009	(–)	PL – NR	[88]
		Chicken meat		Germany/2005	STR	PL – II	[73]
Kiambu	SHV-2 (n=1)	Abattoir chicken cecum		Canada/2006	STR-SUL-SXT	PL – II	[83]
Livingstone	SHV-12 (n=9)	Poultry or turkey carcasses, poultry faeces, broiler faeces		Italy/2005-06	GEN-NAL-(STR-SUL)	NR	[75]
Llandoff	CTX-M-1 (n=1)	Poultry		France/2006	SUL-TET	PL – II	[89]
Manhattan	CTX-M-2 (n=2)	Retail Chicken Products		Japan/2000, 2003	(–)	NR	[86]

Clinical Microbiology and Infection © 2015 European Society of Clinical Microbiology and Infectious Diseases. Published by Elsevier Ltd. All rights reserved, CMI, xxx, 1–12

Please cite this article in press as: Antunes P, et al., Salmonellosis: the role of poultry meat, Clinical Microbiology and Infection (2016), <http://dx.doi.org/10.1016/j.cmi.2015.12.004>

ARTICLE IN PRESS

CMI

Antunes et al. Salmonellosis: the role of poultry meat 7

TABLE 2. Continued

Salmonella serotype	ESBL or AmpC ^a enzymes (no. of isolates)	Source	Concomitant presence in humans ^b	Country(ies)/ year(s)	Non-β-lactam resistance phenotype ^c	Genetic element PL – Inc group	Reference(s)
Minnesota	CTX-M-15 (n=4)	Retail Chicken Products		Japan/2002, 2003	(–)	NR	[86]
	TEM-52 (n=4)	Retail Chicken Products		Japan/2000, 2002–03	(–)	NR	[86]
	SHV-12 (n=2)	Retail Chicken Products		Japan/2002, 2003	(–)	NR	[86]
	CMY-2 (n=2)	Retail Chicken Products		Japan/2002	(–)	PL – NR	[86]
	CTX-M-2 (n=1)	Poultry		Brazil/2009	NR	NR	[71]
	CTX-M-8 (n=1)	Poultry		Brazil/2008	NR	NR	[71]
Newport	CTX-M-14 (n=1)	Poultry		Brazil/2010	NR	NR	[71]
	CMY-2 (n=2)	Poultry		Brazil/2011	NR	NR	[71]
	CTX-M-2 (n=1)	Poultry		Brazil/2008	NR	NR	[71]
	CTX-M-1 (n=6)	Turkey		USA/2011	NR	PL – N	[76]
	CTX-M-1 (n=1)	Poultry		Brazil/2009	NR	NR	[71]
	CTX-M-2 (n=1)	Poultry		Brazil/2009	NR	NR	[71]
Poona	CTX-M-8+CTX-M-14 (n=1)	Poultry		Brazil/2005	NR	NR	[71]
Rissen	CMY-2 (n=1)	Poultry		Brazil/2005	NR	NR	[71]
Saintpaul	CMY-2 (n=1)	Poultry		Brazil/2011	NR	NR	[71]
Schwarzengrund	CTX-M-2 (n=10)	Turkey, chicken, chicken carcass		Brazil/2008	CIP-NAL-STR-TET	PL – P	[72]
Senftenberg	CTX-M-15 (n=1)	Chicken		South Korea/2009	NR	PL – FII	[80]
	CTX-M-1 (n=1)	Poultry meat		Germany/2006	KAN-NEO-STR-SUL-SXT-TET-TMP	PL – II	[73]
	CTX-M-2 (n=10)	Poultry	Yes	Brazil/2004–05, 2007	NR	NR	[71]
	CTX-M-8 (n=1)	Poultry		Brazil/2003–04	SUL-SXT-TET	PL – NR	[90]
	CTX-M-14 (n=1)	Poultry		Brazil/2011	NR	NR	[71]
	CTX-M-15 (n=1)	Poultry		Brazil/2005	NR	NR	[71]
Typhimurium	TEM-52 (n=4)	Broiler	Yes	Belgium/2007	(–)	PL – II	[63]
		Poultry		Belgium/2002, France/2004	(–)	PL – II	[62]
		Poultry		The Netherlands/2001–02	SPT-SUL-TMP-(NEO-TET)	NR	[74]
		Poultry meat		The Netherlands/2002	(–)	NR	[74]
	TEM-52+SHV-12 (n=1)	Poultry		Belgium/2000–01, 2003	NAL-SUL-TET-TMP-(STR)	PL – HI2	[61]
	CTX-M-2 (n=12)	Poultry or poultry product		Ireland/UN	SUL-TET-TMP	PL – P	[77]
Virchow		Broiler	Yes	Belgium/2001, 2004	STR-SUL-TET-TMP	PL – HI2	[63]
		Poultry		Belgium-France/2000–03	NAL-SUL-TMP-(STR-TET)	PL – NR	[59]
	CTX-M-2+SHV-2+TEM-1 (n=1)	Broiler		The Netherlands/2002	NAL-SPT-SUL-TET-TMP	NR	[74]
	CTX-M-9 (n=8)	Chicken faeces		France/2002–03	KAN-NAL-SPT-STR-SUL-TET-TMP	PL – NR	[58]
		Chicken faeces and retail chicken meat	Yes	France/2002–03, Spain/2000	NAL-SPT-STR-SUL-TET-TMP-(KAN)	PL – HI2	[61]
	CTX-M-32 (n=2)	Broiler	Yes	Spain/2000–04	STR-SUL-SXT-NAL-(TET)	NR	[60,64]
Weslaco 35:c:1,2	TEM-52 (n=1)	Poultry		The Netherlands/2002	SPT-STR-SUL-TMP	NR	[74]
	CTX-M-8 (n=1)	Poultry		Brazil/2009	NR	NR	[71]
	SHV-12 (n=5)	Poultry	Yes	Senegal/2000	CHL-GEN-TET-TOB	NR	[92]

Antimicrobial abbreviations: AMK, amikacin; APM, apramycin; CHL, chloramphenicol; CIP, ciprofloxacin; ENR, enrofloxacin; FFC, florfenicol; GEN, gentamicin; KAN, kanamycin; LYX, levofloxacin; NAL, nalidixic acid; NEO, neomycin; OLA, olaquinox; RIF, rifampicin; SPT, spectinomycin; STR, streptomycin; SUL, sulphonamides compound; SXT, trimethoprim/sulfamethoxazole; TET, tetracycline; TMP, trimethoprim; TOB, tobramycin; NR, not reported; NT, plasmids that were not typable with the scheme used; PL, plasmid.

^aOnly references with full-characterized extended-spectrum β-lactamases /AmpC genes were considered.

^bYes, clones or serotypes simultaneously detected in poultry and humans in the same study.

^c(–), Non-β-lactam resistance phenotype was not detected using the tested antibiotics and according to the susceptibility criteria adopted. Variable phenotypes were present between curved brackets.

top MDR serotypes in both poultry and humans (frequently associated with invasive infections) in North America, presented increasing resistance to cephalosporins related with CMY-2 [31,67]. In fact, transmission of these most reported genes in both human and poultry has been associated with diverse plasmid families, such as IncI1 (e.g. *bla*_{CTX-M-1}, *bla*_{TEM-52} and *bla*_{CMY-2}), IncA/C (e.g. *bla*_{CMY-2}) or IncHI2 (e.g. *bla*_{CTX-M-2} and *bla*_{CTX-M-9}) (Table 2). Recently, plasmid-mediated carbapenem-resistant bacteria from food-animals have been reported, alerting us to a new public health problem with indefinable risks [51]. One of the first reports was on a poultry farm in Germany where a *bla*_{VIM-1} gene was detected on *S.*

Infantis [68]. Besides the worldwide resistance to clinically important β-lactams, plasmid-mediated quinolone resistance mechanisms, which typically confer decreased susceptibility to ciprofloxacin, have been widely reported in *Salmonella* [57]. Among them, Qnr proteins (e.g. QnrB2, QnrB19, QnrS1) have been commonly described in different serotypes and geographic locations, with the aminoglycoside acetyltransferase AAC(6′)-Ib-cr and the efflux pump OqxAB associated with *Salmonella* serotypes recovered from Asian poultry and poultry meat (Table 3). In fact, poultry seems to be an important vehicle of non-typhoidal *Salmonella* carrying plasmid-mediated quinolone resistance genes, highlighting the role of food-producing

TABLE 3. *Salmonella* serotypes/clones carrying plasmid-mediated quinolone resistance genes recovered from poultry and poultry products

Salmonella serotype	PMRQ ^a mechanism (no. of Isolates)	Source	Country(ies)/year(s)	Antibiotic resistance phenotype ^b	Genetic element PL – Inc group	Reference(s)
Agona	QnrB2 (n=1)	Turkeys	Germany/NR	NR	NR	[69]
Braenderup	QnrD (n=2)	Fowls	Spain/NR	NR	NR	[69]
Dabou	QnrD (n=1)	Fowls	Spain/NR	NR	NR	[69]
Derby	QnrB2 (n=56)	Fowls or turkeys	Spain/NR	NR	NR	[69]
Enteritidis	QnrB2 (n=1)	Fowls or turkeys	Spain/NR	NR	NR	[69]
	QnrB10/B19 (n=1)	Laying hen flock	Poland/2009	CIP-NAL	NR	[93]
	QnrD (n=3)	Fowls	Spain/NR	NR	NR	[69]
	QnrS1/S3 (n=2)	Broiler meat, broiler flock – faeces	Poland/2008–09	AMP-CIP-(STR-TET)	NR	[93]
	OqxAB (n=3)	Poultry	China/2012–13	AMP-CHL-CIF-CIP-ENR-FFC-NAL-OLA-SXT-(GEN-LVX-TET)	PL – HI2, F, N, B/O	[70]
	OqxAB + AAC(6′)-Ib-cr (n=1)	Poultry	China/2012–13	AMK-AMP-CHL-CIF-CIP-CTX-ENR-FFC-GEN-LVX-NAL-OLA-SXT-TET	PL – HI2	[70]
Give	QnrB19 (n=1)	Imported turkey meat from Brazil	Finland/NR	NR	NR	[69]
Hadar	QnrB2 (n=2)	Fowls or turkeys	Spain/NR	NR	NR	[69]
	QnrB5 (n=4)	Imported turkey meat	Germany/2007	STR-TET	NR	[94]
	QnrB19 (n=15)	Fowls, turkeys	Germany, Denmark/NR	NR	NR	[69]
Havana	QnrB2 (n=3)	Poultry	Portugal/2009–10	(–) or NAL	PL – L/M	[95]
	QnrB19 (n=1)	Poultry	Portugal/2009–10	(–)	PL – HI2	[95]
Heidelberg	QnrB19 (n=2)	Chicken	Venezuela/2005–07, 2008	CIP-GEN	PL – NR	[82]
Indiana	OqxAB (n=50)	Poultry	China/2012–13	AMP-CIF-NAL-SXT-(AMK-CAZ-CHL-CIP-CTX-ENR-FFC-GEN-LVX-OLA-TET)	PL – HI2, F, N, P-Ia, B/O	[70]
		Chicken	China/2008–09	AMK-CHL-FFC-GEN-NAL-OLA-RIF-STR-SXT-(TET)	PL – HI2	[84]
	OqxAB + AAC(6′)-Ib-cr (n=3)	Poultry	China/2012–13	AMP-CHL-CIF-NAL-SXT-(CIP-CTX-ENR-FFC-GEN-LVX-OLA-TET)	PL – HI2, A/C, N	[70]
Infantis	QnrS1 (n=1)	Chicken	Germany/2004	AMP	NR	[96]
	QnrB19 (n=1)	Retail chicken	Colombia/2004	KAN-NAL-NEO-STR-TET	PL – CoIE like	[97]
Kentucky	QnrS1 (n=1)	Chicken	The Netherlands/NR	NR	PL – N	[98]
London	QnrB2 (n=22)	Fowls or turkeys	Spain/NR	NR	NR	[69]
Mbandaka	QnrB19 (n=1)	Poultry	Portugal/2009–10	NAL	PL – HI2	[95]
Montevideo	QnrB2 (n=3)	Fowls or turkeys	Spain/NR	NR	NR	[69]
	QnrD (n=6)	Fowls, turkeys	Italy/NR	NR	NR	[69]
Newport	QnrB5 (n=3)	Imported turkey meat	Poland/2007	(–)	NR	[94]
	QnrS1/S3 (n=12)	Broiler (meat, flock), turkey meat, goose flock – faeces, duck flock – faeces	Poland/2008–11	CIP-(AMP-CHL-FFC-KAN-STR-SUL-TET-NAL)	NR	[93]
	QnrB19 (n=3)	Turkeys	Denmark/NR	NR	NR	[69]
Ohio	QnrD (n=5)	Fowls	Spain/NR	NR	NR	[69]
Saintpaul	QnrS1 (n=15)	Imported turkey meat	Germany/2007–08 Poland/2008	AMC-AMP-STR-TET-(CPD-CHL)	NR	[94]
		Turkeys	Germany, Denmark/NR	NR	NR	[69]
Typhimurium	QnrA1 (n=1)	Turkeys	Germany/NR	NR	NR	[69]
	QnrD (n=2)	Fowls	Spain/NR	NR	NR	[69]
	OqxAB (n=2)	Chicken	China/2007–09	OLA-FFC-SMX-(AMP-CHL-CIF-GEN-NAL-TET)	PL – HI2, F	[99]
	OqxAB + AAC(6′)-Ib-cr (n=8)	Chicken, duck	China/2009–10	NAL-OLA-SMX-(AMP-CHL-CIF-ENR-FFC-GEN-TET)	PL – HI2	[99]
	OqxAB + AAC(6′)-Ib-cr + QnrS1 (n=1)	Duck	China/2010	AMP-CHL-CIF-FFC-GEN-NAL-OLA-SMX-TET	PL – HI2	[99]
Virchow	QnrS1	Chicken carcass	Korea/2002	AMP-CEF-STR-SUL-TET-TMP	N	[100]
		Chicken	UK/2004-05	AMP-CIP	N	[101]
		Chicken	Turkey/2005	AMP-NAL	PL – NT	[102]

Antimicrobial abbreviations: AMC, amoxicillin/clavulanate; AMK, amikacin; AMP, ampicillin; CAZ, ceftazidime; CEF, cefalotin; CHL, chloramphenicol; CIF, ceftiofur; CIP, ciprofloxacin; CPD, cefpodoxime; CTX, cefotaxime; ENR, enrofloxacin; FFC, florfenicol; GEN, gentamicin; KAN, kanamycin; LVX, levofloxacin; NAL, nalidixic acid; NEO, neomycin; OLA, olaquinox; RIF, rifampicin; STR, streptomycin; SUL, sulphonamides compound; SXT, trimethoprim/sulfamethoxazole; TET, tetracycline; TMP, trimethoprim; NR, not reported; NT, plasmids that were not typeable with the scheme used; PL, plasmid.

^aOnly references with full-characterized plasmid-mediated quinolone resistance genes were considered.

^b(–), Antibiotic resistance phenotype was not detected using the tested antibiotics and according to the susceptibility criteria adopted. Variable phenotypes were present between curved brackets.

animals, including the animal/food trade, in its dissemination [69,70].

In addition to the therapeutic consequences, antimicrobial resistance acquisition to agents commonly used in livestock

production has been associated with the spread of particular *Salmonella* clones in the poultry production environment, which might contribute to their association with human diseases. Interestingly, the high-level resistance to ciprofloxacin (due to

ARTICLE IN PRESS

CMI

Antunes et al. Salmonellosis: the role of poultry meat 9

gyrA/*parC* mutations) and other antibiotics (amoxicillin, streptomycin, spectinomycin, gentamicin, sulfamethoxazole and tetracycline) [39–43] of the previously mentioned *S. Kentucky* ST198-XI-SGII strain might also have contributed to its clonal expansion. In the EU, several *S. Stanley* outbreaks linked with the consumption of turkey meat products have been reported, with the clone presenting resistance to nalidixic acid and decreased susceptibility to ciprofloxacin. Additionally, in the recent outbreak in Austria three strains also presented resistance to gentamicin and cephalosporins (CTX-M-15) [36,38]. Several successful *S. Infantis* clones of broiler origin are also characterized by multidrug resistance profiles, including nalidixic acid, tetracyclines, sulphonamides or nitrofurans [26,35,46], antibiotics frequently used in livestock production, which could contribute to their co-selection in the poultry industry.

Data compiled here, highlight the role of the poultry production chain (including poultry meat) as a reservoir of epidemic MDR clones or genes conferring resistance to critical antibiotics, which might spread to humans through the food chain. Globalization of food-animal production is currently posing a major challenge in antimicrobial resistance of zoonotic bacteria like non-typhoidal *Salmonella* with a consequent involvement of human health. In particular in the poultry production pyramid it is crucial to restrict the global use of antimicrobials to minimize the selection of resistant *Salmonella* (from the top of the poultry production pyramid and within flocks), and also to control the spread of MDR epidemic clones by improving biosecurity measures from the farm (e.g. hygiene high standards, vaccination) throughout the food chain (e.g. slaughtering and processing of poultry meat) and control of animal/food trade.

Conclusions

In the last decade, *Salmonella* control programmes, mainly targeting poultry production, have led to a significant reduction in salmonellosis in different countries, including in the EU. This positive effect has been associated with a shift in *Salmonella* serotypes of poultry and human disease, associated with a marked reduction on *S. Enteritidis* and the increase of less common serotypes, driven by the dissemination of particular clones carrying features favouring host (poultry/human) adaptability and frequently MDR.

Poultry has been reported as a source of non-typhoidal *Salmonella* resistant to clinically relevant antibiotics with a remarkable higher incidence reported for middle-income countries, which due to food trade globalization can dramatically challenge worldwide the treatment of severe salmonellosis cases.

Integrated surveillance (collaboration between human health, food safety and animal health—the ‘One Health’ approach) and containment strategies (including farms, retail, catering and consumers) to minimize contamination and reduce the transmission of *Salmonella* (including epidemic clones) along the food chain (from primary production to consumption) are mandatory on a global scale, in particular in the poultry production chain. In addition, continuous surveillance of *Salmonella* resistance levels globally is critical for clinicians to support the best salmonellosis treatment choices, avoid treatment failures, particularly for patients that would benefit from empirical antibiotic treatment.

Continuous monitoring to detect the emergence of any serotype or new clone along the food chain is of critical importance for public health, warning of the emergence of new *Salmonella* food safety risks, involving foodstuffs like poultry meat, which is one of the most consumed and increasing globally traded meat products.

Transparency Declaration

The authors declare no conflict of interest.

References

- [1] Parry CM, Threlfall EJ. Antimicrobial resistance in typhoidal and nontyphoidal salmonellae. *Curr Opin Infect Dis* 2008;21:531–8.
- [2] Chen HM, Wang Y, Su LH, Chiu CH. Nontyphoid *Salmonella* infection: microbiology, clinical features, and antimicrobial therapy. *Pediatr Neonatol* 2013;54:147–52.
- [3] Crump JA, Sjolund-Karlsson M, Gordon MA, Parry CM. Epidemiology, clinical presentation, laboratory diagnosis, antimicrobial resistance, and antimicrobial management of invasive *Salmonella* infections. *Clin Microbiol Rev* 2015;28:901–37.
- [4] Barrow PA, Jones MA, Smith AL, Wigley P. The long view: *Salmonella*—the last forty years. *Avian Pathol* 2012;41:413–20.
- [5] Cosby DE, Cox NA, Harrison MA, Wilson JL, Buhr RJ, Fedorka-Cray PJ. *Salmonella* and antimicrobial resistance in broilers: a review. *J Appl Poultry Res* 2015;24:408–26.
- [6] Majowicz SE, Musto J, Scallan E, Angulo FJ, Kirk M, O'Brien SJ, et al. The global burden of nontyphoidal *Salmonella* gastroenteritis. *Clin Infect Dis* 2010;50:882–9.
- [7] Scallan E, Griffin PM, Angulo FJ, Tauxe RV, Hoekstra RM. Foodborne illness acquired in the united states—unspecified agents. *Emerg Infect Dis* 2011;17:16–22.
- [8] EFSA European Food Safety Authority. The european union summary report on trends and sources of zoonoses, zoonotic agents and foodborne outbreaks in 2013. *EFSA J* 2015a;13:165. <http://dx.doi.org/10.2903/j.efsa.2015.3991>. Available at: www.efsa.europa.eu/en/efsajournal/pub/3991 (last accessed 1 September 2015).
- [9] CDC Centers for Disease Control and Prevention. National Enteric Disease Surveillance: *Salmonella* Annual Report, 2012. 2014. Available at: www.cdc.gov/ncezid/dfwed/pdfs/salmonella-annual-report-2012-508c.pdf (last accessed 1 September 2015).

ARTICLE IN PRESS

10 *Clinical Microbiology and Infection*, Volume ■ Number ■, ■■■ 2016

CMI

- [10] Hugas M, Beloeil P. Controlling *Salmonella* along the food chain in the European Union – progress over the last ten years. *Euro Surveill* 2014;19.
- [11] Butaye P, Michael GB, Schwarz S, Barrett TJ, Brisabois A, White DG. The clonal spread of multidrug-resistant non-typhi *Salmonella* serotypes. *Microbes Infect* 2006;8:1891–7.
- [12] Foley SL, Nayak R, Hanning IB, Johnson TJ, Han J, Ricke SC. Population dynamics of *Salmonella enterica* serotypes in commercial egg and poultry production. *Appl Environ Microbiol* 2011;77:4273–9.
- [13] Antunes P, Machado J, Peixe L. Characterization of antimicrobial resistance and class 1 and 2 integrons in *Salmonella enterica* isolates from different sources in Portugal. *J Antimicrob Chemother* 2006;58:297–304.
- [14] Mazengia E, Samadpour M, Hill HW, Greeson K, Tenney K, Liao G, et al. Prevalence, concentrations, and antibiotic sensitivities of *Salmonella* serovars in poultry from retail establishments in Seattle, Washington. *J Food Prot* 2014;77:885–93.
- [15] USDA/FSIS United States Department of Agriculture/Food Safety and Inspection Service. Progress report on *Salmonella* and *Campylobacter* testing of raw meat and poultry products, 1998–2013. 2014. Available at: www.fsis.usda.gov/wps/wcm/connect/885647f4-2568-48bf-ae5c-4a0d8279f435/Progress-Report-Salmonella-Campylobacter-CY2013.pdf?MOD=AJPERES (last accessed 1 September 2015).
- [16] Antunes P, Réu C, Sousa JC, Peixe L, Pestana N. Incidence of *Salmonella* from poultry products and their susceptibility to antimicrobial agents. *Int J Food Microbiol* 2003;82:97–103.
- [17] EFSA European Food Safety Authority. Scientific opinion on a quantitative estimation of the public health impact of setting a new target for the reduction of *Salmonella* in broilers. *EFSA J* 2011;9:2106. <http://dx.doi.org/10.2903/j.efsa.2011.2106>. Available at: www.efsa.europa.eu/en/efsajournal/pub/2106 (last accessed 1 December 2015).
- [18] EFSA European Food Safety Authority. Scientific opinion on an estimation of the public health impact of setting a new target for the reduction of *Salmonella* in turkeys. *EFSA J* 2012;10:2616. <http://dx.doi.org/10.2903/j.efsa.2012.2616>. Available at: www.efsa.europa.eu/en/efsajournal/pub/2616 (last accessed 1 December 2015).
- [19] Ribeiro A, Kellermann A, Bessa M, dos Santos L, do Nascimento V. *Salmonella* spp. in raw broiler parts: Occurrence, antimicrobial resistance profile and phage typing of the *Salmonella* Enteritidis isolates. *Braz J of Microbiol* 2007;38:296–9.
- [20] Donado-Godoy P, Byrne BA, Leon M, Castellanos R, Vanegas C, Coral A, et al. Prevalence, resistance patterns, and risk factors for antimicrobial resistance in bacteria from retail chicken meat in Colombia. *J Food Prot* 2015;78:751–9.
- [21] Adeyanju GT, Ishola O. *Salmonella* and *Escherichia coli* contamination of poultry meat from a processing plant and retail markets in Ibadan, Oyo State, Nigeria. *Springerplus* 2014;3:139.
- [22] Abd-Elghany SM, Sallam KI, Abd-Elkhalek A, Tamura T. Occurrence, genetic characterization and antimicrobial resistance of *Salmonella* isolated from chicken meat and giblets. *Epidemiol Infect* 2015;143:997–1003.
- [23] Ta YT, Nguyen TT, To PB, Pham da X, Le HT, Thi GN, et al. Quantification, serovars, and antibiotic resistance of *Salmonella* isolated from retail raw chicken meat in Vietnam. *J Food Prot* 2014;77:57–66.
- [24] Yang B, Cui Y, Shi C, Wang J, Xia X, Xi M, et al. Counts, serotypes, and antimicrobial resistance of *Salmonella* isolates on retail raw poultry in the people's Republic of China. *J Food Prot* 2014;77:894–902.
- [25] Yoon RH, Cha SY, Wei B, Roh JH, Seo HS, Oh JY, et al. Prevalence of *Salmonella* isolates and antimicrobial resistance in poultry meat from South Korea. *J Food Prot* 2014;77:1579–82.
- [26] Gal-Mor O, Valinsky L, Weinberger M, Guy S, Jaffe J, Schorr YI, et al. Multidrug-resistant *Salmonella enterica* serovar Infantis, Israel. *Emerg Infect Dis* 2010;16:1754–7.
- [27] Clemente L, Manageiro V, Ferreira E, Jones-Dias D, Correia I, Themudo P, et al. Occurrence of extended-spectrum β -lactamases among isolates of *Salmonella enterica* subsp. *enterica* from food-producing animals and food products, in Portugal. *Int J Food Microbiol* 2013;167:221–8.
- [28] BCCDC BC Center for Disease Control; BC Integrated Surveillance of Foodborne Pathogens (BCISFP). *Salmonella findings* 2013. 2014. Available at: www.bccdc.ca/NR/rdonlyres/3872B775-8B56-459B-9EE9-4DB7774C8D5B/0/2013BC_IntegratedSalmAnnualReport.pdf (last accessed 1 September 2015).
- [29] Figueiredo R, Henriques A, Sereno R, Mendonça N, da Silva GJ. Antimicrobial resistance and extended-spectrum β -lactamases of *Salmonella enterica* serotypes isolated from livestock and processed food in Portugal: an update. *Foodborne Pathog Dis* 2015;12:110–7.
- [30] Dutil L, Irwin R, Finley R, Ng LK, Avery B, Boerlin P, et al. Ceftiofur resistance in *Salmonella enterica* serovar Heidelberg from chicken meat and humans, Canada. *Emerg Infect Dis* 2010;16:48–54.
- [31] Hoffmann M, Zhao S, Pettengill J, Luo Y, Monday SR, Abbott J, et al. Comparative genomic analysis and virulence differences in closely related *Salmonella enterica* serotype Heidelberg isolates from humans, retail meats, and animals. *Genome Biol Evol* 2014;6:1046–68.
- [32] Dhanani AS, Block G, Dewar K, Forgetta V, Topp E, Beiko RG, et al. Genomic comparison of non-typhoidal *Salmonella enterica* serovars Typhimurium, Enteritidis, Heidelberg, Hadar and Kentucky isolates from broiler chickens. *PLoS One* 2015;10:e0128773.
- [33] Nogrady N, Toth A, Kostyak A, Paszti J, Nagy B. Emergence of multidrug-resistant clones of *Salmonella* Infantis in broiler chickens and humans in Hungary. *J Antimicrob Chemother* 2007;60:645–8.
- [34] Nogrady N, Kardos G, Bistyak A, et al. Prevalence and characterization of *Salmonella* Infantis isolates originating from different points of the broiler chicken-human food chain in Hungary. *Int J Food Microbiol* 2008;127:162–7.
- [35] Nogrady N, Kiraly M, Davies R, Nagy B. Multidrug resistant clones of *Salmonella* Infantis of broiler origin in Europe. *Int J Food Microbiol* 2012;157:108–12.
- [36] ECDC/EFSA European Centre for Disease Prevention and Control/ European Food Safety Authority. Multi-country outbreak of *Salmonella* Stanley infections 2014. 2014. Available at: <http://ecdc.europa.eu/en/publications/Publications/salmonella-stanley-multi-country-outbreak-assessment-8-May-2014.pdf> (last accessed 1 September 2015).
- [37] Kinross P, van Alphen L, Martinez Urtaza J, Struelens M, Takkinen J, Coulombier D, et al. Multidisciplinary investigation of a multicountry outbreak of *Salmonella* Stanley infections associated with turkey meat in the European Union, August 2011 to January 2013. *Euro Surveill* 2014;19.
- [38] Springer B, Allerberger F, Kornschöber C. Letter to the editor: *Salmonella* Stanley outbreaks—a prompt to reevaluate existing food regulations. *Euro Surveill* 2014;19.
- [39] Le Hello S, Hendriksen RS, Doublet B, Fisher I, Nielsen EM, Whitchard JM, et al. International spread of an epidemic population of *Salmonella enterica* serotype Kentucky ST198 resistant to ciprofloxacin. *J Infect Dis* 2011;204:675–84.
- [40] Le Hello S, Bekhit A, Granier SA, Barua H, Beutlich J, Zajac M, et al. The global establishment of a highly-fluoroquinolone resistant *Salmonella enterica* serotype Kentucky ST198 strain. *Front Microbiol* 2013;4:395.
- [41] Westrell T, Monnet DL, Gossner C, Heuer O, Takkinen J. Drug-resistant *Salmonella enterica* serotype Kentucky in Europe. *Lancet Infect Dis* 2014;14:270–1.
- [42] Wasyl D, Kern-Zdanowicz I, Domanska-Blicharz K, Zajac M, Hosiowski A. High-level fluoroquinolone resistant *Salmonella enterica* serovar Kentucky ST198 epidemic clone with IncA/C conjugative plasmid carrying *bla_{CTX-M-25}* gene. *Vet Microbiol* 2015;175:85–91.

Clinical Microbiology and Infection © 2015 European Society of Clinical Microbiology and Infectious Diseases. Published by Elsevier Ltd. All rights reserved, CMI, xxx, 1–12

Please cite this article in press as: Antunes P, et al., Salmonellosis: the role of poultry meat, *Clinical Microbiology and Infection* (2016), <http://dx.doi.org/10.1016/j.cmi.2015.12.004>

ARTICLE IN PRESS

CMI

Antunes et al. Salmonellosis: the role of poultry meat 11

- [43] EFSA European Food Safety Authority. Eu Summary Report on antimicrobial resistance in zoonotic and indicator bacteria from humans, animals and food in 2013. EFSA J 2015b;13:4036. <http://dx.doi.org/10.2903/j.efsa.2015.4036>. Available at: www.efsa.europa.eu/en/efsajournal/pub/4036 (last accessed 1 September 2015).
- [44] Antunes P, Mourão J, Pestana N, Peixe L. Leakage of emerging clinically relevant multidrug-resistant *Salmonella* clones from pig farms. J Antimicrob Chemother 2011;66:2028–32.
- [45] Gomes-Neves E, Antunes P, Manageiro V, Gartner F, Caniça M, da Costa JM, et al. Clinically relevant multidrug resistant *Salmonella enterica* in swine and meat handlers at the abattoir. Vet Microbiol 2014;168:229–33.
- [46] Aviv G, Tsyba K, Steck N, Salmon-Divon M, Cornelius A, Rahav G, et al. A unique megaplasmid contributes to stress tolerance and pathogenicity of an emergent *Salmonella enterica* serovar Infantis strain. Environ Microbiol 2014;16:977–94.
- [47] WHO World Health Organization. Critically important antimicrobials for human medicine, 3rd revision, 2012. Available at: http://apps.who.int/iris/bitstream/10665/77376/1/9789241504485_eng.pdf?ua=1&ua=1 (last accessed 1 September 2015).
- [48] BIOHAZ EFSA. Panel on Biological Hazards; Joint Opinion on antimicrobial resistance (AMR) focused on zoonotic infections. EFSA J 2009;7:1372. <http://dx.doi.org/10.2903/j.efsa.2009.1372>. Available at: www.efsa.europa.eu/en/efsajournal/pub/1372 (last accessed 1 September 2015).
- [49] BIOHAZ EFSA. Panel on Biological Hazards. Scientific Opinion on the public health risks of bacterial strains producing extended-spectrum β -lactamases and/or AMP β -lactamases in food and food-producing animals. EFSA J 2011;9:2322. <http://dx.doi.org/10.2903/j.efsa.2011.2322>. Available at: www.efsa.europa.eu/en/efsajournal/pub/2322 (last accessed 1 September 2015).
- [50] Aarestrup FM. The livestock reservoir for antimicrobial resistance: a personal view on changing patterns of risks, effects of interventions and the way forward. Philos Trans R Soc Lond B Biol Sci 2015;370:20140085.
- [51] Seiffert SN, Hilty M, Perreten V, Endimiani A. Extended-spectrum cephalosporin-resistant gram-negative organisms in livestock: an emerging problem for human health? Drug Resist Updat 2013;16:22–45.
- [52] Antunes P, Machado J, Peixe L. Illegal use of nitrofurans in food animals: contribution to human salmonellosis? Clin Microbiol Infect 2006;12:1047–9.
- [53] CDC/USDA/FDA Centers for Disease Control and Prevention/U.S. Department of Agriculture/Food and Drug Administration. The National Antimicrobial Resistance Monitoring System (NARMS): The 2012–2013 Integrated NARMS Report. 2015. Available at: <http://www.fda.gov/downloads/AnimalVeterinary/SafetyHealth/AntimicrobialResistance/NationalAntimicrobialResistanceMonitoringSystem/UCM453387.pdf> (last accessed 1 September 2015).
- [54] Lee HY, Su LH, Tsai MH, Kim SW, Chang HH, Jung SI, et al. High rate of reduced susceptibility to ciprofloxacin and ceftriaxone among nontyphoid *Salmonella* clinical isolates in Asia. Antimicrob Agents Chemother 2009;53:2696–9.
- [55] Van TT, Nguyen HN, Smooker PM, Coloe PJ. The antibiotic resistance characteristics of non-typhoidal *Salmonella enterica* isolated from food-producing animals, retail meat and humans in South East Asia. Int J Food Microbiol 2012;154:98–106.
- [56] Van Boeckel TP, Brower C, Gilbert M, Grenfell Bryan T, Levin Simon A, Robinson Timothy P, et al. Global trends in antimicrobial use in food animals. Proc Natl Acad Sci 2015;112:5649–54.
- [57] Rodríguez-Martínez JM, Cano ME, Velasco C, Martínez-Martínez L, Pascual A. Plasmid-mediated quinolone resistance: an update. J Infect Chemother 2011;17:149–82.
- [58] Weill FX, Lailier R, Praud K, Kerouanton A, Fabre L, Brisabois A, et al. Emergence of extended-spectrum- β -lactamase (CTX-M-9)-producing multiresistant strains of *Salmonella enterica* serotype Virchow in poultry and humans in France. J Clin Microbiol 2004;42:5767–73.
- [59] Bertrand S, Weill FX, Cloeckaert A, Vrints M, Mairiaux E, Praud K, et al. Clonal emergence of extended-spectrum β -lactamase (CTX-M-2)-producing *Salmonella enterica* serovar Virchow isolates with reduced susceptibilities to ciprofloxacin among poultry and humans in Belgium and France (2000 to 2003). J Clin Microbiol 2006;44:2897–903.
- [60] Riano I, Moreno MA, Teshager T, Saenz Y, Dominguez L, Torres C. Detection and characterization of extended-spectrum β -lactamases in *Salmonella enterica* strains of healthy food animals in Spain. J Antimicrob Chemother 2006;58:844–7.
- [61] García Fernández A, Cloeckaert A, Bertini A, Praud K, Doublet B, Weill FX, et al. Comparative analysis of IncHI2 plasmids carrying *bla*_{CTX-M-2} or *bla*_{CTX-M-9} from *Escherichia coli* and *Salmonella enterica* strains isolated from poultry and humans. Antimicrob Agents Chemother 2007;51:4177–80.
- [62] Cloeckaert A, Praud K, Doublet B, Bertini A, Carattoli A, Butaye P, et al. Dissemination of an extended-spectrum- β -lactamase *bla*_{TEM-52} gene-carrying IncI plasmid in various *Salmonella enterica* serovars isolated from poultry and humans in Belgium and France between 2001 and 2005. Antimicrob Agents Chemother 2007;51:1872–5.
- [63] Smet A, Martel A, Persoons D, Dewulf J, Heyndrickx M, Cloeckaert A, et al. Comparative analysis of extended-spectrum- β -lactamase-carrying plasmids from different members of *Enterobacteriaceae* isolated from poultry, pigs and humans: evidence for a shared β -lactam resistance gene pool? J Antimicrob Chemother 2009;63:1286–8.
- [64] Riano I, García-Campello M, Saenz Y, Alvarez P, Vinue L, Lantero M, et al. Occurrence of extended-spectrum β -lactamase-producing *Salmonella enterica* in northern Spain with evidence of CTX-M-9 clonal spread among animals and humans. Clin Microbiol Infect 2009;15:292–5.
- [65] Dierikx C, van Essen-Zandbergen A, Veldman K, Smith H, Mevius D. Increased detection of extended spectrum β -lactamase producing *Salmonella enterica* and *Escherichia coli* isolates from poultry. Vet Microbiol 2010;145:273–8.
- [66] Leverstein-van Hall MA, Dierikx CM, Cohen Stuart J, Voets GM, van den Munckhof MP, van Essen-Zandbergen A, et al. Dutch patients, retail chicken meat and poultry share the same ESBL genes, plasmids and strains. Clin Microbiol Infect 2011;17:873–80.
- [67] Andrysiak AK, Olson AB, Tracz DM, Dore K, Irwin R, Ng LK, et al. Genetic characterization of clinical and agri-food isolates of multi drug resistant *Salmonella enterica* serovar Heidelberg from Canada. BMC Microbiol 2008;8:89.
- [68] Fischer J, Rodríguez I, Schmoger S, Friese A, Roesler U, Helmuth R, et al. *Salmonella enterica* subsp. *enterica* producing VIM-1 carbapenemase isolated from livestock farms. J Antimicrob Chemother 2013;68:478–80.
- [69] Veldman K, Cavaco LM, Mevius D, Battisti A, Franco A, Botteldoorn N, et al. International collaborative study on the occurrence of plasmid-mediated quinolone resistance in *Salmonella enterica* and *Escherichia coli* isolated from animals, humans, food and the environment in 13 European Countries. J Antimicrob Chemother 2011;66:1278–86.
- [70] Li L, Liao XP, Liu ZZ, Huang T, Li X, Sun J, et al. Co-spread of *oqxAB* and *bla*_{CTX-M-9G} in non-typhi *Salmonella enterica* isolates mediated by ST2-IncHI2 plasmids. Int J Antimicrob Agents 2014;44:263–8.
- [71] Fitch FM, Carmo-Rodrigues MS, Oliveira VG, Gaspari MV, Dos Santos A, de Freitas JB, et al. β -lactam resistance genes: characterization, epidemiology, and first detection of *bla*_{CTX-M-1} and *bla*_{CTX-M-14} in *Salmonella* spp. isolated from poultry in Brazil-Brazil Ministry of Agriculture's Pathogen Reduction Program. Microb Drug Resist 2015.

Clinical Microbiology and Infection © 2015 European Society of Clinical Microbiology and Infectious Diseases. Published by Elsevier Ltd. All rights reserved, CMI, xxx, 1–12

Please cite this article in press as: Antunes P, et al., Salmonellosis: the role of poultry meat, Clinical Microbiology and Infection (2016), <http://dx.doi.org/10.1016/j.cmi.2015.12.004>

ARTICLE IN PRESS

12 *Clinical Microbiology and Infection*, Volume ■ Number ■, ■■■ 2016

CMI

- [72] Silva KC, Fontes LC, Moreno AM, Astolfi-Ferreira CS, Ferreira AJ, Lincopan N. Emergence of extended-spectrum- β -lactamase CTX-M-2-producing *Salmonella enterica* serovars Schwarzengrund and Agona in poultry farms. *Antimicrob Agents Chemother* 2013;57:3458–9.
- [73] Rodriguez I, Barownick W, Helmuth R, Mendoza MC, Rodicio MR, Schroeter A, et al. Extended-spectrum β -lactamases and AmpC β -lactamases in ceftiofur-resistant *Salmonella enterica* isolates from food and livestock obtained in Germany during 2003–07. *J Antimicrob Chemother* 2009;64:301–9.
- [74] Hasman H, Møvius D, Veldman K, Olesen I, Aarestrup FM. β -lactamases among extended-spectrum β -lactamase (ESBL)-resistant *Salmonella* from poultry, poultry products and human patients in The Netherlands. *J Antimicrob Chemother* 2005;56:115–21.
- [75] Chiaretto G, Zavagnin P, Bettini F, Mancin M, Minorello C, Saccardin C, et al. Extended spectrum β -lactamase SHV-12-producing *Salmonella* from poultry. *Vet Microbiol* 2008;128:406–13.
- [76] Wittum TE, Mollenkopf DF, Erdman MM. Detection of *Salmonella enterica* isolates producing CTX-M cephalosporinase in U.S. livestock populations. *Appl Environ Microbiol* 2012;78:7487–91.
- [77] Hopkins KL, Liebana E, Villa L, Batchelor M, Threlfall EJ, Carattoli A. Replicon typing of plasmids carrying CTX-M or CMY β -lactamases circulating among *Salmonella* and *Escherichia coli* isolates. *Antimicrob Agents Chemother* 2006;50:3203–6.
- [78] Matsumoto Y, Kitazume H, Yamada M, Ishiguro Y, Muto T, Izumiya H, et al. CTX-M-14 type β -lactamase producing *Salmonella enterica* serovar Enteritidis isolated from imported chicken meat. *Jpn J Infect Dis* 2007;60:236–8.
- [79] Tamang MD, Nam HM, Kim TS, Jang GC, Jung SC, Lim SK. Emergence of extended-spectrum β -lactamase (CTX-M-15 and CTX-M-14)-producing nontyphoid *Salmonella* with reduced susceptibility to ciprofloxacin among food animals and humans in Korea. *J Clin Microbiol* 2011;49:2671–5.
- [80] Kang MS, Kwon YK, Oh JY, Kim MJ, Call DR, An BK, et al. Evidence for recent acquisition and successful transmission of *bla*_{CTX-M-15} in *Salmonella enterica* in South Korea. *Antimicrob Agents Chemother* 2013;57:2383–7.
- [81] Clemente L, Manageiro V, Jones-Dias D, Correia I, Themudo P, Albuquerque T, et al. Antimicrobial susceptibility and oxymino- β -lactam resistance mechanisms in *Salmonella enterica* and *Escherichia coli* isolates from different animal sources. *Res Microbiol* 2015;166:574–83.
- [82] Gonzalez F, Araque M. Association of transferable quinolone resistance determinant *qnrB19* with extended-spectrum β -lactamases in *Salmonella* GIVE and *Salmonella* Heidelberg in Venezuela. *Int J Microbiol* 2013;2013:628185.
- [83] Pouget JG, Coutinho FJ, Reid-Smith RJ, Boerlin P. Characterization of *bla*_{SHV} genes on plasmids from *Escherichia coli* and *Salmonella enterica* isolates from Canadian food animals (2006–2007). *Appl Environ Microbiol* 2013;79:3864–6.
- [84] Lai J, Wang Y, Shen J, Li R, Han J, Foley SL, et al. Unique class I integron and multiple resistance genes co-located on IncH12 plasmid is associated with the emerging multidrug resistance of *Salmonella* Indiana isolated from chicken in China. *Foodborne Pathog Dis* 2013;10:581–8.
- [85] Rayamajhi N, Jung BY, Cha SB, Shin MK, Kim A, Kang MS, et al. Antibiotic resistance patterns and detection of *bla*_{DHA-1} in *Salmonella* species isolates from chicken farms in South Korea. *Appl Environ Microbiol* 2010;76:4760–4.
- [86] Noda T, Murakami K, Etoh Y, Okamoto F, Yatsuyanagi J, Sera N, et al. Increase in resistance to extended-spectrum cephalosporins in *Salmonella* isolated from retail chicken products in Japan. *PLoS One* 2015;10:e0116927.
- [87] Shahada F, Chuma T, Dahshan H, Akiba M, Sueyoshi M, Okamoto K. Detection and characterization of extended-spectrum β -lactamase (TEM-52)-producing *Salmonella* serotype Infantis from broilers in Japan. *Foodborne Pathog Dis* 2010;7:515–21.
- [88] Boyle F, Morris D, O'Connor J, Delappe N, Ward J, Cormican M. First report of extended-spectrum- β -lactamase-producing *Salmonella enterica* serovar Kentucky isolated from poultry in Ireland. *Antimicrob Agents Chemother* 2010;54:551–3.
- [89] Cloeckaert A, Praud K, Lefevre M, Doublet B, Pardo M, Granier SA, et al. IncI plasmid carrying extended-spectrum- β -lactamase gene *bla*_{CTX-M-1} in *Salmonella enterica* isolates from poultry and humans in France, 2003 to 2008. *Antimicrob Agents Chemother* 2010;54:4484–6.
- [90] Fernandes SA, Paterson DL, Ghilardi-Rodrigues AC, Adams-Haduch JM, Tavechio AT, Doi Y. CTX-M-2-producing *Salmonella* Typhimurium isolated from pediatric patients and poultry in Brazil. *Microb Drug Resist* 2009;15:317–21.
- [91] Politi L, Tassios PT, Lambiri M, Kansouzidou A, Pasiotou M, Vatopoulos AC, et al. Repeated occurrence of diverse extended-spectrum β -lactamases in minor serotypes of food-borne *Salmonella enterica* subsp. *enterica*. *J Clin Microbiol* 2005;43:3453–6.
- [92] Cardinale E, Colbachini P, Perrier-Gros-Claude JD, Gassama A, Aidara-Kane A. Dual emergence in food and humans of a novel multiresistant serotype of *Salmonella* in Senegal: *Salmonella enterica* subsp. *enterica* serotype 35:c:1,2. *J Clin Microbiol* 2001;39:2373–4.
- [93] Wasyl D, Hosiowski A, Zajac M. Prevalence and characterisation of quinolone resistance mechanisms in *Salmonella* spp. *Vet Microbiol* 2014;171:307–14.
- [94] Cavaco LM, Korsgaard H, Sorensen G, Aarestrup FM. Plasmid-mediated quinolone resistance due to *qnrB5* and *qnrS1* genes in *Salmonella enterica* serovars Newport, Hadar and Saintpaul isolated from turkey meat in Denmark. *J Antimicrob Chemother* 2008;62:632–4.
- [95] Jones-Dias D, Manageiro V, Francisco AP, Martins AP, Domingues G, Louro D, et al. Assessing the molecular basis of transferable quinolone resistance in *Escherichia coli* and *Salmonella* spp. from food-producing animals and food products. *Vet Microbiol* 2013;167:523–31.
- [96] Kehrenberg C, Friederichs S, de Jong A, Michael GB, Schwarz S. Identification of the plasmid-borne quinolone resistance gene *qnrS* in *Salmonella enterica* serovar Infantis. *J Antimicrob Chemother* 2006;58:18–22.
- [97] Karczmarczyk M, Martins M, McCusker M, Mattar S, Amaral L, Leonard N, et al. Characterization of antimicrobial resistance in *Salmonella enterica* food and animal isolates from Colombia: Identification of a *qnrB19*-mediated quinolone resistance marker in two novel serovars. *FEMS Microbiol Lett* 2010;313:10–9.
- [98] Dolejska M, Villa L, Hasman H, Hansen L, Carattoli A. Characterization of IncN plasmids carrying *bla*_{CTX-M-1} and *qnr* genes in *Escherichia coli* and *Salmonella* from animals, the environment and humans. *J Antimicrob Chemother* 2013;68:333–9.
- [99] Li L, Liao X, Yang Y, Sun J, Li L, Liu B, et al. Spread of *oqxAB* in *Salmonella enterica* serotype Typhimurium predominantly by IncH12 plasmids. *J Antimicrob Chemother* 2013;68:2263–8.
- [100] Kim JH, Cho JK, Kim KS. Prevalence and characterization of plasmid-mediated quinolone resistance genes in *Salmonella* isolated from poultry in Korea. *Avian Pathol* 2013;42:221–9.
- [101] Hopkins KL, Wootton L, Day MR, Threlfall EJ. Plasmid-mediated quinolone resistance determinant *qnrS1* found in *Salmonella enterica* strains isolated in the UK. *J Antimicrob Chemother* 2007;59:1071–5.
- [102] Avsaroglu MD, Helmuth R, Junker E, Hertwig S, Schroeter A, Akcelik M, et al. Plasmid-mediated quinolone resistance conferred by *qnrS1* in *Salmonella enterica* serovar Virchow isolated from Turkish food of avian origin. *J Antimicrob Chemother* 2007;60:1146–50.

II. Protocols optimized for assessment of metal and biocide susceptibility by genotypic and/or phenotypic methods

Polymerase chain reaction (PCR) assays for detecting metal tolerance genes

TABLE 1 - Primers used in PCR assays to search acquired metal tolerance genes

Operon or cluster	Target Gene	Gene coding for	Primer	Primer sequence (5'-3') ^a	Nucleotide position (GenBank accession no.)	Reference
<i>sil</i>	<i>silA</i>	Silver inner-membrane proton/cation antiporter	<i>silA</i> _Fw	GCAAGACCGGTAAAGCAGAG	8936-8955	Mourão <i>et al.</i> , 2015
			<i>silA</i> _Rv	CCTGCCAGTACAGGAACCAT	9871-9852	
	<i>silC</i>	Outer membrane protein	<i>silC</i> _4090_F	CGGGCTGGCGHAMCTTTTGTG	(AF067954.1) 4090-4110	This study
			<i>silC</i> _4470_R	CCAGTTGCTGRTGAAATARC	4470-4450 (BX664015.1)	
	<i>silE</i>	Silver/copper periplasmic binding protein	<i>silE</i> _1105_F	GTTCTGTCATGTYTCATGAGC	1105-1125	Mourão <i>et al.</i> , 2015
			<i>silE</i> _1368_R	GTACTYCCCCGGACATCAATAATT	1368-1345 (AF067954.1)	
	<i>silP</i>	P-type cation ATPase	<i>silP</i> _11882_F	GGCGATAAGCTCCGCATCAGA	11882-11902	Kremer <i>et al.</i> , 2014
			<i>silP</i> _12405_R	TCCACTTTTTCAGACGCTCA	12405-12385 (BX664015.1)	
	<i>silR</i>	Transcriptional regulator responder	<i>silR</i> _3244_F	CCCTGATGGCGAAKGAAGAA	3244-3263	Kremer <i>et al.</i> , 2014
			<i>silR</i> _3572_R	AACGGCTGGGATATCRTCSCG	3572-3553 (BX664015.1)	
<i>pco</i>	<i>pcoA</i>	Multicopper oxidase	<i>pcoA</i> _978_F	CTCGCGGATGTCAGTGGCTACACCT	978-1002	Mourão <i>et al.</i> , 2015
			<i>pcoA</i> _1481_R	ATCCGGAAGGTCAGCACCGTCCATAGAC	1481-1454 (X83541.1)	
	<i>pcoD</i>	Copper inner membrane pump	<i>pcoD</i> _F	CTGGCCACACTTGCTGGGG	3801-3820	Mourão <i>et al.</i> , 2015
			<i>pcoD</i> _R	CACGCTACGGCGCCGAGAAT	4300-4281 (X83541.1)	
	<i>pcoE</i>	Periplasmic chaperone	<i>pcoE</i> _6585_F	CCTGGTTCTCGAGTGATGA	6585-6604	This study
			<i>pcoE</i> _7057_R	TGACCAITATTTCGCTTCT	7057-7039 (X83541.1)	
	<i>pcoR</i>	Regulator of <i>pcoABCD</i> cluster	<i>pcoR</i> _4460_F	AGGCTATCAGCCCGATCTCT	4460-4478	This study
			<i>pcoR</i> _4928_R	CCAGACCAGGGACGAGATAA	4928-4909 (X83541.1)	
	<i>pcoS</i>	Sensor of <i>pcoABCD</i> cluster	<i>pcoS</i> _5182_F	TAATCAGGACCGCGATTTC	5182-5201	This study
			<i>pcoS</i> _6236_R	CACTGTACGGCTCAAGGTGT	6236-6217	

Operon or cluster	Target Gene	Gene coding for	Primer	Primer sequence (5'-3') ^a	Nucleotide position (GenBank accession no.)	Reference
ter	<i>terB</i>	Copper export ATPase	terB_F	CATACGGTAGCTTTAAGGAGATTTC	(X83541.1) 4954-4980	Hasman <i>et al.</i> , 2006
			terB_R	ATAGAGGACTCCGCCACCATG	5616-5595 (AY048044.2)	
mer	<i>merA</i>	Mercuric reductase	merA_1F	ACCATCGGGGGCACCCTGCGT	2140-2159	Liebert <i>et al.</i> , 1997
			merA_5R	ACCATCGTCAGGTAGGGGAAC	3377-3357 (K03089.1)	
ars	<i>arsB</i>	Arsenite transmembrane pump	arsB_Fw	ACTGAAAGACAGACGAAAGCG	159735-159754	García-Fernández <i>et al.</i> , 2012
			arsB_Rv	GGCAGATAGTGTGGAATGCG	160870-160851 (BX664015.1)	
ter	<i>terF</i>	Tellurite resistance protein	terF_Fw1	ATAGCACTGGATCGTGTTC	80174-80193	This study
			terF_Rv	TTCATCGATCCACGGTCTG	81163-81145 (BX664015.1)	

^aH = A, C or T; K = G or T; M = A or C; R = A or G; S = G or C; Y = C or T.

TABLE 2 - Conditions used for the PCR assays to search acquired metal tolerance genes

Operon or cluster	Target Gene	Primer	Taq Polymerase kit used (V _{final} = 25 µl)	Amplification conditions	Annealing temperature (°C)	Amplicon size (bp)
sil	<i>silA</i>	<i>silA_Fw</i>	Buffer 0.5x	94°C - 10 minutes (1 cycle)	59	936
		<i>silA_Rv</i>	2mM of MgCl ₂	94° - 1 minute		
	<i>silC</i>	<i>silC_4090_F</i>	200 µM of each dNTP	T _A ° - 1 minute	60	381
		<i>silC_4470_R</i>	0.2 µM of each Primer	72° - 2 minutes (30 cycles)		
	<i>silE</i>	<i>silE_1105_F</i>	KAPA Taq DNA Polymerase 1U	72° - 10 minutes (1 cycle)	62	264
		<i>silE_1368_R</i>				
	<i>silP</i>	<i>silP_11882_F</i>			60	524
		<i>silP_12405_R</i>				
	<i>silR</i>	<i>silR_3244_F</i>			62	329
		<i>silR_3572_R</i>				
pco		<i>silS_2279_F</i>			60	741
		<i>silS_3019_R</i>				
	<i>pcoA</i>	<i>pcoA_978_F</i>	Buffer 1x	95°C - 10 minutes (1 cycle)	60	504
		<i>pcoA_1481_R</i>	1.5mM of MgCl ₂	94° - 30 seconds		

Operon or cluster	Target Gene	Primer	Taq Polymerase kit used (V _{final} = 25 µl)	Amplification conditions	Annealing temperature (°C)	Amplicon size (bp)
	<i>pcoD</i>	pcoD_F pcoD_R	200 µM of each dNTP 0.2 µM of each Primer PROMEGA Go Taq® DNA Polymerase 0.5U	T _A ⁰ - 30 seconds 72° - 30 seconds (30 cycles) 72° - 10 minutes (1 cycle)	55	500
	<i>pcoE</i>	pcoE_6585_F pcoE_7057_R			58	473
	<i>pcoR</i>	pcoR_4460_F pcoR_4928_R			58	469
	<i>pcoS</i>	pcoS_5182_F pcoS_6236_R			63	1055
	<i>tcr</i>	tcrB_F tcrB_R	Buffer 1x 1.5 mM of MgCl ₂ 200 µM of each dNTP 0.125 µM of each Primer PROMEGA Go Taq® DNA Polymerase 1.25U	95°C- 10 min (1 cycle) 94°C-30 seconds T _A ⁰ -45 seconds 72°C-45 seconds (25 cycles) 72°C-10 minutes (1 cycle)	55	990
<i>mer</i>	<i>merA</i>	merA_1F merA_5R	Buffer 1x 1.5mM of MgCl ₂ 200 µM of each dNTP 0.2 µM of each Primer KAPA Taq DNA Polymerase 0.5U	95°C – 10 minutes (1 cycle) 94° - 30 seconds T _A ⁰ - 1 minute 72° - 2 minutes (30 cycles) 72° - 10 minutes (1 cycle)	65	1238
<i>ars</i>	<i>arsB</i>	arsB_Fw arsB_Rv	Buffer 1x 1.5mM of MgCl ₂ 200 µM of each dNTP 0.2 µM of each Primer PROMEGA Go Taq® DNA Polymerase 1U	95°C – 10 minutes (1 cycle) 94° - 30 seconds T _A ⁰ - 30 seconds 72° - 30 seconds (25 cycles) 72° - 10 minutes (1 cycle)	60	1136
<i>ter</i>	<i>terF</i>	terF_Fw1 terF_Rv	Buffer 1x 1.5mM of MgCl ₂ 200 µM of each dNTP 0.2 µM of each Primer PROMEGA Go Taq® DNA Polymerase 0.5U	95°C – 10 minutes (1 cycle) 94° - 1 minute T _A ⁰ - 1 minute 72° - 2 minutes (30 cycles) 72° - 10 minutes (1 cycle)	60	990

Appendix references:

Garcia Fernandez A, Cloeckaert A, Bertini A, Praud K, Doublet B, Weill FX, *et al.* Comparative analysis of IncHI2 plasmids carrying *bla*_{CTX-M-2} or *bla*_{CTX-M-9} from *Escherichia coli* and *Salmonella enterica* strains isolated from poultry and humans. *Antimicrob Agents Chemother.* 2007;51(11):4177-80.

Hasman H, Kempf I, Chidaine B, Cariolet R, Ersboll AK, Houe H, *et al.* Copper resistance in *Enterococcus faecium*, mediated by the *tcrB* gene, is selected by supplementation of pig feed with copper sulfate. *Appl Environ Microbiol.* 2006;72(9):5784-9.

Kremer AN, Hoffmann H. Subtractive hybridization yields a silver resistance determinant unique to nosocomial pathogens in the *Enterobacter cloacae* complex. *J Clin Microbiol.* 2012;50(10):3249-57.

Liebert CA, Wireman J, Smith T, Summers AO. Phylogeny of mercury resistance (*mer*) operons of gram-negative bacteria isolated from the fecal flora of primates. *Appl Environ Microbiol.* 1997;63(3):1066-76.

Mourão J, Novais C, Machado J, Peixe L, Antunes P. Metal tolerance in emerging clinically relevant multidrug-resistant *Salmonella enterica* serotype 4,[5],12:i:- clones circulating in Europe. *Int J Antimicrob Agents.* 2015;45(6):610-6.

Determination of copper sulphate and silver nitrate minimum inhibitory concentrations (MICs) by the agar dilution method in aerobic and anaerobic atmospheres

Strains:

Day 1

- a) Prepare and isolate test cultures and quality control strain cultures on CLED (MIC_{CuSO₄}) or Mueller-Hinton Agar/MHA (MIC_{AgNO₃}) and check purity before.

Control strains:

For CuSO₄ susceptibility assays

- *Enterococcus faecium* BM4105RF*
- *Escherichia coli* ED8739 (plasmid pRJ1004 with *pco* cluster) – offered by Jill Williams

For AgNO₃ susceptibility assays

- *S. aureus* ATCC 29213*
- *Escherichia coli* J53 (plasmid pMG101 with *sil* cluster) -offered by Christopher Randall

* Both negative for all genes tested (*pcoD*⁻, *silA*⁻, *tcrB*⁻, *cueO*⁻)

TABLE 3 – MICs values of quality control strains to CuSO₄ and AgNO₄ in aerobic (with/without induction) and anaerobic atmospheres

Control strains	O ₂ environment	<i>sil</i> genes	<i>pco</i> genes	MIC _{CuSO₄} (mM)	MIC _{AgNO₃} (mM)
<i>Enterococcus faecium</i> BM4105RF	Aerobic	(-)	(-)	8–12	
	Anaerobic			2–4	
<i>Escherichia coli</i> ED8739 (pRJ1004)	Aerobic	(+)	(+)	32–36	
	Anaerobic			16–20	
<i>S. aureus</i> ATCC 29213	Aerobic	(-)	(-)		0.16–0.25
	Induction <i>sil</i>				0.16–0.25
	Anaerobic				0.32–0.5
<i>Escherichia coli</i> J53 (pMG101)	Aerobic	(+)	(+)		> 3
	Induction <i>sil</i>				>3
	Anaerobic				> 3

Material:**Day 2**

- Prepare first Mueller-Hinton Agar 2, cation adjusted (Biomérieux – Ref. 51075) following manufacturer's instructions. We will need 25 mL of MHA per plate to prepare all the concentrations in the range to test and to prepare control plates for growth controls (MHA_{initial} and MHA_{final}).
- Autoclave the MHA at 121° C for 15 minutes.
- Cool to 60° C in water bath before aseptically add metal solutions.
- Pre-warm 50, 100 mL or 250 mL of measurement glasses (for mixing and pouring) in 60° C water bath until needed. We will need one Erlenmeyer flask for each concentration and graduated cylinders.
- After cooling down the media check pH using a pH meter, accordingly to the metal (7.2 – copper sulphate; 7.4 – silver nitrate) at room temperature before adding the metal solutions.

NOTE: We have **three systems** for MIC's determination: 2 systems of 21 wells and 1 system of 17 wells.

We will need the following concentrations for each metal:

1. Cu⁺/Cu²⁺ (aerobic and anaerobic conditions)

0.25 mM – 0.5 mM – 1 mM – 2 mM – 4 mM – 8 mM – 12 mM – 16 mM – 20 mM – 24 mM – 28 mM – 32 mM – 36 mM (**aerobic**)

0.25 mM – 0.5 mM – 1 mM – 2 mM – 4 mM – 8 mM – 12 mM – 16 mM – 20 mM – 24 mM – 28 mM – 32 mM – 36 mM (**anaerobic**)

Mueller-Hinton Agar for one system = (13 concentrations aerobic + 13 concentrations anaerobic + 4 plates for MH_{initial} and MH_{final}) x 25 mL agar = 750 mL MHA. Prepare MHA in excess (e.g. 900 mL).

2. Ag²⁺ (aerobic with/without induction and anaerobic conditions)

0.025 mM – 0.06 mM – 0.08 mM – 0.125 mM – 0.16 mM – 0.25 mM – 0.32 mM – 0.5 mM – 0.75 mM – 1 mM – 1.5 mM – 3 mM (**aerobic with/without induction**)

0.025 mM – 0.06 mM – 0.08 mM – 0.125 mM – 0.16 mM – 0.25 mM – 0.32 mM – 0.5 mM – 0.75 mM – 1 mM – 1.5 mM – 3 mM (**anaerobic**)

Mueller-Hinton Agar for one system = (12 concentrations aerobic + 12 concentrations anaerobic + 4 plates for $MH_{initial}$ and MH_{final}) x 25 mL agar = 700 mL MHA. Prepare MHA in excess (e.g. 900 mL).

Metal solutions:

...Day 2

The recommendation is preparing stock solutions from the compounds at least 10X the highest concentration of the range. Copper sulphate is prepared at 1M stock solution and silver nitrate at 10 mM stock solution in ultra-pure distilled water (Ex: if the highest concentration to test is 1mM the stock solution should be prepared at least at 10mM). See section above.

MW ($CuSO_4$) = 159.61 g/mol

MW ($AgNO_3$) = 169.87 g/mol

SAFETY- WEIGHING OF THE POWDER NEEDS TO BE DONE IN A HIGH SAFETY FUME HOOD WITH DOUBLE GLOVES) AND VERY CAREFULLY DUE TO THE POWDER TOXICITY! ADD WATER AS SOON AS POSSIBLE TO REDUCE CHANCES OF POWDER INHALATION!

Preparation of plates with copper sulphate, pH = 7.2

4 systems

PROCEDURE

A. Prepare 25 ml of the stock solution of $CuSO_4$ at 1M

$$c = \frac{n}{v} \leftrightarrow 1M = \frac{n}{0.025 L} \leftrightarrow n = 0.025 mol$$

$$n = \frac{m}{M_w} \leftrightarrow 0.025 mol = \frac{m}{159.61} \leftrightarrow m = 3.99025 g + 25 ml of distilled water$$

B. Make 2-fold dilutions – in 2 orders**1.**

24 mM – From 200 mL MH Agar, discard 4.8 mL of agar and add 4.8 mL of 1M Copper Sulphate stock solution mix and adjust pH to 7.2.

12 mM – Mix 50 ml 24 mM with 50 ml of MHA, mix and adjust pH to 7.2.

See the first example to apply to the rest of the concentrations:

$$C_c \times V_c = C_d \times V_d \leftrightarrow 1M \times V_c = 0.024 M \times 200 mL \leftrightarrow V_c = \mathbf{4.8 mL}$$

2.

16 mM – From 200 mL MH Agar, discard 3.2 mL of agar and add 3.2 mL of 1M Copper Sulphate stock solution mix and adjust pH to 7.2.

8 mM – Mix 100 ml 16 mM with 100 ml of MHA, mix and adjust pH to 7.2.

4 mM – Mix 100 ml 8 mM with 100 ml of MHA, mix and adjust pH to 7.2.

2 mM – Mix 100 ml 4 mM with 100 ml of MHA, mix and adjust pH to 7.2.

1 mM – Mix 100 ml 2 mM with 100 ml of MHA, mix and adjust pH to 7.2.

0.5 mM – Mix 100 ml 1 mM with 100 ml of MHA, mix and adjust pH to 7.2.

0.25 mM – Mix 50 ml 0.5 mM with 50 ml of MHA, mix and adjust pH to 7.2.

Separately prepare:

20 mM – From 100 mL MH Agar, discard 2.0 mL of agar and add 2.0 mL of 1M Copper Sulphate stock solution mix and adjust pH to 7.2.

28 mM – From 100 mL MH Agar, discard 2.8 mL of agar and add 2.8 mL of 1M Copper Sulphate stock solution mix and adjust pH to 7.2.

32 mM – From 100 mL MH Agar, discard 3.2 mL of agar and add 3.2 mL of 1M Copper Sulphate stock solution mix and adjust pH to 7.2.

36 mM – From 100 mL MH Agar, discard 3.6 mL of agar and add 3.6 mL of 1M Copper Sulphate stock solution mix and adjust pH to 7.2.

C. Prepare all dilutions using CuSO₄ at recommended concentrations as follows:

Concentration (mM)	Order 1	Order 2	Volume of copper sulphate (ml)	Volume of MHA (ml)
MH _{initial} + MH _{final}		Separately	0 mL	2 x 100 mL
2 x 0 mM				
0.25 mM		9	50 ml 0.5 mM agar	50 ml
0,5 mM		8	100 ml 1 mM agar	100 ml
1 mM		7	100 ml 2 mM agar	100 ml
2 mM		6	100 ml 4 mM agar	100 ml
4 mM		5	100 ml 8 mM agar	100 ml
8 mM		4	100 ml 16 mM agar	100 ml
12 mM	2		50 ml 24 mM agar	50 ml
16 mM		3	3.2 ml stock solution at 1M	200 ml
20 mM		Separately	2.0 ml stock solution at 1M	100 ml
24 mM	1		4.8 ml stock solution at 1M	200 ml
28 mM		Separately	2.8 ml stock solution at 1M	100 ml
32 mM		Separately	3.2 ml stock solution at 1M	100 ml
36 mM		Separately	3.6 ml stock solution at 1M	100 ml

Preparation of plates with silver nitrate, pH = 7.4

4 systems

PROCEDURE

D. Prepare 90 ml of the stock solution of AgNO₃ at 10 mM

$$c = \frac{n}{v} \leftrightarrow 0.010M = \frac{n}{0.090 L} \leftrightarrow n = 0.0009 \text{ mol}$$

$$n = \frac{m}{M_w} \leftrightarrow 0.0009 \text{ mol} = \frac{m}{169.87} \leftrightarrow m = 0.152883 \text{ g} + 90 \text{ ml of distilled water}$$

E. Prepare 200 ml of a solution AgNO₃ at 1 mM in Mueller-Hinton Agar

$$C_c \times V_c = C_d \times C_v \leftrightarrow 10 \text{ mM} \times V_c = 1 \text{ mM} \times 200 \text{ ml}$$

$$V_c = 20 \text{ ml of the 10 mM silver solution} + 180 \text{ ml of MHA, adjust pH to 7.4}$$

F. Prepare all dilutions using mix of agar at 1 mM and 10 mM as follows:

Concentration (mM) of stock solution	Volume of the AgNO ₃ stock solution (ml)	Volume of Mueller- Hinton Agar (ml)	Final concentration (mM)
			MH_{initial} + MH_{final}
0	0	2 x 100	2 x 0 mM
1	2.5	97.5	0.025 mM
1	6	94	0.06 mM
1	8	92	0.08 mM
1	12.5	87.5	0.125 mM
1	16	84	0.16 mM
1	25	75	0.25 mM
1	32	68	0.32 mM
1	50	50	0.5 mM
10	7.5	92.5	0.75 mM
10	10	90	1 mM
10	15	85	1.5 mM
10	30	70	3 mM

Preparation of plates:**...Day 2**

- Distribute Mueller-Hinton Agar with metals (Cu or Ag) into Petri dishes at depth of about 3 to 4 mm (± 25 ml each), mark metal concentration on plates inside LAF bench.
- Plates must solidify at room temperature, and must be used in the same day.
- Evaluate potency and stability with control strains in all rounds of MIC testing – all the control strains are in the first page.
- Prepare 2 (MH_{initial} and MH_{final}) control plates without metal solution as growth controls for each round.

Inoculum preparation:**...Day 2**

- Prepare a suspension at 0.5 McFarland ($1-2 \times 10^8$ CFU/ml) in 2 mL of sterile saline (NaCl) using a nephelometer.
- Dilute 1:10 in sterile saline to approximated 10^7 CFU/ml (100 μ l of the former suspension at 0.5 McF + 900 μ l saline) in 1.5 mL Eppendorf tubes before adding the suspension to the inoculum replicator block.

Inoculum replicator:**...Day 2**

Prepare the **scheme** of the block inoculum replicator including 21 inocula, the position of 19 test strains and the 2 control strains.

Remember that 20 and 21 positions are the controls.

- Inoculum suspension (10^7 CFU/ml prepared before) in 1.5 mL eppendorf tubes should be ordered in a rack and 500 μ l aliquots placed in the corresponding places in the replicator inoculum block.
- Agar plates need to be marked for orientation. ATTENTION VERY IMPORTANT: All the plates must be marked on the same side of the plate to allow to locate strains when reading. Mark the plate and NOT the cover.
- The samples are applied to the agar surface, always orientated in the same way and according to the marking applied, by use of an inocula-replicating device (or standardized loops or pipettes). This protocol is designed to obtain a final concentration of 10^4 CFU/mL.
- A growth control plate (without antimicrobial $MH_{initial}$) is inoculated in the first place and then the plates containing the antimicrobial, starting with the lowest concentration. After all the previous plates are inoculated, a second control plate is inoculated in last place (without antimicrobial MH_{final}) to ensure that there was no contamination or significant antimicrobial carry over during the inoculation.
- A sample of each inoculum is streaked on a suitable agar plate (Blood Agar or Mueller-Hinton agar sterility test) and incubated overnight to detect mixed cultures and to provide freshly isolated colonies in case retesting proves necessary.

The inoculator pins and blocks are disinfected with Virkon for 1 hour (3 liters of water + 2.5 spoons of the powder), then washed thoroughly with distilled water and scrubbed with a swab to guarantee that all disinfectant is removed, otherwise it could affect the growth of bacteria in the following assays. Finally pack and sent to autoclaving after use.

Incubating agar plates:**...Day 2**

AEROBIC: The inoculated plates are allowed to stand in a flow chamber until the moisture of inoculum has been absorbed into the agar, but **NO more than 30 minutes**. **!!!DO NOT INVERT THE PLATES!!!** Incubate at 37° C for 16-24h and then again after 48 hours.

ANAEROBIC ATMOSPHERE: After the inoculum was completely absorbed, incubate the plates in an anaerobic jar (42 plates = 7 litres) with four generators (GENbox anaer – Biomérieux Ref. 36124) and one methylene blue indicator strip (Anaer indicator – Biomérieux Ref. 36118). When incubating in an anaerobic condition is only at 37° C for 20-24h.

Reading and determining agar dilution end points:

...Day 2

After incubation, the plates should be placed on a dark non-reflecting surface to determine end points. The MIC is recorded as the lowest concentration of antimicrobial agent that completely inhibits growth disregarding a single colony or little growth caused by the inoculum.

- If two or more colonies persist after an obvious end point or there is no growth at lower concentrations but it occurs at higher concentrations, culture purity should be checked and the test repeated.
- Register all the results in terms of MIC determined and correspond to tested strains according to the previously prepared scheme for each round of tests and check results of control strains.

Induction assays of *sil* gene cluster

Day 1

1. All induction assays should be performed few days before starting the protocol for the determination of MICs to silver nitrate.
2. Prepare and isolate test cultures and quality control strain cultures (*E. coli* J53 plasmid pMG101) on MHA and check purity before performing the assay. Incubate in the static oven at 37°C.
3. Prepare BBL™ Mueller-Hinton Broth (BD/reference 212322) for using in all the steps. We will need 10 mL of MHB per test strains plus the controls until the end of the assay.

Day 2

- Pick up one to two pure colonies of the MHA plates (test isolates + control) and put them into a glass test tube containing 5 mL of MHB. Incubate in a static oven at 37°C for 14 hours.

Day 3

- Prepare 10 ml of the stock solution of AgNO₃ at 10 mM

$$c = \frac{n}{v} \leftrightarrow 0.010M = \frac{n}{0.010 L} \leftrightarrow n = 0.0001 \text{ mol}$$

$$n = \frac{m}{M_w} \leftrightarrow 0.0001 \text{ mol} = \frac{m}{169.87} \leftrightarrow m = 0.016987 \text{ g} + 10 \text{ ml of distilled water}$$

- Prepare 10 ml of a solution of AgNO₃ at 1 mM in water using the stock at 10 mM

$$C_c \times V_c = C_d \times C_v \leftrightarrow 10 \text{ mM} \times V_c = 1 \text{ mM} \times 10 \text{ ml}$$

$$V_c = 1 \text{ ml of the 10 mM silver solution} + 9 \text{ ml of water}$$

- Pick up the test tubes from the static oven and homogenise the suspension slowly. Take 100 µl from each tube containing the test strains and control, for two new glass test tubes containing:



Tube 1:
100 µl strain +
5 mL MHB



Tube 2 (induction):
100 µl strain +
5 mL MHB +
0.025 mM AgNO₃ (use the previously
prepared AgNO₃ solutions at 1 mM)*

* Prepare 0.025mM AgNO₃ using the solution at 1 mM:

$$C_c \times V_c = C_d \times C_v \leftrightarrow 1 \text{ mM} \times V_c = 0.025 \text{ mM} \times 5 \text{ ml}$$

$$V_c = 0.125 \text{ ml of the 1 mM silver solution} + 4.875 \text{ ml of MHB}$$


8. Incubate all the previous test tube in a shaking water bath with strong agitation (180 rpm) at 37°C for 16 hours.

Day 4

9. Pick up the test tube from the shaking water bath and proceed with determination of MICs to AgNO₃ following the standard protocol.

Determination of benzalkonium chloride-BZK minimum inhibitory concentrations by the broth microdilution method

The scheme below represents the main steps of the adopted protocol in this work and described in the items below.



Day 1

1. Prepare and isolate test cultures and quality control strain cultures on MHA and check purity before performing the assay.
2. Prepare BBL™ Mueller-Hinton Broth for using in all the steps, taking into account that in each of the 96 wells of the microtiter plate we will have a final volume of 100 µl. Additionally, we will need 20 mL of MHB for the dilution of the stock solution of BZK and 5 mL for each test strain/control.

Day 2

3. Prepare the stock solution of the biocide benzalkonium chloride (powder, B6295-Sigma-Aldrich®) according to the number of isolates that will be tested. In each microtiter plate we can test 6 strains (line A-F) plus the controls (line G-H). In each plate we can use the column 11 and 12 to control the growth of the bacterial suspension and the biocide (without any inoculated strain).

- Concentration range of BZK to use in the assay

Column	1	2	3	4	5	6	7	8	9	10	11	12
Concentration (mg/L)	256	128	64	32	16	8	4	2	1	0.5	C ⁺	C ⁻

- Prepare 1 mL of a stock solution of BZK at 100 mg/mL or g/L (Mw = 360 g/mol)

$$M \text{ (mol/L)} = \frac{C \left(\frac{\text{g}}{\text{L}} \right)}{M_w} \leftrightarrow M = \frac{100}{360} \leftrightarrow n = 0.27777777 \text{ mol/L}$$

$$c = \frac{n}{v} \leftrightarrow 0.27777777 = \frac{n}{0.001 \text{ L}} \leftrightarrow n = 0.0002777 \text{ mol}$$

$$n = \frac{m}{M_w} \leftrightarrow 0.0002777 \text{ mol} = \frac{m}{360} \leftrightarrow m = 0.099972 \text{ g} + 1 \text{ ml of distilled water}$$

4. Dilute with MHB the previous stock of BZK at 100 mg/mL in sterile polypropylene tubes to make a final solution at twice the maximum concentration used for the susceptibility testing (e.g. if your maximum concentration of BZK is 256 mg/L you will need to prepare a BZK solution in MHB at 512 mg/L).

- Prepare 20 mL of a MHB solution of BZK at 512 mg/L using the stock at 100 mg/mL

$$C_c \times V_c = C_d \times C_v \leftrightarrow 100 \text{ mg/mL} \times V_c = 0.512 \text{ mg/mL} \times 20 \text{ ml}$$

$$V_c = 0.1024 \text{ ml of the } 100 \text{ mg/mL BZK stock solution} + 19.8976 \text{ ml of MHB}$$

5. With a multichannel add 50 µl of MHB at each column with the exception of number 1. Then add 50 µl of BZK solution in MHB to columns 1 and 2. With the multichannel make serial dilutions starting at column 2 and ending at the last desired concentration. Discard the last 50 µl of the final concentration.
6. Prepare the inoculum suspension of each strains plus the controls. For the preparation of the inoculum suspension, use 2-3 pure colonies to make a final suspension in sterile saline (NaCl 0,9%) at 0.5 McFarland ($1,5 \times 10^8$ CFU/mL) using a photometric device/nephelometer. Dilute 1:100 the previous suspension to achieve 1×10^6 CFU/mL in 5 mL of MHB (e.g. 50 µl of the final suspension in 5 mL of MHB). The last suspension should be distributed (50 µl) in each well of the microtiter plate, optimally within 15 minutes.
7. Incubate the microtiter plate in a static oven at 37°C for 16-20 hours. To maintain the same temperature incubation for all cultures, the trays should not be staked more than 4 highs.

III. Permissions for inclusion of papers, figures and tables

12/01/2016

Rightslink® by Copyright Clearance Center



RightsLink®

Home

Account
Info

Help



Title: Leakage of emerging clinically relevant multidrug-resistant Salmonella clones from pig farms:

Logged in as:
Joana Mourao
Account #:
3000982054

Author: Patrícia Antunes, Joana Mourão, Nazaré Pestana, Luísa Peixe

[LOGOUT](#)

Publication: Journal of Antimicrobial Chemotherapy

Publisher: Oxford University Press

Date: 09/01/2011

Copyright © 2011, Oxford University Press

Order Completed

Thank you very much for your order.

This is a License Agreement between Joana Mourao ("You") and Oxford University Press ("Oxford University Press"). The license consists of your order details, the terms and conditions provided by Oxford University Press, and the [payment terms and conditions](#).

[Get the printable license.](#)

License Number	3786170804909
License date	Jan 11, 2016
Licensed content publisher	Oxford University Press
Licensed content publication	Journal of Antimicrobial Chemotherapy
Licensed content title	Leakage of emerging clinically relevant multidrug-resistant Salmonella clones from pig farms:
Licensed content author	Patrícia Antunes, Joana Mourão, Nazaré Pestana, Luísa Peixe
Licensed content date	09/01/2011
Volume number	66
Issue number	9
Type of Use	Thesis/Dissertation
Requestor type	Academic/Educational institute
Format	Print
Portion	Full article
Will you be translating?	No
Author of this OUP article	Yes
Order reference number	None
Title of your thesis / dissertation	CONTRIBUTION OF BIOCIDES TO THE SELECTION OF SALMONELLA NON-TYPHOID RESISTANT TO ANTIBIOTICS: A MULTILAYERED APPROACH
Expected completion date	Feb 2016
Estimated size(pages)	250
Publisher VAT ID	GB 125 5067 30
Total	0.00 EUR

[ORDER MORE...](#)
[CLOSE WINDOW](#)

Copyright © 2016 [Copyright Clearance Center, Inc.](#) All Rights Reserved. [Privacy statement](#). [Terms and Conditions](#).
Comments? We would like to hear from you. E-mail us at customer@copyright.com

15/01/2016

Rightslink® by Copyright Clearance Center



RightsLink®

Home

Account
Info

Help



Title: Characterization of the emerging clinically-relevant multidrug-resistant *Salmonella enterica* serotype 4,[5],12:i:- (monophasic variant of *S. Typhimurium*) clones

Author: J. Mourão

Publication: European Journal of Clinical Microbiology and Infectious Diseases

Publisher: Springer

Date: Jan 1, 2014

Logged in as:

Joana Mourao

Account #:

3000982054

LOGOUT

Copyright © 2014, Springer-Verlag Berlin Heidelberg

Order Completed

Thank you very much for your order.

This is a License Agreement between Joana Mourao ("You") and Springer ("Springer"). The license consists of your order details, the terms and conditions provided by Springer, and the [payment terms and conditions](#).

[Get the printable license.](#)

License Number	3787890799997
License date	Jan 14, 2016
Licensed content publisher	Springer
Licensed content publication	European Journal of Clinical Microbiology and Infectious Diseases
Licensed content title	Characterization of the emerging clinically-relevant multidrug-resistant <i>Salmonella enterica</i> serotype 4,[5],12:i:- (monophasic variant of <i>S. Typhimurium</i>) clones
Licensed content author	J. Mourão
Licensed content date	Jan 1, 2014
Volume number	33
Issue number	12
Type of Use	Thesis/Dissertation
Portion	Full text
Number of copies	10
Author of this Springer article	Yes and you are the sole author of the new work
Title of your thesis / dissertation	CONTRIBUTION OF BIOCIDES TO THE SELECTION OF <i>SALMONELLA</i> NON-TYPHOID RESISTANT TO ANTIBIOTICS: A MULTILAYERED APPROACH
Expected completion date	Feb 2016
Estimated size(pages)	250
Total	0.00 EUR

CLOSE WINDOW

Copyright © 2016 Copyright Clearance Center, Inc. All Rights Reserved. [Privacy statement](#). [Terms and Conditions](#). Comments? We would like to hear from you. E-mail us at customercare@copyright.com

18/01/2016

Rightslink® by Copyright Clearance Center



RightsLink®

Account
Info

Help

Live Chat



Title: Metal tolerance in emerging clinically relevant multidrug-resistant *Salmonella enterica* serotype 4,[5],12:i:– clones circulating in Europe

Author: Joana Mourão, Carla Novais, Jorge Machado, Luísa Peixe, Patrícia Antunes

Publication: International Journal of Antimicrobial Agents

Publisher: Elsevier

Date: Jun 1, 2015

Copyright © 2015, Elsevier

Logged in as:

Joana Mourao

Account #:

3000982054

LOGOUT

Order Completed

Thank you for your order.

This Agreement between Joana Mourao ("You") and Elsevier ("Elsevier") consists of your order details and the terms and conditions provided by Elsevier and Copyright Clearance Center.

License number	Reference confirmation email for license number
License date	Jan 14, 2016
Licensed content publisher	Elsevier
Licensed content publication	International Journal of Antimicrobial Agents
Licensed content title	Metal tolerance in emerging clinically relevant multidrug-resistant <i>Salmonella enterica</i> serotype 4,[5],12:i:– clones circulating in Europe
Licensed content author	Joana Mourão, Carla Novais, Jorge Machado, Luísa Peixe, Patrícia Antunes
Licensed content date	June 2015
Licensed content volume number	45
Licensed content issue number	6
Number of pages	7
Type of Use	reuse in a thesis/dissertation
Portion	full article
Format	print
Are you the author of this Elsevier article?	Yes
Will you be translating?	No
Title of your thesis/dissertation	CONTRIBUTION OF BIOCIDES TO THE SELECTION OF <i>SALMONELLA</i> NON-TYPHOID RESISTANT TO ANTIBIOTICS: A MULTILAYERED APPROACH
Expected completion date	Feb 2016
Elsevier VAT number	GB 494 6272 12
Billing Type	Invoice
Billing address	

Permissions price	0.00 EUR
VAT/Local Sales Tax	0.00 EUR / 0.00 GBP
Total	0.00 EUR

CLOSE WINDOW

Copyright © 2016 Copyright Clearance Center, Inc. All Rights Reserved. [Privacy statement](#). [Terms and Conditions](#). Comments? We would like to hear from you. E-mail us at customer@copyright.com

18/01/2016


Rightslink® by Copyright Clearance Center



RightsLink®

Account
Info

Help



Title: Salmonella enterica serotype Bovismorbificans, a new host for CTX-M-9

Author: Patrícia Antunes, Joana Mourão, Tatiana Alves, Joana Campos, Carla Novais, Ângela Novais, Luísa Peixe

Publication: International Journal of Antimicrobial Agents

Publisher: Elsevier

Date: Jan 1, 2013

Copyright © 2013, Elsevier

Logged in as:
Joana Mourao
Account #:
3000982054

LOGOUT

Order Completed

Thank you for your order.

This Agreement between Joana Mourao ("You") and Elsevier ("Elsevier") consists of your order details and the terms and conditions provided by Elsevier and Copyright Clearance Center.

License number	Reference confirmation email for license number
License date	Jan 14, 2016
Licensed content publisher	Elsevier
Licensed content publication	International Journal of Antimicrobial Agents
Licensed content title	Salmonella enterica serotype Bovismorbificans, a new host for CTX-M-9
Licensed content author	Patrícia Antunes, Joana Mourão, Tatiana Alves, Joana Campos, Carla Novais, Ângela Novais, Luísa Peixe
Licensed content date	January 2013
Licensed content volume number	41
Licensed content issue number	1
Number of pages	3
Type of Use	reuse in a thesis/dissertation
Portion	full article
Format	print
Are you the author of this Elsevier article?	Yes
Will you be translating?	No
Title of your thesis/dissertation	CONTRIBUTION OF BIOCIDES TO THE SELECTION OF SALMONELLA NON-TYPHOID RESISTANT TO ANTIBIOTICS: A MULTILAYERED APPROACH
Expected completion date	Feb 2016
Elsevier VAT number	GB 494 6272 12
Billing Type	Invoice
Billing address	

Permissions price	0.00 EUR
VAT/Local Sales Tax	0.00 EUR / 0.00 GBP
Total	0.00 EUR

CLOSE WINDOW

Copyright © 2016 Copyright Clearance Center, Inc. All Rights Reserved. [Privacy statement](#). [Terms and Conditions](#).
Comments? We would like to hear from you. E-mail us at customercare@copyright.com

15/01/2016

Rightslink® by Copyright Clearance Center



RightsLink®

Home

Account
Info

Help



Title: Clinical Salmonella Typhimurium ST34 with metal tolerance genes and an IncHI2 plasmid carrying oqxAB-aac(6')-Ib-cr from Europe:

Logged in as:
Joana Mourao
Account #:
3000982054

LOGOUT

Author: Joana Campos, Joana Mourão, Sara Marçal, Jorge Machado, Carla Novais, Luísa Peixe, Patrícia Antunes

Publication: Journal of Antimicrobial Chemotherapy

Publisher: Oxford University Press

Date: 12/17/2015

Copyright © 2015, Oxford University Press

Order Completed

Thank you very much for your order.

This is a License Agreement between Joana Mourao ("You") and Oxford University Press ("Oxford University Press"). The license consists of your order details, the terms and conditions provided by Oxford University Press, and the [payment terms and conditions](#).

[Get the printable license](#).

License Number	3787960688972
License date	Jan 14, 2016
Licensed content publisher	Oxford University Press
Licensed content publication	Journal of Antimicrobial Chemotherapy
Licensed content title	Clinical Salmonella Typhimurium ST34 with metal tolerance genes and an IncHI2 plasmid carrying oqxAB-aac(6')-Ib-cr from Europe:
Licensed content author	Joana Campos, Joana Mourão, Sara Marçal, Jorge Machado, Carla Novais, Luísa Peixe, Patrícia Antunes
Licensed content date	12/17/2015
Type of Use	Thesis/Dissertation
Requestor type	Academic/Educational institute
Format	Print
Portion	Full article
Will you be translating?	No
Author of this OUP article	Yes
Order reference number	None
Title of your thesis / dissertation	CONTRIBUTION OF BIOCIDES TO THE SELECTION OF SALMONELLA NON-TYPHOID RESISTANT TO ANTIBIOTICS: A MULTILAYERED APPROACH
Expected completion date	Feb 2016
Estimated size(pages)	250
Publisher VAT ID	GB 125 5067 30
Total	0.00 EUR

ORDER MORE...

CLOSE WINDOW

Copyright © 2016 [Copyright Clearance Center, Inc.](#) All Rights Reserved. [Privacy statement](#). [Terms and Conditions](#).
Comments? We would like to hear from you. E-mail us at customer@copyright.com

01/02/2016

Rightslink® by Copyright Clearance Center



RightsLink®

Account
Info

Help

Live Chat


Title: CMI Themed Issue "Poultry and human infections" (2016) Salmonellosis: the role of poultry-meat

Author: Patrícia Antunes, Joana Mourão, Joana Campos, Luísa Peixe

Publication: Clinical Microbiology and Infection

Publisher: Elsevier

Date: Dec 31, 1969

Copyright © 1969, Elsevier

Logged in as:
Joana Mourao
Account #:
3000982054

[LOGOUT](#)

Order Completed

Thank you for your order.

This Agreement between Joana Mourao ("You") and Elsevier ("Elsevier") consists of your order details and the terms and conditions provided by Elsevier and Copyright Clearance Center.

License number	Reference confirmation email for license number
License date	Jan 31, 2016
Licensed content publisher	Elsevier
Licensed content publication	Clinical Microbiology and Infection
Licensed content title	CMI Themed Issue "Poultry and human infections" (2016) Salmonellosis: the role of poultry-meat
Licensed content author	Patrícia Antunes, Joana Mourão, Joana Campos, Luísa Peixe
Licensed content date	Available online 17 December 2015
Licensed content volume number	n/a
Licensed content issue number	n/a
Number of pages	1
Type of Use	reuse in a thesis/dissertation
Portion	full article
Format	print
Are you the author of this Elsevier article?	Yes
Will you be translating?	No
Title of your thesis/dissertation	CONTRIBUTION OF BIOCIDES TO THE SELECTION OF SALMONELLA NON-TYPHOID RESISTANT TO ANTIBIOTICS: A MULTILAYERED APPROACH
Expected completion date	Feb 2016
Elsevier VAT number	GB 494 6272 12
Billing Type	
Billing address	

Permissions price	0.00 EUR
VAT/Local Sales Tax	0.00 EUR / 0.00 GBP
Total	0.00 EUR

[CLOSE WINDOW](#)

Copyright © 2016 [Copyright Clearance Center, Inc.](#) All Rights Reserved. [Privacy statement](#). [Terms and Conditions](#).
Comments? We would like to hear from you. E-mail us at customercare@copyright.com

17/12/2015

Rightslink® by Copyright Clearance Center



RightsLink®

Home

Account
Info

Help



Title: Extended-spectrum cephalosporin-resistant gram-negative organisms in livestock: An emerging problem for human health?

Logged in as:
Joana Mourao
Account #:
3000982054

LOGOUT

Author: Salome N. Seiffert, Markus Hilty, Vincent Perreten, Andrea Endimiani

Publication: Drug Resistance Updates

Publisher: Elsevier

Date: February–April 2013

Copyright © 2013 Elsevier Ltd. All rights reserved.

Order Completed

Thank you very much for your order.

This is a License Agreement between Joana Mourao ("You") and Elsevier ("Elsevier"). The license consists of your order details, the terms and conditions provided by Elsevier, and the [payment terms and conditions](#).

[Get the printable license.](#)

License Number	3771590158837
License date	Dec 17, 2015
Licensed content publisher	Elsevier
Licensed content publication	Drug Resistance Updates
Licensed content title	Extended-spectrum cephalosporin-resistant gram-negative organisms in livestock: An emerging problem for human health?
Licensed content author	Salome N. Seiffert, Markus Hilty, Vincent Perreten, Andrea Endimiani
Licensed content date	February–April 2013
Licensed content volume number	16
Licensed content issue number	1-2
Number of pages	24
Type of Use	reuse in a thesis/dissertation
Portion	figures/tables/illustrations
Number of figures/tables/illustrations	1
Format	both print and electronic
Are you the author of this Elsevier article?	No
Will you be translating?	No
Original figure numbers	Table 2. Antimicrobials approved for use in food-producing animals
Title of your thesis/dissertation	CONTRIBUTION OF BIOCIDES TO THE SELECTION OF SALMONELLA NON-TYPHOID RESISTANT TO ANTIBIOTICS: A MULTILAYERED APPROACH
Expected completion date	Feb 2016
Estimated size (number of pages)	250
Elsevier VAT number	GB 494 6272 12
Permissions price	0.00 EUR
VAT/Local Sales Tax	0.00 EUR / 0.00 GBP
Total	0.00 EUR

ORDER MORE...

CLOSE WINDOW

Copyright © 2015 [Copyright Clearance Center, Inc.](#) All Rights Reserved. [Privacy statement](#). [Terms and Conditions](#).
Comments? We would like to hear from you. E-mail us at customercare@copyright.com

11/12/2015

Rightslink® by Copyright Clearance Center



RightsLink®

Home

Account
Info

Help



Title: *Salmonella*, a cross-kingdom pathogen infecting humans and plants

Logged in as:
Joana Mourao

Author: Casandra Hernández-Reyes, Adam Schikora

LOGOUT

Publication: FEMS Microbiology Letters

Publisher: Oxford University Press

Date: 2013-06-01

Copyright © 2013, Oxford University Press

Order Completed

Thank you very much for your order.

This is a License Agreement between Joana Mourao ("You") and Oxford University Press ("Oxford University Press"). The license consists of your order details, the terms and conditions provided by Oxford University Press, and the [payment terms and conditions](#).

[Get the printable license.](#)

License Number	3765990379750
License date	Dec 11, 2015
Licensed content publisher	Oxford University Press
Licensed content publication	FEMS Microbiology Letters
Licensed content title	<i>Salmonella</i> , a cross-kingdom pathogen infecting humans and plants
Licensed content author	Casandra Hernández-Reyes, Adam Schikora
Licensed content date	2013-06-01
Volume number	343
Issue number	1
Type of Use	Thesis/Dissertation
Requestor type	Academic/Educational institute
Format	Print and electronic
Portion	Figure/table
Number of figures/tables	1
Will you be translating?	No
Author of this OUP article	No
Order reference number	None
Title of your thesis / dissertation	CONTRIBUTION OF BIOCIDES TO THE SELECTION OF SALMONELLA NON-TYPHOID RESISTANT TO ANTIBIOTICS: A MULTILAYERED APPROACH
Expected completion date	Feb 2016
Estimated size(pages)	250
Publisher VAT ID	GB 125 5067 30
Total	0.00 EUR

ORDER MORE...

CLOSE WINDOW

Copyright © 2015 [Copyright Clearance Center, Inc.](#) All Rights Reserved. [Privacy statement.](#) [Terms and Conditions.](#)
Comments? We would like to hear from you. E-mail us at customercare@copyright.com

14/12/2015

Rightslink® by Copyright Clearance Center



RightsLink®

Home

Account
Info

Help



Title: What is a resistance gene?
Ranking risk in resistomes

Author: José L. Martínez, Teresa M.
Coque, Fernando Baquero

Publication: Nature Reviews Microbiology

Publisher: Nature Publishing Group

Date: Dec 15, 2014

Logged in as:
Joana Mourao
Account #:
3000982054

LOGOUT

Copyright © 2014, Rights Managed by Nature
Publishing Group

Order Completed

Thank you very much for your order.

This is a License Agreement between Joana Mourao ("You") and Nature Publishing Group ("Nature Publishing Group"). The license consists of your order details, the terms and conditions provided by Nature Publishing Group, and the [payment terms and conditions](#).

[Get the printable license](#).

License Number	3767860797824
License date	Dec 14, 2015
Licensed content publisher	Nature Publishing Group
Licensed content publication	Nature Reviews Microbiology
Licensed content title	What is a resistance gene? Ranking risk in resistomes
Licensed content author	José L. Martínez, Teresa M. Coque, Fernando Baquero
Licensed content date	Dec 15, 2014
Type of Use	reuse in a dissertation / thesis
Volume number	13
Issue number	2
Requestor type	academic/educational
Format	print and electronic
Portion	figures/tables/illustrations
Number of figures/tables/illustrations	1
High-res required	no
Figures	Figure 1 Determination of epidemiological breakpoints of susceptibility to antibiotics
Author of this NPG article	no
Your reference number	None
Title of your thesis / dissertation	CONTRIBUTION OF BIOCIDES TO THE SELECTION OF SALMONELLA NON-TYPHOID RESISTANT TO ANTIBIOTICS: A MULTILAYERED APPROACH
Expected completion date	Feb 2016
Estimated size (number of pages)	250
Total	0.00 EUR

ORDER MORE...

CLOSE WINDOW

Copyright © 2015 [Copyright Clearance Center, Inc.](#) All Rights Reserved. [Privacy statement](#). [Terms and Conditions](#).
Comments? We would like to hear from you. E-mail us at customercare@copyright.com

17/12/2015

Rightslink® by Copyright Clearance Center



RightsLink®

Home

Account
Info

Help



Title: Platforms for antibiotic discovery
Author: Kim Lewis
Publication: Nature Reviews Drug Discovery
Publisher: Nature Publishing Group
Date: Apr 30, 2013

Logged in as:
 Joana Mourao
 Account #:
 3000982054

LOGOUT

Copyright © 2013, Rights Managed by Nature
 Publishing Group

Order Completed

Thank you very much for your order.

This is a License Agreement between Joana Mourao ("You") and Nature Publishing Group ("Nature Publishing Group"). The license consists of your order details, the terms and conditions provided by Nature Publishing Group, and the [payment terms and conditions](#).

[Get the printable license.](#)

License Number	3771550743194
License date	Dec 17, 2015
Licensed content publisher	Nature Publishing Group
Licensed content publication	Nature Reviews Drug Discovery
Licensed content title	Platforms for antibiotic discovery
Licensed content author	Kim Lewis
Licensed content date	Apr 30, 2013
Type of Use	reuse in a dissertation / thesis
Volume number	12
Issue number	5
Requestor type	academic/educational
Format	print and electronic
Portion	figures/tables/illustrations
Number of figures/tables/illustrations	2
High-res required	no
Figures	Figure1 Targets of antibiotics Figure of Box1 Antibiotic resistance and tolerance
Author of this NPG article	no
Your reference number	None
Title of your thesis / dissertation	CONTRIBUTION OF BIOCIDES TO THE SELECTION OF SALMONELLA NON-TYPHOID RESISTANT TO ANTIBIOTICS: A MULTILAYERED APPROACH
Expected completion date	Feb 2016
Estimated size (number of pages)	250
Total	0.00 EUR

ORDER MORE...

CLOSE WINDOW

Copyright © 2015 Copyright Clearance Center, Inc. All Rights Reserved. [Privacy statement](#). [Terms and Conditions](#).
 Comments? We would like to hear from you. E-mail us at customercare@copyright.com

17/12/2015

Rightslink® by Copyright Clearance Center



RightsLink®

Home

Account
Info

Help



Title: How antibiotics kill bacteria:
from targets to networks

Author: Michael A. Kohanski, Daniel J.
Dwyer and James J. Collins

Publication: Nature Reviews Microbiology

Publisher: Nature Publishing Group

Date: May 4, 2010

Logged in as:
Joana Mourao

Account #:
3000982054

LOGOUT

Copyright © 2010, Rights Managed by Nature
Publishing Group

Order Completed

Thank you very much for your order.

This is a License Agreement between Joana Mourao ("You") and Nature Publishing Group ("Nature Publishing Group"). The license consists of your order details, the terms and conditions provided by Nature Publishing Group, and the [payment terms and conditions](#).

[Get the printable license](#).

License Number	3771551002205
License date	Dec 17, 2015
Licensed content publisher	Nature Publishing Group
Licensed content publication	Nature Reviews Microbiology
Licensed content title	How antibiotics kill bacteria: from targets to networks
Licensed content author	Michael A. Kohanski, Daniel J. Dwyer and James J. Collins
Licensed content date	May 4, 2010
Type of Use	reuse in a dissertation / thesis
Volume number	8
Issue number	6
Requestor type	academic/educational
Format	print and electronic
Portion	figures/tables/illustrations
Number of figures/tables/illustrations	1
High-res required	no
Figures	Figure 1 Drug-target interactions and associated cell death mechanisms
Author of this NPG article	no
Your reference number	None
Title of your thesis / dissertation	CONTRIBUTION OF BIOCIDES TO THE SELECTION OF SALMONELLA NON-TYPHOID RESISTANT TO ANTIBIOTICS: A MULTILAYERED APPROACH
Expected completion date	Feb 2016
Estimated size (number of pages)	250
Total	0.00 EUR

ORDER MORE...

CLOSE WINDOW

Copyright © 2015 [Copyright Clearance Center, Inc.](#) All Rights Reserved. [Privacy statement](#). [Terms and Conditions](#).
Comments? We would like to hear from you. E-mail us at customer@copyright.com

21/12/2015

Rightslink® by Copyright Clearance Center



RightsLink®

Home

Account
Info

Help



Title: Mobile genetic elements: the agents of open source evolution

Author: Laura S. Frost, Raphael Leplae, Anne O. Summers and Ariane Toussaint

Logged in as:
Joana Mourao
Account #:
3000982054

LOGOUT

Publication: Nature Reviews Microbiology**Publisher:** Nature Publishing Group**Date:** Sep 1, 2005

Copyright © 2005, Rights Managed by Nature Publishing Group

Order Completed

Thank you very much for your order.

This is a License Agreement between Joana Mourao ("You") and Nature Publishing Group ("Nature Publishing Group"). The license consists of your order details, the terms and conditions provided by Nature Publishing Group, and the [payment terms and conditions](#).

[Get the printable license.](#)

License Number	3773400084931
License date	Dec 20, 2015
Licensed content publisher	Nature Publishing Group
Licensed content publication	Nature Reviews Microbiology
Licensed content title	Mobile genetic elements: the agents of open source evolution
Licensed content author	Laura S. Frost, Raphael Leplae, Anne O. Summers and Ariane Toussaint
Licensed content date	Sep 1, 2005
Type of Use	reuse in a dissertation / thesis
Volume number	3
Issue number	9
Requestor type	academic/educational
Format	print and electronic
Portion	figures/tables/illustrations
Number of figures/tables/illustrations	1
High-res required	no
Figures	Figure 1 Transfer of DNA between bacterial cells
Author of this NPG article	no
Your reference number	None
Title of your thesis / dissertation	CONTRIBUTION OF BIOCIDES TO THE SELECTION OF SALMONELLA NON-TYPHOID RESISTANT TO ANTIBIOTICS: A MULTILAYERED APPROACH
Expected completion date	Feb 2016
Estimated size (number of pages)	250
Total	0.00 EUR

ORDER MORE...

CLOSE WINDOW

Copyright © 2015 [Copyright Clearance Center, Inc.](#) All Rights Reserved. [Privacy statement](#). [Terms and Conditions](#).
Comments? We would like to hear from you. E-mail us at customercare@copyright.com

29/12/2015

Rightslink® by Copyright Clearance Center



RightsLink®

Home

Account
Info

Help



Title: Antimicrobial Resistance:
Implications for the Food
System

Publication: Comprehensive Reviews in Food
Science and Food Safety

Publisher: John Wiley and Sons

Date: Aug 2, 2006

Copyright © 2006, John Wiley and Sons

Logged in as:
Joana Mourao
Account #:
3000982054

LOGOUT

Order Completed

Thank you for your order.

This Agreement between Joana Mourao ("You") and John Wiley and Sons ("John Wiley and Sons") consists of your license details and the terms and conditions provided by John Wiley and Sons and Copyright Clearance Center.

Your confirmation email will contain your order number for future reference.

[Get the printable license.](#)

License Number	3777940679241
License date	Dec 28, 2015
Licensed Content Publisher	John Wiley and Sons
Licensed Content Publication	Comprehensive Reviews in Food Science and Food Safety
Licensed Content Title	Antimicrobial Resistance: Implications for the Food System
Licensed Content Author	None
Licensed Content Date	Aug 2, 2006
Licensed Content Pages	67
Type of use	Dissertation/Thesis
Requestor type	University/Academic
Format	Print and electronic
Portion	Figure/table
Number of figures/tables	1
Original Wiley figure/table number(s)	Figure 2 - Application of antimicrobials from farm to table
Will you be translating?	No
Title of your thesis / dissertation	CONTRIBUTION OF BIOCIDES TO THE SELECTION OF SALMONELLA NON-TYPHOID RESISTANT TO ANTIBIOTICS: A MULTILAYERED APPROACH
Expected completion date	Feb 2016
Expected size (number of pages)	250
Requestor Location	Joana Mourao

Billing Type
Billing address

Total 0.00 EUR

Would you like to purchase the full text of this article? If so, please continue on to the content ordering system located here: [Purchase PDF](#)

<https://s100.copyright.com/AppDispatchServlet>

1/2

30/12/2015

Rightslink® by Copyright Clearance Center



RightsLink®

Home

Account
Info

Help



Title: Nutritional immunity: transition metals at the pathogen–host interface

Author: M. Indriati Hood and Eric P. Skaar

Publication: Nature Reviews Microbiology

Publisher: Nature Publishing Group

Date: Jul 16, 2012

Copyright © 2012, Rights Managed by Nature Publishing Group

Logged in as:

Joana Mourao

Account #:

3000982054

LOGOUT

Order Completed

Thank you very much for your order.

This is a License Agreement between Joana Mourao ("You") and Nature Publishing Group ("Nature Publishing Group"). The license consists of your order details, the terms and conditions provided by Nature Publishing Group, and the [payment terms and conditions](#).

[Get the printable license.](#)

License Number	3778511098572
License date	Dec 29, 2015
Licensed content publisher	Nature Publishing Group
Licensed content publication	Nature Reviews Microbiology
Licensed content title	Nutritional immunity: transition metals at the pathogen–host interface
Licensed content author	M. Indriati Hood and Eric P. Skaar
Licensed content date	Jul 16, 2012
Type of Use	reuse in a dissertation / thesis
Volume number	10
Issue number	8
Requestor type	academic/educational
Format	print and electronic
Portion	figures/tables/illustrations
Number of figures/tables/illustrations	1
High-res required	no
Figures	Figure 5 New insights into the roles for Cu+ in innate immunity
Author of this NPG article	no
Your reference number	None
Title of your thesis / dissertation	CONTRIBUTION OF BIOCIDES TO THE SELECTION OF SALMONELLA NON-TYPHOID RESISTANT TO ANTIBIOTICS: A MULTILAYERED APPROACH
Expected completion date	Feb 2016
Estimated size (number of pages)	250
Total	0.00 EUR

ORDER MORE...

CLOSE WINDOW

Copyright © 2015 [Copyright Clearance Center, Inc.](#) All Rights Reserved. [Privacy statement](#). [Terms and Conditions](#).
Comments? We would like to hear from you. E-mail us at customercare@copyright.com

23/01/2016

Rightslink® by Copyright Clearance Center



RightsLink®

Home

Account
Info

Help



Title: Antimicrobial activity of metals: mechanisms, molecular targets and applications

Author: Joseph A. Lemire, Joe J. Harrison, Raymond J. Turner

Publication: Nature Reviews Microbiology

Publisher: Nature Publishing Group

Date: May 13, 2013

Copyright © 2013, Rights Managed by Nature Publishing Group

Logged in as:

Joana Mourao

Account #:

3000982054

[LOGOUT](#)

Order Completed

Thank you very much for your order.

This is a License Agreement between Joana Mourao ("You") and Nature Publishing Group ("Nature Publishing Group"). The license consists of your order details, the terms and conditions provided by Nature Publishing Group, and the [payment terms and conditions](#).

[Get the printable license.](#)

License Number	3794541385005
License date	Jan 22, 2016
Licensed content publisher	Nature Publishing Group
Licensed content publication	Nature Reviews Microbiology
Licensed content title	Antimicrobial activity of metals: mechanisms, molecular targets and applications
Licensed content author	Joseph A. Lemire, Joe J. Harrison, Raymond J. Turner
Licensed content date	May 13, 2013
Type of Use	reuse in a dissertation / thesis
Volume number	11
Issue number	6
Requestor type	academic/educational
Format	print and electronic
Portion	figures/tables/illustrations
Number of figures/tables/illustrations	2
High-res required	no
Figures	Figure 2. Antibacterial mechanisms of metal toxicity. Figure of Box 2. Adapt and resist: how microorganisms deal with deadly metals.
Author of this NPG article	no
Your reference number	None
Title of your thesis / dissertation	CONTRIBUTION OF BIOCIDES TO THE SELECTION OF SALMONELLA NON-TYPHOID RESISTANT TO ANTIBIOTICS: A MULTILAYERED APPROACH
Expected completion date	Feb 2016
Estimated size (number of pages)	250
Total	0.00 EUR

[ORDER MORE...](#)
[CLOSE WINDOW](#)

Copyright © 2016 [Copyright Clearance Center, Inc.](#) All Rights Reserved. [Privacy statement](#). [Terms and Conditions](#).
Comments? We would like to hear from you. E-mail us at customer@copyright.com

09/01/2016

Rightslink® by Copyright Clearance Center



RightsLink®

Home

Account
Info

Help



Title: Survival in amoeba—a major selection pressure on the presence of bacterial copper and zinc resistance determinants? Identification of a “copper pathogenicity island”

Author: Xiuli Hao

Publication: Applied Microbiology and Biotechnology

Publisher: Springer

Date: Jan 1, 2015

Copyright © 2015, Springer-Verlag Berlin Heidelberg

Logged in as:

Joana Mourao

Account #:

3000982054

LOGOUT

Order Completed

Thank you very much for your order.

This is a License Agreement between Joana Mourao ("You") and Springer ("Springer"). The license consists of your order details, the terms and conditions provided by Springer, and the [payment terms and conditions](#).

[Get the printable license.](#)

License Number	3785010353411
License date	Jan 09, 2016
Licensed content publisher	Springer
Licensed content publication	Applied Microbiology and Biotechnology
Licensed content title	Survival in amoeba—a major selection pressure on the presence of bacterial copper and zinc resistance determinants? Identification of a “copper pathogenicity island”
Licensed content author	Xiuli Hao
Licensed content date	Jan 1, 2015
Volume number	99
Issue number	14
Type of Use	Thesis/Dissertation
Portion	Figures/tables/illustrations
Number of figures/tables/illustrations	1
Author of this Springer article	No
Original figure numbers	Fig. 2 Pco and Sil mechanisms in action.
Title of your thesis / dissertation	CONTRIBUTION OF BIOCIDES TO THE SELECTION OF SALMONELLA NON-TYPHOID RESISTANT TO ANTIBIOTICS: A MULTILAYERED APPROACH
Expected completion date	Feb 2016
Estimated size(pages)	250
Total	0.00 EUR

CLOSE WINDOW

Copyright © 2016 Copyright Clearance Center, Inc. All Rights Reserved. [Privacy statement](#). [Terms and Conditions](#). Comments? We would like to hear from you. E-mail us at customercare@copyright.com